

CATEGORICITY OF AN ABSTRACT
ELEMENTARY CLASS IN
TWO SUCCESSIVE CARDINALS
SH576

SAHARON SHELAH

Institute of Mathematics
The Hebrew University
Jerusalem, Israel

Rutgers University
Department of Mathematics
New Brunswick, NJ USA

ABSTRACT. We investigate categoricity of abstract elementary classes without any remnants of compactness (like non-definability of well ordering, existence of E.M. models, or existence of large cardinals). We prove (assuming a weak version of GCH around λ) that if $\mathfrak{K}$ is categorical in λ, λ^+ , $LS(\mathfrak{K}) \leq \lambda$ and $1 \leq I(\lambda^{++}, \mathfrak{K}) < 2^{\lambda^{++}}$ then $\mathfrak{K}$ has a model in λ^{+++} .

Partially supported by the United States-Israel Binational Science Foundation.

I thank Alice Leonhardt for the excellent typing.

Done 10/94, (set theoretic, pseudo descriptive set theory) 22-29/11/94 (essentially §1-§2); more 9-14/12/95

(fulfilling §1, §3 eliminating use of the maximal model in λ^{+2} , (new §4) of categoricity in λ^{+2} (see §7, §8), more 18-2 of 4/95 (redo §3, §10).

§10 written 1/2/95

First typed at Rutgers - 95/Feb/10

Latest Revision 98/May/15

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

ANNOTATED CONTENT

§0 Introduction

[We give three versions of the main theorem in 0.2 - 0.4. In 0.5 - 0.32 we review the relevant knowledge of abstract elementary classes to help make this paper self-contained. This includes the representation by PC-classes defined by omission of quantifier free types (0.13, 0.14); types and stability (based on $\leq_{\mathfrak{K}}$); and the equivalence of saturation to model homogeneity (0.26).]

§1 Weak Diamond

[We mainly present necessary material on the weak diamond, a combinatorial principle whose main variant holds for λ^+ if $2^\lambda < 2^{\lambda^+}$.

We state cases provable from ZFC together with suitable cardinal arithmetic assumptions (1.2). We present applications of weak diamond to the number of models of fixed cardinality (notably in 1.10). We deal also with the definable weak diamond, and introduce a smaller ideal of “small” sets: $\text{UDmId}^{\mathcal{F}}(\lambda)$ (1.14 - 1.17).]

§2 First Attempts

[We define the class K_λ^3 of triples (M, N, a) representing types in $\mathcal{S}(M)$ for $M \in K_\lambda$, and start to investigate it, dealing with the weak extension property, the extension property, minimality, reduced types (except for minimality, in the first order case, these hold trivially). Our aims are to have the extension property or at least the weak extension property for all triples in K_λ^3 , and the density of minimal triples. The first property makes the model theory more like the first order case, and the second is connected with categoricity. We start by proving the weak extension property under reasonable assumptions. We prove the density of minimal triples under the strong assumption $K_{\lambda^{++}} = \emptyset$ and an extra cardinal arithmetic assumption ($2^{\lambda^+} > \lambda^{++}$). In the end, under the additional assumption $K_{\lambda^{++}} = 0$ we prove that all triples have the extension property and that we have disjoint amalgamation in K_λ . Now the assumption $K_{\lambda^{++}} = 0$ does no harm if we just want to prove Theorem 0.2. The reader willing to accept these assumptions may skip some proofs later. The proof of the extension property makes essential use of categoricity in λ^+ .]

§3 Non-structure

[We try to present clearly and in some generality the construction of many models in λ^{++} based on knowledge of models of size λ^+ , using weak diamond on λ^+ and on λ^{++} . This is done by forming a tree $\langle \bar{M}^\eta : \eta \in {}^{\lambda^{++}}2 \rangle$ with $\bar{M}^\eta$ an $\leq_{\mathfrak{K}}$ -increasing continuous sequence of members of K_λ with limit $\bigcup_{i < \lambda^+} M_i^\eta$ increasing with η (and an additional restriction). Actually λ^+ can be replaced by a regular uncountable λ' (so $\|M_i^\eta\| = \lambda$ is replaced by $\|M_i^\eta\| < \lambda'$).

§4 Minimal types

[We prove that every member of K_λ^3 has the extension property, by proving it for minimal triples. We use: if $M_\ell \in K_\lambda$ and for every minimal $p \in \mathcal{S}(M_0)$ the set $\mathcal{S}_{\geq p}(M_1)$ has cardinality $\leq \lambda^+$, then the $M \in K_{\lambda^+}$ is saturated for minimal types and hence the number of minimal types in $\mathcal{S}(M_1)$ is $\leq \lambda^+$ (for $M_1 \in K_\lambda$), which is a step toward stability in λ .]

§5 Inevitable types and stability in λ

[We continue to “climb the ladder”, using the amount of structure we already have (and sometimes categoricity) to get more. We start by assuming there are minimal types, and show that some minimal types are inevitable, construct $p_i \in \mathcal{S}(N_i)$ minimal ($i \leq \lambda^+$) both strictly increasing continuous and with p_0, p_δ inevitable, and then as in the proof of the equivalence of saturativity and model homogeneity, we show N_δ is universal over N_0 . We can then deduce stability in λ , so the model in λ^+ is saturated. Then we note that we have disjoint amalgamation in K_λ .]

§6 A proof for $\aleph$ categorical in λ^{+2}

[We give a shortcut to proving the main theorem by using stronger assumptions. If $I(\lambda^{+2}, K) = 1$ and $I(\lambda^{+3}, K) = 0$ then in some triple $(M, N, a) \in K_{\lambda^+}$, a is essentially algebraic over M , i.e. this is a maximal triple. Now first assuming for some pair $M_0 \leq_{\aleph} M_2$ in K_λ we have unique amalgamation for every possible M_1 with $M_0 \leq_{\aleph} M_1 \in K_\lambda$ (and using stability), we get models in λ^{++} thus contradicting the existence of maximal triples. We then use the methods of §3 to prove there are enough cases of unique amalgamation.]

§7 Extensions and Conjugacy

[We investigate types. We prove that in $\mathcal{S}(N), N \in K_\lambda$ that reduced implies inevitable, and that non-algebraic extensions preserve the conjugacy classes (so solving the realize/ materialize problem). Hence if $\langle N_i : i < \alpha \rangle$ is $<_{\aleph}$ -increasing in K_λ and λ divides α then N_α is $(\lambda, \text{cf}(\alpha))$ -saturated over N_0 .]

§8 Uniqueness of Amalgamation in $\aleph_\lambda$

[We have by §6 only $M_0^* <_{\aleph} M_2^*$ in $\aleph_\lambda$ such that $M_0^* \leq_{\aleph} M_1 \Rightarrow M_0^*, M_1, M_2^*$ has unique (disjoint) amalgamation. Now if we have a $\leq_{\aleph}$ -increasing continuous sequence $\langle N_i : i \leq \alpha \rangle$ such that $(N_i, N_{i+1}) \cong (M_0^*, M_2^*)$, we can amalgamate N_0, M_1, N_α whenever $N_0 \leq_{\aleph} M_1$, step by step. So some uniqueness is preserved and N_α can be any $(\lambda, \text{cf}(\alpha))$ -saturated model over N_0 . When we require also saturativity of M_i and of the resulting model, we get a nonforking relation denoted $\text{NF}_{\lambda, \delta}$. We define the general nonforking relation NF_λ by closing $\text{NF}_{\lambda, \delta}$ downward. So we succeed to define a relation which should behave as a nice nonforking relation. But we have to work to prove that this relation satisfies the expected properties, first for the “saturated” version and then in the general case by a diagram chase.]

§9 Nice extensions in K_{λ^+}

[As we have a notion of “nonforking” amalgamation in K_λ , we can use it to build $\leq_{\aleph}$ -extensions $M_1 \in K_{\lambda^+}$ for any given $M_0 \in K_{\lambda^+}$. This defines

naturally a two-place relation $\leq_{\lambda+}^*$ on $K_{\lambda+}$; “being a nice $\leq_{\mathfrak{R}}$ -submodel”. We investigate it and variants. In particular, we prove the existence of disjoint amalgamation.]

§10 Non-structure for $\leq_{\lambda+}^*$

[Instead proving that all disjoint amalgamations in K_{λ} are nonforking ones, we prove that on $K_{\lambda+}$ the relation $\leq_{\lambda+}^*$ is the same as $\leq_{\lambda}^*$, which is just as good for our purpose. Toward this we assume a failure and get many pairwise non-isomorphic models in $K_{\lambda+2}$, contradicting an assumption of 0.2(2). But once we have that $\leq_{\mathfrak{R}}$ agrees on $K_{\lambda+}$ with $\leq_{\lambda+}^*$ we have disjoint amalgamation, which suffices for building a model in $K_{\lambda+3}$.]

§0 INTRODUCTION

Makowski [Mw85] is a readable and good exposition concerning categoricity in abstract elementary classes around $\aleph_1$.

Our primary concern is:

Problem 0.1. Can we have some (not necessarily much) classification theory for reasonable non-first order classes $\mathfrak{K}$ of models, with no uses of even traces of compactness and only mild set theoretic assumptions?

Let me try to clarify the meaning of Problem 0.1.

What is the meaning of “mild set theoretic assumptions?” We are allowing requirements on cardinal arithmetic like GCH and weaker relatives. Preferably, assumptions like diamonds and squares and even mild large cardinals will not be used (apart from cases provable in ZFC, or in ZFC plus allowable assumptions).

In fact we try to continue [Sh 88], where results about the number of non-isomorphic models in $\aleph_1$ and $\aleph_2$ of a sentence $\psi \in L_{\omega_1, \omega}$ are obtained. Now in [Sh 88] the theorem parallel to the present one is proved assuming $2^{\aleph_0} < 2^{\aleph_1}$, so it is quite natural to use such assumptions here.

What is the meaning of “some classification theory?” While the dream is to have a classification theory as “full” as the one obtained in [Sh:c], we will be glad to have theorems speaking just on having few models in some cardinals or even categoricity and at least one in others. E.g. by [Sh 88] if $\psi \in L_{\omega_1, \omega}$ satisfies $1 \leq I(\aleph_1, \psi) < 2^{\aleph_1}$ (and $2^{\aleph_0} < 2^{\aleph_1}$) then $I(\aleph_2, \psi) > 0$. Here $I(\mu, \mathfrak{K})$ is the number of models in $\mathfrak{K}$ of cardinality μ , up to isomorphism.

What are “reasonable non first order classes?” This means we allow classes of “locally finite” or “atomic” structures, or structures “omitting a type”, or more generally the class of models of a sentence in $L_{\kappa, \omega}$, (i.e. allowing conjunction $< \kappa$ but quantification only over a finite string) but not one restricting ourselves to e.g. well orderings. In fact, we use “abstract elementary classes” from [Sh 88] (reviewed below).

What is the meaning of “uses traces of compactness?” For non first order classes we cannot use the powerful compactness theorem, but there are many ways to get weak forms of it: one way is using large cardinals (compact cardinals in Makkai Shelah [MaSh 285], or just measurable cardinals as in Kolman Shelah [KlSh 362], or in [Sh 472]). Another way is to use “non-definability of well ordering” which follows from the existence of Ehrenfeucht-Mostowski models, and also from $\psi \in L_{\omega_1, \omega}$ having uncountable models (used extensively in [Sh 88]). Our aim is to use none of this and we would like to see if any theory is left.

Above all, we hope the proofs will initiate classification theory in this case, so we hope the flavour will be one of introducing and investigating notions of a model theoretic character. Proofs of, say, a descriptive set theory character, will not satisfy this hope.

It seems to us that this goal is met here. We prove:

Theorem 0.2. $(2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}})$. Let $\mathfrak{K}$ be an abstract elementary class. If $\mathfrak{K}$ categorical in λ, λ^+ and λ^{++} then $I(\lambda^{++}, \mathfrak{K}) > 0$.

Here $\lambda^{++} = \lambda^{+++}$ and in general $\aleph_\alpha^{+\beta}$ means $\aleph_{\alpha+\beta}$.

Of course, the categoricity in three successive cardinals is a strong assumption.

Now note that in [Sh 88], the categoricity in $\aleph_0$ is gained “freely”, so the gap is smaller than seems at first glance. Still it is better to have

Theorem 0.3. $(2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}})$. Let $\mathfrak{K}$ be an abstract elementary class. If $\mathfrak{K}$ is categorical in λ and λ^+ , and $1 \leq I(\lambda^{++}, \mathfrak{K}) < 2^{\lambda^{++}}$ then $I(\lambda^{+3}, \mathfrak{K}) > 0$. Here we do not fully prove 0.3 but we prove a slightly weaker version:

Theorem 0.4. If $\mathfrak{K}$ is categorical in λ and λ^+ with $\lambda > \aleph_0$ and $1 \leq I(\lambda^{++}, \mathfrak{K}) < \mu_{\text{wd}}(\lambda^{++})$, then $I(\lambda^{+3}, \mathfrak{K}) > 0$.

Note however:

- (α) A silly point; at exactly one point in the proof of 0.4 we assume $\lambda > \aleph_0$ (in the proof of 4.6). This is silly as our intent is to prove for general λ what we know for $\lambda = \aleph_0$ by [Sh 88]; however, there we assume $\mathfrak{K}$ is $\text{PC}_{\aleph_1, \aleph_0}$, a reasonable assumption, but one which is not assumed here. We shall complete this in [Sh 600, §2], so we do not mention the assumption $\lambda > \aleph_0$ in theorems relying on 4.6.
- (β) More seriously, at some point we assume toward a contradiction that $K_{\lambda^{+3}} = \emptyset$. This is fine for proving theorem 0.2, but is not desirable if we want to develop a classification theory. This will also be dealt with in [Sh 600, §2].
- (γ) Concerning $\mu_{\text{wd}}(\lambda^{+2})$: in 6.12 we get $I(\lambda^{++}, K) \geq \mu_{\text{wd}}(\lambda^{++})$ instead of $I(\lambda^{++}, K) = 2^{\lambda^{++}}$ (eliminated in [Sh 600], too). The cardinal $\mu_{\text{wd}}(\chi)$ is defined in 1.1(6). It is almost 2^χ ; i.e. always $2^\chi < \mu_{\text{wd}}(\chi^+) \leq 2^{\chi^+} \leq \mu_{\text{wd}}(\chi^+)^{\aleph_0}$. Already in [Sh 87b] sometimes instead of getting 2^χ we get only $\mu_{\text{wd}}(\chi)$ (again see [Sh 600]). If you are satisfied with $I(\chi, K) \geq \mu_{\text{wd}}(\chi)$ in those cases, you can ignore some proofs.

We present below, as background, the following open questions which appeared in [Sh 88], for $\mathfrak{K}$ an abstract elementary class, of course, e.g. the class of models of $\psi \in L_{\lambda^+, \omega}$ with the relation $M \leq_{\mathfrak{K}} N$ being $M \prec_{\mathcal{L}} N$ for $\mathcal{L}$ a fragment of $L_{\kappa^+, \omega}$ to which ψ belongs. In [Sh 87a], [Sh 87b], [Sh 88] we prove:

- (*)₃ categoricity (of $\psi \in L_{\omega_1, \omega}(Q)$) in $\aleph_1$ implies the existence of a model of ψ of cardinality $\aleph_2$;
- (*)₄ if $n > 0$, $2^{\aleph_0} < 2^{\aleph_1} < \dots < 2^{\aleph_n}$, $\psi \in L_{\omega_1, \omega}$ and $1 \leq I(\aleph_\ell, \psi) < \mu_{\text{wd}}(\aleph_\ell)$ for $1 \leq \ell \leq n$, then ψ has a model of cardinality $\aleph_{n+1}$.

Now the problems were:

Problem 1 Prove (*)₃, (*)₄ in the context of an abstract elementary class $\mathfrak{K}$ which is $\text{PC}_{\aleph_0}$.

Problem 2 Parallel results in ZFC; e.g. prove (*)₃ when $n = 1$, $2^{\aleph_0} = 2^{\aleph_1}$.

Problem 3 Construct examples; e.g. $\mathfrak{K}$ (or $\psi \in L_{\omega_1, \omega}$), categorical in $\aleph_0, \aleph_1, \dots, \aleph_n$ but not in $\aleph_{n+1}$.

Problem 4 If $\mathfrak{K}$ is PC_λ (abstract elementary class), and is categorical in λ and λ^+ , does it necessarily have a model in λ^{++} ?

Concerning Problem 3, by Hart Shelah [HaSh 323], 2.10(2) + 3.8 there is $\psi_n \in \mathcal{L}_{\omega_1, \omega}$ categorical in $\aleph_0, \aleph_1, \dots, \aleph_{k-1}$, but not categorical in λ if $2^\lambda > 2^{\aleph_{k-1}}$.

The direct motivation for the present work, is that Grossberg asked me (Oct. 94) some questions in this neighborhood, in particular:

Problem 5) Assume $K = \text{Mod}(T)$, (i.e. K is the class of models of T), $T \subseteq L_{\omega_1, \omega}$, $|T| = \lambda$, $I(\lambda, K) = 1$ and $1 \leq I(\lambda^+, K) < 2^{\lambda^+}$. Does it follow that $I(\lambda^{++}, K) > 0$?

We think of this as a test problem and much prefer a model theoretic to a set theoretic solution. This is closely related to Problem 4 above and to [Sh 88, Theorem 3.7] (where we assume categoricity in λ^+ , do not require $2^\lambda < 2^{\lambda^+}$ but take $\lambda = \aleph_0$ or some similar cases) and [Sh 88, Theorem 5.17(4)] (and see [Sh 88, 5.1.4.5] on the assumptions) (there we require $2^\lambda < 2^{\lambda^+}$, $1 \leq I(\lambda^+, K) < 2^{\lambda^+}$ and $\lambda = \aleph_0$).

As said above, we are dealing with a closely related problem. Problem 0.1 was stated *a posteriori* but is, I think, the real problem.

In a first try we used more set theory, i.e. we used the definitional weak diamond on both λ^+ and λ^{++} (see Definition 1.13) and things like “a nice equivalence relation on $\mathcal{P}(\lambda)$ has either few or many classes” (see §1).

Here we take a model theoretic approach.

We feel that this paper provides a reasonable positive solution to problem 0.1, with a classification theory flavor. We shall continue in [Sh 600] toward a parallel of [Sh 87a], [Sh 87b]. Grossberg and Shelah, in the mid-eighties, started to write a paper [GrSh 266] to prove that: if $\psi \in L_{\lambda^+, \omega}$ has models of arbitrarily large cardinality, and is categorical in μ^{+n} for each n and if $\mu \geq \lambda$ and $2^{\mu^{+n}} < 2^{\mu^{+n+1}}$ for $n < \omega$, then ψ is categorical in every $\mu' \geq \mu$; this is a weak form of the upward part of Los’ conjecture. See Makkai Shelah [MaSh 285] on $T \subseteq \mathcal{L}_{\kappa, \omega}$ where κ is a compact cardinal; where we get downward and upward theorems for successor cardinals which are sufficiently bigger than $\kappa + |T|$. On the downward part, see [KlSh 362], [Sh 472] which deals with a downward theorem for successor cardinals which are sufficiently larger than $\kappa + |T|$ when the theory T is in the logic $\mathcal{L}_{\kappa, \omega}$ and κ is measurable. See also [Sh 394] which deals with abstract elementary classes with amalgamation, getting similar results with no large cardinals.

Part of §1 and §3 have a combinatorial character. Most of the paper forms the content of a course given in Fall ’94 (essentially without §3, §9, §10). The paper is written with an eye to developing the model theory, rather than just proving Theorem 0.2.

* * *

On abstract elementary classes see [Sh 88] and [Sh 300, II, §3]. To make the paper self-contained, we will review some relevant definitions and results. We thank Gregory Cherlin for improving the writing of §1-§3 and for breaking 3.25, 3.26 and 3.27 to three proofs.

This work is continued in [Sh 600].

ABSTRACT ELEMENTARY CLASSES

0.5 Conventions. $\mathfrak{K} = (K, \leq_{\mathfrak{K}})$, where K is a class of τ -models for a fixed vocabulary $\tau = \tau_K = \tau_{\mathfrak{K}}$ and $\leq_{\mathfrak{K}}$ is a two-place relation on the models in K . We do not always strictly distinguish between K and $(K, \leq_{\mathfrak{K}})$. We shall assume that $K, \leq_{\mathfrak{K}}$ are fixed, and $M \leq_{\mathfrak{K}} N \Rightarrow M, N \in K$; and we assume that the following axioms hold. When we use $<$ in the sense of elementary submodel for first order logic, we write $<_{\mathcal{L}_{\omega, \omega}}$.

0.6 Definition.

Ax 0: The validity of $M \in K$ or of $N \leq_{\mathfrak{K}} M$ depends on N and M only up to isomorphism - in the second case, isomorphism of the pair.

Ax I: If $M \leq_{\mathfrak{K}} N$ then $M \subseteq N$ (i.e. M is a submodel of N).

Ax II: $\leq_{\mathfrak{K}}$ is transitive and reflexive on K .

Ax III: If λ is a regular cardinal, M_i ($i < \lambda$) is $\leq_{\mathfrak{K}}$ -increasing (i.e. $i < j < \lambda$ implies $M_i \leq_{\mathfrak{K}} M_j$) and continuous (i.e. for limit ordinals $\delta < \lambda$ we have

$$M_\delta = \bigcup_{i < \delta} M_i) \text{ then } M_0 \leq_{\mathfrak{K}} \bigcup_{i < \lambda} M_i.$$

Ax IV: If λ is a regular cardinal, M_i ($i < \lambda$) is $\leq_{\mathfrak{K}}$ -increasing continuous, $M_i \leq_{\mathfrak{K}} N$ then $\bigcup_{i < \lambda} M_i \leq_{\mathfrak{K}} N$.

Ax V: If $M_0 \subseteq M_1$ and $M_\ell \leq_{\mathfrak{K}} N$ for $\ell = 0, 1$, then $M_0 \leq_{\mathfrak{K}} M_1$.

Ax VI: There is a cardinal λ such that: if $A \subseteq N$ and $|A| \leq \lambda$ then for some $M \leq_{\mathfrak{K}} N$ we have $A \subseteq |M| \leq \lambda$. We define the Löwenheim-Skolem number $LS(\mathfrak{K})$ as the least such λ with $\lambda \geq |\tau|$.

Notation: $K_\lambda = \{M \in K : \|M\| = \lambda\}$ and $K_{<\lambda} = \bigcup_{\mu < \lambda} K_\mu$.

$\mathcal{L}_{\omega, \omega}$ is first order logic.

A theory in $\mathcal{L}(\tau)$ is a set of sentences from $\mathcal{L}(\tau)$.

0.7 Definition. The embedding $f : N \rightarrow M$ is a $\mathfrak{K}$ -embedding or a $\leq_{\mathfrak{K}}$ -embedding if its range is the universe of a model $N' \leq_{\mathfrak{K}} M$, (so $f : N \rightarrow N'$ is an isomorphism (onto)).

Very central in [Sh 88], but peripheral here is:

0.8 Definition. 1) For a logic $\mathcal{L}$ and vocabulary τ , $\mathcal{L}(\tau)$ is the set of $\mathcal{L}$ -formulas in this vocabulary.

2) Let T_1 be a theory in $\mathcal{L}_{\omega, \omega}(\tau_1)$, $\tau \subseteq \tau_1$ vocabularies, Γ a set of types in $\mathcal{L}_{\omega, \omega}(\tau_1)$; (i.e. for some m , a set of formulas $\varphi(x_0, \dots, x_{m-1}) \in \mathcal{L}_{\omega, \omega}(\tau_1)$).

3) $EC(T_1, \Gamma) = \{M : M \text{ a } \tau_1\text{-model of } T_1 \text{ which omits every } p \in \Gamma\}$.

4) $PC_\tau(T_1, \Gamma) = PC(T_1, \Gamma, \tau) = \{M : M \text{ is a } \tau\text{-reduct of some } M_1 \in EC(T_1, \Gamma)\}$.

5) We say that $\mathfrak{K}$ is PC_λ^μ if for some $T_1, T_2, \Gamma_1, \Gamma_2$ and τ_1 and τ_2 we have: T_ℓ is a first order theory in the vocabulary τ_ℓ , Γ_ℓ is a set of types and $K = PC(T_1, \Gamma_1, \tau_{\mathfrak{K}})$ and $\{(M, N) : M \leq_{\mathfrak{K}} N, M, N \in K\} = PC(T_2, \Gamma_2, \tau')$ where

$\tau' = \tau_1 \cup \{P\}$ (P a new one place predicate), $|T_\ell| \leq \lambda, |\Gamma_\ell| \leq \mu$ for $\ell = 1, 2$. If $\mu = \lambda$, we may omit it.

0.9 Example. If $\tau_1 = \tau, T_1, \Gamma$ are as above, and $(K, \leq_{\mathfrak{K}})$ are defined by $K =: EC(T_1, \Gamma)$, $\leq_{\mathfrak{K}} =: \prec_{\mathcal{L}_{\omega, \omega}}$, then the pair satisfies the Axioms from 1.2 and $LS(\mathfrak{K}) \leq |T_1| + \aleph_0$.

0.10 Example. ($V = L$).

Let $\text{cf}(\lambda) \geq \aleph_1, n < \omega$ then for some $\psi \in L_{\lambda^+, \omega}$ we have: ψ has no model of cardinality $\lambda^{+(n+1)}$, and is categorical in λ^{+n} (i.e. has one and only one model up to isomorphism).

Let $M^* = (L_{\lambda^{+n}}, \in, i)_{i \leq \lambda}$ and let ψ be

$$\bigwedge \left\{ \varphi : \varphi \text{ is a first order sentence which } M^* \text{ satisfies} \right\} \\ \wedge (\forall x) \left(x \in \lambda \equiv \bigvee_{i < \lambda} x = i \right).$$

0.11 Lemma. Let I be a directed set (i.e. partially ordered by $\leq$, such that any two elements have a common upper bound).

1) If M_t is defined for $t \in I$, and $t \leq s \in I$ implies $M_t \leq_{\mathfrak{K}} M_s$ then for every $t \in I$ we have $M_t \leq_{\mathfrak{K}} \bigcup_{s \in I} M_s$.

2) If in addition $t \in I$ implies $M_t \leq_{\mathfrak{K}} N$ then $\bigcup_{s \in I} M_s \leq_{\mathfrak{K}} N$.

Proof. By induction on $|I|$ (simultaneously for 1) and 2)), or see [Sh 88, 1.6]. $\square_{0.11}$

0.12 Lemma.

1) Let $\tau_1 = \tau \cup \{F_i^n : i < LS(\mathfrak{K}) \text{ and } n < \omega\}$, F_i^n an n -place function symbol (assuming, of course, $F_i^n \notin \tau$). If M_1 is an expansion of M to a τ_1 -model and $\bar{a} \in {}^n|M|$ for some n , let $M_{\bar{a}}$ be the submodel of M with universe $\{F_i^n(\bar{a}) : i < LS(k)\}$. Every model M from K can be expanded to a τ_1 -model M_1 such that:

- (A) $M_{\bar{a}} \leq_{\mathfrak{K}} M$ for any $\bar{a} \in {}^n|M|$;
- (B) if $\bar{a} \in {}^n|M|$ then $\|M_{\bar{a}}\| \leq LS(\mathfrak{K})$;
- (C) if $\bar{b}$ is a subsequence of $\bar{a}$ (even up to rearrangement), then $M_{\bar{b}} \leq_{\mathfrak{K}} M_{\bar{a}}$;
- (D) for every $N_1 \subseteq M_1$ we have $N_1 \upharpoonright \tau \leq_{\mathfrak{K}} M$.

2) If $N \leq_{\mathfrak{K}} M$ is given, then we can choose the expansion M_1 so that clauses (A) - (D) hold and

- (E) $N = N_1 \upharpoonright \tau$ for some $N_1 \subseteq M_1$.

Proof. We define by induction on n , the values of $F_i^n(\bar{a})$ for every $i < LS(\mathfrak{K})$, $\bar{a} \in {}^n|M|$ such that $\bar{a} \subseteq N \Rightarrow M_{\bar{a}} \subseteq N$ when we are proving (2). By $Ax VI$ there is an $M_{\bar{a}} \leq_{\mathfrak{K}} M$, $\|M_{\bar{a}}\| \leq LS(\mathfrak{K})$ such that $|M_{\bar{a}}|$ includes

$$\bigcup \{M_{\bar{b}} : \bar{b} \text{ a subsequence of } \bar{a} \text{ of length } < n\} \cup \bar{a}$$

and $M_{\bar{a}}$ does not depend on the order of $\bar{a}$. Let $|M_{\bar{a}}| = \{c_i : i < i_0 \leq LS(\mathfrak{K})\}$, and define $F_i^n(\bar{a}) = c_i$ for $i < i_0$ and $F_i^n(\bar{a}) = c_0$ for $i_0 \leq i < LS(\mathfrak{K})$ (so we can demand “ F_i^n is symmetric”).

Clearly our conditions are satisfied: if $\bar{b}$ is a subsequence of $\bar{a}$ then $M_{\bar{b}} \leq_{\mathfrak{K}} M_{\bar{a}}$ by $Ax V$. $\square_{0.12}$

0.13 Lemma. 1) *There is a set Γ of types in $\mathcal{L}_{\omega, \omega}(\tau_1)$ (where τ_1 is as in Lemma 0.12) such that $K = PC(\emptyset, \Gamma, \tau)$, so it is a $PC_{LS(\mathfrak{K})}^{2^{LS(\mathfrak{K})}}$ -class.*

The types above consist of quantifier-free formulas.

2) *Moreover, if $N_1 \subseteq M_1 \in EC(\emptyset, \Gamma)$, and N, M are the τ -reducts of N_1, M_1 respectively then $N \leq_{\mathfrak{K}} M$. Also, if $N \leq_{\mathfrak{K}} M$ then there is an τ_1 -expansion M_1 of M and a submodel N_1 of M_1 such that $M_1 \in EC(\emptyset, \Gamma)$ and $N_1 \upharpoonright \tau = N$.*

3) *Also $\{(M, N) : N \leq_{\mathfrak{K}} M\}$ is a $PC_{LS(\mathfrak{K})}^{(2^{LS(\mathfrak{K})})}$ -class, hence $\mathfrak{K}$ is as well.*

Proof. 1) Let Γ_n be the set of complete quantifier free n -types $p(x_0, \dots, x_{n-1})$ in $\mathcal{L}_{\omega, \omega}(\tau_1)$ such that: if M_1 is an L_1 -model, $\bar{a}$ realizes p in M_1 and M is the L -reduct of M_1 , then $M_{\bar{b}} \leq_{\mathfrak{K}} M_{\bar{a}}$ for any subsequence of $\bar{b}$ of $\bar{a}$. Where $M_{\bar{c}}$ (for $\bar{c} \in {}^m|M_1|$) is the submodel of M whose universe is $\{F_i^m(\bar{c}) : i < LS(\mathfrak{K})\}$ (and there are such submodels).

Let Γ be the set of p which are complete quantifier free n -types for some $n < \omega$ in $\mathcal{L}_{\omega, \omega}(\tau_1)$ and which do not belong to Γ_n . So if M^1 is in $PC(\emptyset, \Gamma, \tau_1)$ then by 0.11 we have $M_1 \upharpoonright \tau \in K$ hence $PC(\emptyset, \Gamma, L) \subseteq K$ and by 0.12(1) we have $K \subseteq PC(\emptyset, \Gamma, L)$.

2) The first phrase is proven as in (1). For the second phrase, use 0.12(2).

3) Follows from (2). $\square_{0.13}$

0.14 Conclusion. There is τ_1 with $\tau \subseteq \tau_1$ and $|\tau_1| \leq LS(\mathfrak{K})$ such that: for any $M \in K$ and any τ_1 -expansion M_1 of M which is in $EC(\emptyset, \Gamma)$,

$$N_1 \subseteq M_1 \Rightarrow N_1 \upharpoonright \tau \leq_{\mathfrak{K}} M$$

$$N_1 \subseteq N_2 \subseteq M_1 \Rightarrow N_1 \upharpoonright \tau \leq_{\mathfrak{K}} N_2 \upharpoonright \tau.$$

0.15 Conclusion. If for every $\alpha < (2^{LS(\mathfrak{K})})^+$, K has a model of cardinality $\geq \beth_\alpha$ then K has a model in every cardinality $\geq LS(\mathfrak{K})$.

Proof. Use 0.13 and the value of the Hanf number for: models of a first order theory omitting a given set of types, for languages of cardinality $LS(K)$ (see [Sh:c, VII, §5]).

$\square_{0.15}$

0.16 Definition. For λ regular $> LS(\mathfrak{K})$ and $N \in \mathfrak{K}_\lambda$ we say $\bar{N} = \langle N_\alpha : \alpha < \lambda \rangle$ is a representation of N if $\bar{N}$ is $\leq_{\mathfrak{K}}$ -increasing continuous, $\|N_\alpha\| < \lambda$ and $N = \bigcup_{\alpha < \lambda} N_\alpha$.

If $\lambda = \mu^+$ then if not said otherwise, we require $\|N_\alpha\| = \mu$.

How will we define types, and in particular, the set $\mathcal{S}(M)$ of complete types over M , when no formulas are present? If we have a “monster model” $\mathfrak{C}$ we can use automorphisms; but any such “monster” is far down the road. So we will “chase diagrams” in K_λ (being careful, not to use excessively large models). This gives us a relation of “having the same type” we call E_μ^{at} , but this relation in general is not transitive (if we do not have amalgamation in K_λ). So E_μ will be defined as the transitive closure of E_μ^{at} .

0.17 Definition. 1) The two-place relation E_M^{at} is defined on triples (M, N, a) with M fixed, $M \leq_{\mathfrak{K}} N \in K_{\|M\|}$, and $a \in N$ by:

$$(M, N_1, a_1) E_M^{at} (M, N_2, a_2) \text{ iff there is } N \in K_\mu \text{ and } \leq_{\mathfrak{K}}\text{-embeddings} \\ f_\ell : N_\ell \rightarrow N \text{ for } \ell = 1, 2 \text{ such that:} \\ f_1 \upharpoonright M = \text{id}_M = f_2 \upharpoonright M \text{ and } f_1(a_1) = f_2(a_2).$$

Let E_M be the transitive closure of E_M^{at} .

2) For $\mu \geq LS(\mathfrak{K})$ and $M \in K_\mu$ we define $\mathcal{S}(M)$ as $\{\text{tp}(a, M, N) : M \leq_{\mathfrak{K}} N \in K_\mu \text{ and } a \in N\}$ where $\text{tp}(a, M, N) = (M, N, a)/E_M$.

3) We say “ a realizes p in N ” if $a \in N$, $p \in \mathcal{S}(M)$ and for some $N' \in K_\mu$ we have $M \leq_{\mathfrak{K}} N' \leq_{\mathfrak{K}} N$, $a \in N'$, and $p = \text{tp}(a, M, N')$.

4) We say “ a_2 strongly realizes $(M, N^1, a^1)/E_M^{at}$ in N ” if for some N^2, a^2 we have $M \leq_{\mathfrak{K}} N^2 \leq_{\mathfrak{K}} N$, $a_2 \in N^2$, and $(M, N^1, a^1) E_M^{at} (M, N^2, a^2)$.

5) We say $M_0 \in \mathfrak{K}$ is an amalgamation base if letting $\lambda = \|M_0\|$ we have: for every $M_1, M_2 \in \mathfrak{K}_\lambda$ and $\leq_{\mathfrak{K}}$ -embeddings $f_\ell : M_0 \rightarrow M_\ell$ (for $\ell = 1, 2$) there is $M_3 \in \mathfrak{K}_\lambda$ and $\leq_{\mathfrak{K}}$ -embeddings $g_\ell : M_\ell \rightarrow M_3$ (for $\ell = 1, 2$) such that $g_1 \circ f_1 = g_2 \circ f_2$.

6) We say $\mathfrak{K}$ is stable in λ iff $LS(\mathfrak{K}) \leq \lambda$ and for all $M \in K_\lambda$ we have $|\mathcal{S}(M)| \leq \lambda$.

0.18 Observation. If M is an amalgamation base then $E_M = E_M^{at}$, and we have:

$$“a \text{ strongly realizes } (M, N, b)/E_M \text{ in } N” \text{ iff } “a \text{ realizes } (M, N, b)/E_M.”$$

0.19 Definition.

1) $\mathfrak{K}$ has the λ -amalgamation property or has amalgamation in λ , if every $M_0 \in \mathfrak{K}_\lambda$ is an amalgamation base (see 0.17(5) above).

2) N is universal over M if: $M \leq_{\mathfrak{K}} N$ and if $M \leq_{\mathfrak{K}} N' \in K_{\leq \|N\|}$ then N' can be $\leq_{\mathfrak{K}}$ -embedded into N over M .

3) $\mathfrak{K}$ has universal extensions in λ iff for every $M \in K_\lambda$ there is N such that:

- (a) $M \leq_{\mathfrak{K}} N \in K_\lambda$
- (b) N is universal over M .

4) N_1, N_2 have a joint embedding if for some $N \in K$ there are $\leq_{\mathfrak{K}}$ -embeddings h_ℓ of N_ℓ into N for $\ell = 1, 2$. Let JEP_μ (JEP) means this holds for N_1, N_2 in K_μ (in K).

5) $\mathfrak{K}_\lambda$ has unique amalgamation (or $\mathfrak{K}$ has unique amalgamation in λ) when: if $M_i^\ell \in K_\lambda$ for $\ell < 2, i < 4$, and for $i = 1, 2$ we have $h_{i,0}^\ell$ is a $\leq_{\mathfrak{K}}$ -embedding of M_0^ℓ into M_i^ℓ , $h_{3,i}^\ell$ is a $\leq_{\mathfrak{K}}$ -embedding of M_i^ℓ into M_3^ℓ , $h_{3,1}^\ell \circ h_{1,0}^\ell = h_{3,2}^\ell \circ h_{2,0}^\ell$ and $\text{Rang}(h_{3,1}^\ell) \cap \text{Rang}(h_{3,2}^\ell) = \text{Rang}(h_{3,1}^\ell \circ h_{1,0}^\ell)$ and for $i < 3, f_i$ is an isomorphism from M_i^0 onto $M_i^1, f_0 \subseteq f_1, f_0 \subseteq f_2$ then for some $N \in K_\lambda$ there are $\mathfrak{K}$ -embedding $h_\ell : M_3^\ell \rightarrow N$ such that $h_0 \circ h_{3,i}^0 = h_1 \circ h_{3,i}^1$, for $i = 1, 2$.

6) Let $p_\ell \in \mathcal{S}(M_\ell)$ for $\ell = 0, 1$. We say $p_0 \leq p_1$ iff $M_0 \leq_{\mathfrak{K}} M_1$ and for some N and a we have $M_1 \leq_{\mathfrak{K}} N \in K_{\|M_1\|+LS(\mathfrak{K})}, a \in N$ and $\text{tp}(a, M_\ell, N) = p_\ell$ for $\ell = 0, 1$. We also write $p_0 = p_1 \upharpoonright M_0$ (p_0 is unique knowing M_0, M_1, p_1 hence $p_1 \upharpoonright M_0$ is well defined).

0.20 Claim.

1) If $\mathfrak{K}$ is categorical in λ and $LS(\mathfrak{K}) \leq \lambda$ then: there is a model in K_{λ^+} iff for some (equivalently, every) model $M \in K_\lambda$ there is N such that $M <_{\mathfrak{K}} N \in K_\lambda$ iff for some (equivalently every) $M \in K_\lambda$ there is N such that $M <_{\mathfrak{K}} N \in K_{\lambda^+}$.

2) If $\mathfrak{K}$ has amalgamation in $\lambda, LS(\mathfrak{K}) \leq \lambda$, and $M_0 \leq_{\mathfrak{K}} M_1$ are in K_λ with $M_0 \leq_{\mathfrak{K}} N_0 \in K_{\lambda^+}$ then we can find h and N_1 such that $N_0 \leq_{\mathfrak{K}} N_1 \in K_{\lambda^+}$ and h is a $\leq_{\mathfrak{K}}$ -embedding of M_1 into N_1 extending id_{M_0} . We can allow $N_0 \in K_\mu$ with $\mu > \lambda$ if $\mathfrak{K}$ has amalgamation in every $\lambda' \in [\lambda, \mu)$.

3) Assume $\mathfrak{K}$ has amalgamation in λ and $LS(\mathfrak{K}) \leq \lambda$. If $M_0 \leq M_1$ are from K_λ and $p_0 \in \mathcal{S}(M_0)$ then we can find an extension $p_1 \in \mathcal{S}(M_1)$ of p_0 .

Proof. 1) As then we can choose by induction on $i < \lambda^+$ models

$M_i \in K_\lambda, \leq_{\mathfrak{K}}$ -increasing continuous, $M_i \neq M_{i+1}$, for $i = 0$ use $K_{\lambda^+} \neq \emptyset$, for

i limit take union, for $i = j + 1$ use the previous sentence; so

$M_{\lambda^+} = \cup\{M_i : i < \lambda^+\} \in K_{\lambda^+}$ as required.

2), 3) Left to the reader. $\square_{0.20}$

0.21 Remark. We can here add the content of 6.5, 6.7, 6.12.

0.22 Definition. 1) For $\lambda > LS(\mathfrak{K})$ we say “ $N \in \mathfrak{K}$ is λ -saturated” iff for every $M \leq_{\mathfrak{K}} N$ of cardinality $< \lambda$, if $M \leq_{\mathfrak{K}} N'$ in $K_{<\lambda}$ and $a' \in N'$ then some $a \in N$ strongly realizes $(M, N', a')/E_M^{\text{at}}$ (in the interesting cases this suffices).

2) We say “ $N \in \mathfrak{K}$ is λ -saturated above μ (or is λ -saturated $\geq \mu$)” iff above we restrict ourselves to M of cardinality $\geq \mu$.

(Cf. (λ, κ) -saturated in Definition 0.28, 0.29 below).

0.23 Fact. 1) If $\mu + LS(\mathfrak{K}) < \lambda, \lambda$ is regular, $\mathfrak{K}$ has the amalgamation property in every $\mu' \in [\mu, \lambda)$, and for all $M \in K_{[\mu, \lambda)}$ we have $|\mathcal{S}(M)| \leq \lambda$ and $\lambda = \text{cf}(\lambda)$, then there is some $M \in K_\lambda$ saturated above μ .

2) Assume $\lambda > \mu \geq LS(\mathfrak{K})$ and $N \in K$ and $\mathfrak{K}$ has amalgamation in K_μ , for every $\mu_1 \in [\mu, \lambda)$. Then: N is λ -saturated above μ iff for every $M \leq_{\mathfrak{K}} N$ of cardinality $< \lambda$, every $p \in \mathcal{S}(M)$ is realized in N (i.e. for some $a \in N$ we have $\text{tp}(a, M, N) = p$).

3) If $LS(\mathfrak{K}) \leq \mu_0 \leq \mu'_0 \leq \mu' \leq \mu$ and M is μ -saturated above μ_0 , then it is μ' -saturated above μ'_0 . If $LS(\mathfrak{K}) \leq \mu_0 < \mu$ then: M is μ -saturated above μ_0 iff for every $\lambda \in [\mu_0, \mu)$, M is λ^+ -saturated above λ .

0.24 Definition. The type $p \in \mathcal{S}(M)$ is local when: if for any directed partial order I and models $M_t \leq_{\mathfrak{K}} M$ for $t \in I$ with $I \models t \leq s \Rightarrow M_t \leq_{\mathfrak{K}} M_s$ and $M = \bigcup_{t \in I} M_t$, and any $p' \in \mathcal{S}(M)$ with $(p \upharpoonright M_t = p' \upharpoonright M_t)$ for all $t \in I$, $p = p'$. We say M is local if every $p \in \mathcal{S}(M)$ is, and $\mathcal{K}$ is local if every $M \in \mathcal{K}$ is. We can add “above μ ” as in Definition 0.22(2).

0.25 Definition. 1) $I(\lambda, K) = I(\lambda, \mathfrak{K})$ is the number of $M \in K_\lambda$ up to isomorphism. 2) $\mathfrak{K}$ (or K) is categorical in λ if $I(\lambda, K) = 1$. 3) $IE(\lambda, \mathfrak{K}) = \sup\{|K'| : K' \subseteq K_\lambda \text{ and for } M \neq N \text{ in } K'_\lambda, M \text{ is not } \leq_{\mathfrak{K}}\text{-embeddable into } N\}$. If we write $IE(\lambda, K) \geq \mu$, we mean that for some $K' \subseteq K_\lambda$ as above, $|K'| \geq \mu$, and similarly for $= \mu$. If there is a problem with attainment of the supremum we shall say explicitly.

0.26 The Model-homogeneity = Saturativity Lemma. *Let $\lambda > \mu \geq LS(\mathfrak{K})$ and $M \in K$.*

- 1) M is λ -saturated above μ iff M is $(\mathcal{D}_{\mathfrak{K}}, \lambda)$ -homogeneous above μ , which means: for every $N_1 \leq_{\mathfrak{K}} N_2 \in K$ such that $\mu \leq \|N_1\| \leq \|N_2\| < \lambda$ and $N_1 \leq_{\mathfrak{K}} M$ there is a $\leq_{\mathfrak{K}}$ -embedding f of N_2 into M over N_1 .
- 2) If $M_1, M_2 \in K_\lambda$ are λ -saturated above $\mu < \lambda$ and for some $N_1 \leq_{\mathfrak{K}} M_1, N_2 \leq_{\mathfrak{K}} M_2$, both of cardinality $\in [\mu, \lambda)$, we have $N_1 \cong N_2$ then $M_1 \cong M_2$; in fact, any isomorphism f from N_1 onto N_2 can be extended to an isomorphism from M_1 onto M_2 .
- 3) If in (2) we demand only “ M_2 is λ -saturated” and $M_1 \in K_{\leq \lambda}$ then f can be extended to a $\leq_{\mathfrak{K}}$ -embedding from M_1 into M_2 .
- 4) In part (2) instead of $N_1 \cong N_2$ it suffices to assume that N_1 and N_2 can be $\leq_{\mathfrak{K}}$ -embedded into some $N \in K$, which holds if $\mathfrak{K}$ has the JEP.

Proof. 1) The “if” direction is easy as $\lambda > LS(\mathfrak{K})$. Let us prove the other direction, so without loss of generality $\|N_1\| = \mu$.

By 0.19(6) without loss of generality λ is regular, moreover N_2 has cardinality $\|N_1\|$.

Let $|N_2| = \{a_i : i < \kappa\}$, and we know $\kappa = \|N_1\| = \|N_2\| < \lambda$. We define by induction on $i \leq \kappa$, N_1^i, N_2^i, f_i such that:

- (a) $N_1^i \leq_{\mathfrak{K}} N_2^i$ and $\|N_2^i\| < \lambda$
- (b) N_1^i is $\leq_{\mathfrak{K}}$ -increasing continuous in i
- (c) N_2^i is $\leq_{\mathfrak{K}}$ -increasing continuous in i
- (d) f_i is a $\leq_{\mathfrak{K}}$ -embedding of N_1^i into M
- (e) f_i is increasing continuous in i
- (f) $a_i \in N_1^{i+1}$
- (g) $N_1^0 = N_1, N_2^0 = N_2, f_0 = id_{N_1}$
- (h) N_1^i and N_2^i has cardinality κ .

For $i = 0$, clause (g) gives the definition. For i limit let:

$$N_1^i = \bigcup_{j < i} N_1^j \text{ and}$$

$$N_2^i = \bigcup_{j < i} N_2^j \text{ and}$$

$$f_i = \bigcup_{j < i} f_j.$$

Now (a)-(f) continues to hold by continuity (and $\|N_2^i\| < \lambda$ as λ is regular).

For i successor we use our assumption; more elaborately, let $M_1^{i-1} \leq_{\mathfrak{K}} M$ be $f_{i-1}(N_1^{i-1})$ and M_2^{i-1}, g_{i-1} be such that g_{i-1} is an isomorphism from N_2^{i-1} onto M_2^{i-1} extending f_{i-1} , so $M_1^{i-1} \leq_{\mathfrak{K}} M_2^{i-1}$ (but without loss of generality $M_2^{i-1} \cap M = M_1^{i-1}$). Now apply the saturation assumption with M, M_1^{i-1} , $\text{tp}(g_{i-1}(a_{i-1}), M_1^{i-1}, M_2^{i-1})$ here standing for N, M, p there (note: $a_{i-1} \in N_2 = N_2^0 \subseteq N_2^{i-1}$ and $\mu > \|N_2^{i-1}\| = \|M_2^{i-1}\|$ and $\|M_1^{i-1}\| = \|N_1^{i-1}\| \geq \|N_1^0\| = \|N_1\| \geq \mu_0$ so the requirements and also the cardinality in Definition 0.22 holds). So there is $b \in M$ such that $\text{tp}(b, M_1^{i-1}, M) = \text{tp}(g_{i-1}(a_{i-1}), M_1^{i-1}, M_2^{i-1})$. Moreover (if $\mathfrak{K}$ has amalgamation in μ the proof is slightly shorter) remembering the second sentence in 0.22(1) which speaks about “strongly realizes” there is $b \in M$ such that b strongly realizes $(M_1^{i-1}, M_2^{i-1}, g_{i-1}(a_{i-1}))/E_{M_1^{i-1}}^{\text{at}}$ in M . This means (see Definition 0.17 (3)) that for some $M_1^{i,*}$ we have $M_1^{i-1} \leq_{\mathfrak{K}} M_1^{i,*} \leq_{\mathfrak{K}} M$ and $(M_1^{i-1}, M_2^{i-1}, g_{i-1}(a_{i-1}))/E_{M_1^{i-1}}^{\text{at}}(M_1^{i-1}, M_1^{i,*}b)$. This means (see Definition 0.17(1) that M' too has cardinality κ and there is $M_2^{i,*} \in K_\mu$ such that $M_1^{i-1} \leq_{\mathfrak{K}} M_2^{i,*}$ and there are $\leq_{\mathfrak{K}}$ -embeddings h_2^i, h_1^i of $M_2^{i-1}, M_1^{i,*}$ into $M_2^{i,*}$ over M_1^{i-1} respectively, such that $h_2^i(g_{i-1}(a_{i-1})) = h_1^i(b)$.

Now changing names, without loss of generality h_2^i is the identity.

Let N_2^i, h_i be such that $N_2^{i-1} <_{\mathfrak{K}} N_2^i, h_i$ an isomorphism from N_2^i onto $M_2^{i,*}$ extending g_{i-1} . Let $N_1^i = h_i^{-1} \circ h_1^i(M_1^{i,*}), f_i = (h_1^i)^{-1} \circ (h_i \upharpoonright N_1^i)$.

We have carried the induction. Now f_κ is a $\leq_{\mathfrak{K}}$ -embedding of N_1^κ into M over N_1 , but $|N_2| = \{a_i : i < \kappa\} \subseteq N_1^\kappa$, so $f_\kappa \upharpoonright N_2 : N_2 \rightarrow M$ is as required.

2), 3) By the hence and forth argument (or see [Sh 300, II, §3] = [Sh:e, II, §3]).

4) Easy, too. $\square_{0.26}$

0.27 Remark. Note that by 0.26(2) if M is μ -saturated above μ_0 and $\mathfrak{K}$ has the JEP $_{\mu_0}$ then $\mathfrak{K}$ has λ -amalgamation for each $\lambda \in [\mu_0, \mu)$.

0.28 Definition. Fix $\lambda \geq \kappa$ with κ regular.

1) We say that $N_1 \in K_\lambda$ is (λ, κ) -saturated over N_0 or that $(N_1, c)_{c \in N_0}$ is (λ, κ) -saturated (and λ -saturated of cofinality κ means (λ, κ) -saturated) if:

there is a sequence $\langle M_i : i < \kappa \rangle$ which is $\leq_{\mathfrak{K}}$ -increasing continuous with $M_0 = N, M_\kappa = N_1$ and $M_{i+1} \in K_\lambda$ universal over M_i (see 0.19(2)).

2) If we omit κ , we mean $\kappa = \text{cf}(\lambda)$; (λ, α) -saturated means $(\lambda, \text{cf}(\alpha))$ -saturated; and N_1 is $(\lambda, 1)$ -saturated over N_0 means just $N_0 \leq_{\mathfrak{K}} N_1$ are in K_λ .

0.29 Claim. Fix $\lambda \geq \kappa$ with κ regular.

- 1) If N_1 is (λ, κ) -saturated over N , then N_1 is unique over N .
- 2) If $K_\lambda \neq \emptyset$, $\lambda \geq \kappa = \text{cf}(\kappa)$ and over every $M \in K_\lambda$ there is N with $M \leq_{\mathfrak{K}} N \in K_\lambda$ universal over M , then for every $N \in K_\lambda$ there is $N_1 \in K_\lambda$ which is (λ, κ) -saturated over N .
- 3) If $\mathfrak{K}_\lambda$ has amalgamation and $\mathfrak{K}$ is stable in λ (i.e. $M \in K_\lambda \Rightarrow |\mathcal{S}(M)| \leq \lambda$) then every $M \in K_\lambda$ has a universal extension (so part (2)'s conclusion holds).

Proof. See [Sh 300, Ch.II] or check. $\square_{0.29}$

We do not need at present but recall from [Sh 88]:

0.30 Claim. There is $\tau' \supseteq \tau \cup \{P_0, P_1, P_2, c\}$ of cardinality $\leq LS(\mathfrak{K})$ with c an individual constant, with P_ℓ unary predicates, and a set Γ of quantifier free types such that:

- (a) if $M' \in PC_{\tau'}(\emptyset, \Gamma)$ and $M_\ell = (M' \upharpoonright \tau) \upharpoonright P_\ell^{M'}$ for $\ell = 0, 1, 2$, then $M_\ell \in K$, $M_0 \leq_{\mathfrak{K}} M_1$, $M_0 \leq_{\mathfrak{K}} M_2$, $c^{M'} \in M_2$, and $N' \subseteq M' \Rightarrow (N' \upharpoonright \tau) \upharpoonright P_\ell^{N'} \leq_{\mathfrak{K}} M_\ell$, and there is no $b \in M_1$ satisfying:
 - ⊗ for every $\bar{a} \in {}^{\omega>}(P_0^{M'})$, letting $N_{\bar{a}}$ be the τ' -submodel of M' generated by $\bar{a}$ and $M_{\bar{a}}^\ell = M_\ell \upharpoonright (M_\ell \cap N_{\bar{a}})$, we have $M_{\bar{a}}^\ell \leq_{\mathfrak{K}} M_\ell$, and b strongly realizes $tp(c^{M'}, M_{\bar{a}}^0, M_{\bar{a}}^2)$ in $M_{\bar{a}}^1$
- (b) if M_ℓ^*, c are as in (a) and $M_0 = M_1 \cap M_2$. then for some M' we have clause (a).

Proof. Should be clear; see [Sh 88]. $\square_{0.30}$

0.31 Remark.

- 1) Claim 0.30 enables us to translate results of the form: two cardinal with omitting types theorem in λ_2 implies the existence of one in λ_1 , provided that types are local in the sense that $p \in \mathcal{S}(M)$ is determined by $\langle p \upharpoonright N : N \leq_{\mathfrak{K}} M, \|N\| \leq \lambda \rangle$.
- 2) This enables us to prove implications between cases of λ -categoricity, if we have a nice enough theory of types as in [Sh:c, VIII, §4]; if we have in λ_2 a saturated model, categoricity in λ_1 implies categoricity in λ_2 . Also (if we know a little more) categoricity in λ_2 is equivalent to the non-existence of a non-saturated model in λ_2 .

0.32 Claim. 1) Assume $M_n \leq_{\mathfrak{K}} M_{n+1}$, $M_n \in K_\lambda$, $\mathfrak{K}$ has amalgamation in λ . If $p_n \in \mathcal{S}(\mathcal{M}_n)$, $p_n \leq p_{n+1}$ (i.e. $p_n = p_{n+1} \upharpoonright M_n$, see 0.19(6)), then there is $p \in \mathcal{S}(\bigcup_{n < \omega} M_n)$ such that $n < \omega \Rightarrow p_n \leq p$.

- 2) If $\langle M_i : i \leq \delta \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous, $p_i \in \mathcal{S}(M_i)$, $(j < i \Rightarrow p_j \leq p_i)$, $p_i = tp(a_i, M_i, N_i)$ and $h_{i,j}$ is a $\leq_{\mathfrak{K}}$ -embedding of N_j into N_i (for $j < i < \delta$) such that $h_{i,j} \upharpoonright M_j = id_{M_j}$, $h_{i,j}(a_j) = a_i$, then there is $p_\delta \in \mathcal{S}(M_\delta)$, $i \leq \delta \Rightarrow p_i \leq p_\delta$.
- 3) If $\mathfrak{K}$ has amalgamation in λ and is stable in λ (i.e. $M \in K_\lambda \Rightarrow |\mathcal{S}(M)| \leq \lambda$), then

- (a) *every $M \in K_\lambda$ has a universal extension;*
- (b) *for every $M \in K_\lambda$ and regular $\theta \leq \lambda$ there is $N \in K_\lambda$ which is (λ, θ) -saturated over M .*

§1 WEAK DIAMOND

1.1 Definition. Fix λ regular and uncountable.

1)

$$\text{WDmTId}(\lambda, S, \bar{\chi}) = \left\{ A : A \subseteq \prod_{\alpha \in S} \chi_\alpha, \text{ and for some function } F \text{ with domain } \bigcup_{\alpha < \lambda} {}^\alpha(2^{<\lambda}) \text{ mapping } {}^\alpha(2^{<\lambda}) \text{ into } \chi_\alpha, \right. \\ \left. \text{for every } \eta \in A, \text{ for some } f \in {}^\lambda(2^{<\lambda}) \text{ the set } \{ \delta \in S : \eta(\delta) = F(f \upharpoonright \delta) \} \text{ is not stationary} \right\}.$$

(Note: WDmTId stand for weak diamond target ideal)¹.

Here we can replace $2^{<\lambda}$ by any set of this cardinality, and so we can replace $f \in {}^\lambda(2^{<\lambda})$ by $f_1, \dots, f_n \in {}^\lambda(2^{<\lambda})$ with F being n -place.

$$2) \quad \text{cov}_{\text{wdmt}}(\lambda, S, \bar{\chi}) = \text{Min} \left\{ |\mathcal{P}| : \mathcal{P} \subseteq \text{WDmTId}(\lambda, S) \text{ and } {}^S 2 \subseteq \bigcup_{A \in \mathcal{P}} A \right\}$$

3)

$$\text{WDmTId}_{<\mu}(\lambda, S, \bar{\chi}) = \left\{ A : \text{for some } i^* < \mu \text{ and } A_i \in \text{WDmTId}(\lambda, S) \text{ for } i < i^* \text{ we have } A \subseteq \bigcup_{i < i^*} A_i \right\}$$

$$4) \quad \text{WDmId}_{<\mu}(\lambda, \bar{\chi}) = \left\{ S \subseteq \lambda : \text{cov}_{\text{wdmt}}(\lambda, S) < \mu \right\}.$$

5) Instead of “ $< \mu^+$ ” we write μ , if we omit μ we mean $(2^{<\lambda})$. If $\bar{\chi}$ is constantly 2 we may omit it, if $\chi_\alpha = 2^{|\alpha|}$ we may write pow instead of $\bar{\chi}$.

$$6) \quad \text{Let } \mu_{\text{wd}}(\lambda, \bar{\chi}) = \text{cov}_{\text{wdmt}}(\lambda, \lambda, \bar{\chi}).$$

7) We say that the weak diamond holds on λ if $\lambda \notin \text{WDmId}(\lambda)$.

By [DvSh 65], [Sh:b, XIV, 1.18(2), 1.8] (presented better in [Sh:f, AP, §1], note: 1.2(4) below rely on [Sh 460]) we have:

¹in [Sh:b, AP, §1], [Sh:f, AP, §1] we express $\text{cov}_{\text{wdmt}}(\lambda, S) > \mu^*$ by allowing $f(0) \in \mu^* < \mu$

1.2 Theorem.

1) If $\lambda = \aleph_1, 2^{\aleph_0} < 2^{\aleph_1}, \mu \leq (2^{\aleph_0})$ or even $2^\theta = 2^{<\lambda} < 2^\lambda, \mu = (2^\theta)^+$, or just: for some $\theta, 2^\theta = 2^{<\lambda} < 2^\lambda, \mu \leq 2^\lambda$, and $\chi^{<\lambda} < \mu$ for $\chi < \mu$, then $\lambda \notin \text{WDmId}_{<\mu}(\lambda)$.
 If in addition $(*)_{<\mu, \lambda}$ below holds, then $\lambda \notin \text{WDmId}_{<\mu}(\lambda, \text{pow})$, where:

$(*)_{\mu, \lambda}$ there are no $A_i \in [\mu]^{\lambda^+}$ for $i < 2^\lambda$ such that $i \neq j \Rightarrow |A_i \cap A_j| < \aleph_0$

and $(*)_{<\mu, \lambda}$ means $(*)_{\chi, \lambda}$ holds for $\chi < \mu$.

2) If $\mu \leq \lambda^+$ or $\text{cf}([\mu_1]^{\leq \lambda}, \subseteq) < \mu$ for $\mu_1 < \mu$ or $\mu = \aleph_0$ then $\text{WDmId}_{<\mu}(\lambda, \bar{\chi})$ is a normal ideal on λ . If this ideal is not trivial, then $\lambda = \text{cf}(\lambda) > \aleph_0, 2^{<\lambda} < 2^\lambda$.

3) A sufficient condition for $(*)_{<\mu, \lambda}$ is:

(a) $\mu \leq 2^\lambda$ & $(\forall \alpha < \mu)(|\alpha|^{\aleph_0} < 2^\lambda)$.

4) Another sufficient condition for $(*)_{<\mu, \lambda}$ is:

(b) $\mu \leq 2^\lambda$ & $\lambda \geq \beth_\omega$.

1.3 Remark. 1) So if $\text{cf}(2^\lambda) < \mu$ (which holds if 2^λ is singular and $\mu = 2^\lambda$) then $(*)_{<\mu, \lambda}$ implies that there is $A \subseteq {}^\lambda 2, |A| < 2^\lambda, A \notin \text{WDmTId}(\lambda)$.

2) Some related definitions appear in 1.13; we use them below (mainly $\text{DfWD}_{<\mu}(\lambda)$), but as in a first reading it is recommended to ignore them, the definition is given later.

3) We did not look again at the case $(\forall \sigma < \lambda)(2^\sigma < 2^{<\lambda} < 2^\lambda)$.

As in [Sh 88, 3.5], ([Sh 87a, 2.7], [Sh 87b, 6.3]):

1.4 Claim. Assume $\lambda \notin \text{WDmId}_{<\mu}(\lambda)$ or at least $\text{DfWD}_{<\mu}(\lambda)$

(where $\lambda = \text{cf}(\lambda) > \aleph_0$) and $\mathfrak{K}$ is an abstract elementary class.

1) Assume $\mathfrak{K}$ is categorical in $\chi, \lambda = \chi^+$, and $\mathfrak{K}$ has a model in λ (if $LS(\mathfrak{K}) \leq \chi$ this is equivalent to: the model $M \in K_\chi$ is not $\leq_{\mathfrak{K}}$ -maximal). Assume further $\mathfrak{K}$ does not have the λ -amalgamation property in χ . Then for any $M_i \in \mathfrak{K}_\lambda$ for $i < i^* < \mu$, there is $N \in \mathfrak{K}_\lambda$ not $\leq_{\mathfrak{K}}$ -embeddable into any M_i (and the assumptions of part (2) below holds).

2) Assume $M_\eta \in K_{<\lambda}$ for $\eta \in {}^\lambda 2, M_\eta = \bigcup_{\alpha < \text{lg}(\eta)} M_{\eta \restriction (\alpha+1)}, \nu \triangleleft \eta \Rightarrow M_\nu \leq_{\mathfrak{K}} M_\eta$,

and $M_{\eta \restriction \langle 0 \rangle}, M_{\eta \restriction \langle 1 \rangle}$ cannot be amalgamated over M_η (hence $M_\eta \neq M_{\eta \restriction \langle \ell \rangle}$). Set

$M_\eta =: \bigcup_{\alpha < \lambda} M_{\eta \restriction \alpha}$ for $\eta \in {}^\lambda 2$. Clearly M_η belongs to K_λ . For the $\text{DfWD}_{<\mu}(\lambda)$

version assume also

$(*) \langle M_\eta : \eta \in {}^\lambda 2 \rangle$ is definable (even just by $\mathcal{L}_{\lambda, \lambda}$) in

$\mathfrak{B} = (\mathcal{H}(\chi), \in, <_\chi^*, \mathfrak{K}_{<\chi}, \lambda, \mu)$.

Then for any $N_i \in \mathfrak{K}_\lambda$ for $i < i^* < \mu$, there is $\eta \in {}^\lambda 2$ such that: M_η is not $\leq_{\mathfrak{K}}$ -embeddable into any N_i .

3) In part (2), if $LS(\mathfrak{K}) \leq \lambda$ we can allow $N_i \in K_{\kappa_i}$ if $\sum_{i < i^*} \text{cf}([\kappa_i]^\lambda, \subseteq) < \mu$.

4) If we use $\text{DfWD}_{<\mu}(\lambda)$: assume only $\lambda \notin \text{WDmId}_{<\mu}(\lambda, \bar{\chi})$. Part (2) holds if M_η

is defined for $\eta \in \bigcup_{\alpha < \lambda} \prod_{i < \alpha} \chi_i$, and $\varepsilon < \zeta < \chi_i, \eta \in \prod_{j < i} \chi_j \Rightarrow M_{\eta \hat{\ } \langle \varepsilon \rangle}, M_{\eta \hat{\ } \langle \zeta \rangle}$ cannot be amalgamated over M_η . The assumption of Part (4) holds (hence the conclusion of Part (2)) holds if we assume that for $M \in K_\chi, i < \lambda$ there are $\chi_i \leq_{\mathfrak{K}}$ -extensions of M in K_λ , which pairwise cannot be amalgamated over M .

Proof. 1) It is straightforward to choose $M_\eta \in K_\chi$ for $\eta \in {}^\alpha 2$ by induction on α , as required in part (2). Then use part (2) to get the desired conclusion.

2) Without loss of generality the universe of N_i is λ and the universe of M_η is an ordinal γ_η such that $\eta \in {}^{\lambda > 2} \Rightarrow \gamma_\eta < \lambda$ and $\eta \in {}^\lambda 2 \Rightarrow \gamma_\eta = \lambda$. The reader can ignore the “DfWD $_{<\mu}(\lambda)$ ” version (ignoring the h_η ’s, g) if he likes. For $\alpha < \lambda$ and $\eta \in {}^\alpha 2$ let the function h_η be $h_\eta(i) = M_{\eta \upharpoonright (i+1)}$ for $i < \ell g(\eta)$. Let $\langle M_\eta : \eta \in {}^{\lambda > 2} \rangle$ be the $<_\chi^*$ -first such object. For each $i < i^*$ we define $A_i \subseteq {}^\lambda 2$ by $A_i = \{\eta \in {}^\lambda 2 : M_\eta \text{ can be } \leq_{\mathfrak{K}}\text{-embedded into } N_i\}$.

For $\eta \in A_i$ choose $f_\eta : M_\eta \rightarrow N_i$, a $\leq_{\mathfrak{K}}$ -embedding, hence $f_\eta \in {}^\lambda \lambda$. We also define a function F_i from $\bigcup_{\alpha < \lambda} ({}^\alpha 2 \times {}^\alpha \lambda)$ to $\{0, 1\}$ by:

$$F_i(\eta, f) \text{ is : } 0 \quad \text{if } f \text{ is a } \leq_{\mathfrak{K}}\text{-embedding of } M_\eta \text{ into } N_i \\ \text{with range } \subseteq \ell g(\eta) \text{ which can be extended to a} \\ \leq_{\mathfrak{K}}\text{-embedding of } M_{\eta \hat{\ } \langle 0 \rangle} \text{ into } N_i$$

$$F_i(\eta, f) \text{ is } 1 \quad \text{otherwise.}$$

Now for any $\eta \in A_i$, the set

$$E = \{\delta < \lambda : \gamma_\eta \upharpoonright \delta = \delta \text{ and } f_\eta \upharpoonright \delta \text{ is a function from } \delta \text{ to } \delta\}$$

is a club of λ . For every $\delta \in E$ clearly $F(\eta \upharpoonright \delta, f_\eta \upharpoonright \delta) = \eta(\delta)$ (as $M_{\eta \hat{\ } \langle 0 \rangle}, M_{\eta \hat{\ } \langle 1 \rangle}$ cannot be amalgamated over M_η).

Hence (for the “Def” version see Definition 1.13(2) using 1.14(1), 1.14(3)) we have $A_i \in \text{WDmTId}^{\text{Def}}(\lambda)$. As $i^* < \mu$ clearly $\bigcup_{i < i^*} A_i \in \text{WDmId}_{<\mu}^{\text{Def}}(\lambda)$ and hence

by assumption of the claim ${}^\lambda 2 \neq \bigcup_i A_i$. Take $\eta \in {}^\lambda 2 \setminus \bigcup_{i < i^*} A_i$. Then M_η is as required.

3) Without loss of generality the universe of N_i is κ_i . Let $\mathcal{P}_i \subseteq [\kappa_i]^\lambda$ be a set of minimal cardinality such that $(\forall B)[B \subseteq \kappa_i \ \& \ |B| \leq \lambda \rightarrow (\exists B' \in \mathcal{P}_i)(B \subseteq B')]$. As $LS(\mathfrak{K}) \leq \lambda$ we can find for each $A \in \mathcal{P}_i$, a model $N_A^i \leq_{\mathfrak{K}} N_i$ of cardinality λ^+ whose universe includes A . Now apply part (2) to $\{N_A^i : i < i^* \text{ and } A \in \mathcal{P}_i\}$.

4) Same proof. $\square_{1.4}$

We give three variants of the preceding:

1.5 Claim. 1) Assume

- (*)₁ $\lambda = \text{cf}(\lambda) > \aleph_0$
 - (a) M_η is a τ -model of cardinality $< \lambda$ for $\eta \in {}^\lambda 2$ and
 - (b) for each $\eta \in {}^\lambda 2$, $\langle M_{\eta \upharpoonright \alpha} : \alpha < \lambda \rangle$ is $\subseteq$ -increasing continuous with union, called $M_\eta \in K$, of cardinality λ
 - (c) if $\eta \in {}^\lambda 2$, $\eta \triangleleft \rho_\ell \in {}^\lambda 2$ for $\ell = 1, 2$ then M_{ρ_1}, M_{ρ_2} are not isomorphic over M_η

- (*)₂ $\lambda \notin \text{WDmId}_{<\mu}(\lambda)$

Then $I(\lambda, K) \geq \mu$, and in fact we can find $X \subseteq {}^\lambda 2$ of cardinality $\geq \mu$ such that $\{M_\rho : \rho \in X\}$ are pairwise non-isomorphic.

2) Assume

- (*)₁^d like (*)₁ in part (1) but in addition
 - (d) $\langle M_\eta : \eta \in {}^\lambda 2 \rangle$ is definable in $\mathfrak{B} = \mathfrak{B}_X$

- (*)₂^d $\lambda \notin \text{DfWD}_{<\mu}(\lambda)$.

Then the conclusion of part (1) holds.

3) The parallel of 1.4(4) holds.

Proof. 1) Let $\{N_i : i < i^*\}$ be a maximal subset of $\{M_\rho : \rho \in {}^\lambda 2\}$ consisting of pairwise non-isomorphic models, and use the proof of 1.4(2) with $f_\eta : N_i \simeq M_\eta$.

2), 3) Left to the reader. $\square_{1.5}$

1.6 Claim. 1) Assume

- (*)₁ $M_\eta \in K_{<\lambda}$ for $\eta \in {}^\lambda 2$, $\langle M_{\eta \upharpoonright \alpha} : \alpha \leq \ell g(\eta) \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous, and $M_{\eta \upharpoonright \langle 1 \rangle}$ cannot be $\leq_{\mathfrak{K}}$ -embedded into M_ν over M_η when $\eta \hat{\ } \langle 0 \rangle \leq \nu \in {}^\lambda 2$
- (*)₂ $\lambda = \text{cf}(\lambda) > \aleph_0$, $\lambda \notin \text{WDmId}_{<\mu}(\lambda)$, and λ is a successor cardinal, or at least there is no λ -saturated normal ideal on λ , or at least $\text{WDmId}(\lambda)$ is not λ -saturated (which holds if for some $\theta < \lambda$, $\{\delta < \lambda : \text{cf}(\delta) = \theta\} \notin \text{WDmId}(\lambda)$).

Then there is $A \subseteq {}^\lambda 2$, $|A| = 2^\lambda$ such that: if $\eta_1 \neq \eta_2$ are in A then (taking $M_\eta = \bigcup_{\alpha < \lambda} M_{\eta \upharpoonright \alpha}$)

- (α) $M_{\eta_1} \not\cong M_{\eta_2}$ for $\eta_1 \neq \eta_2 \in A$ and
- (β) if $(2^\chi)^+ < 2^\lambda$ for $\chi < \lambda$ then we can also achieve: M_{η_1} cannot be $\leq_{\mathfrak{K}}$ -embedded into M_{η_2} .

2) Under the assumptions of 1.4(1) we can find $\langle M_\eta : \eta \in {}^\lambda 2 \rangle$ as in the assumption of 1.4(2).

3) Under the assumption of 1.4(2) the assumption of 1.6(1) holds.

- 4) Under the assumption of 1.6(1) we have $I(\lambda, \aleph) = 2^\lambda$ and if $(2^\lambda)^+ < 2^\lambda$ then $IE(\lambda, \aleph) = 2^\lambda$.
 5) The parallel of 1.4(4) holds.

Proof of 1.6. 1) The proof of [Sh 88, 3.5] works (see the implications preceding it). More elaborately, we divide the proof into cases according to the answer to the following:

Question: Is there $\eta^* \in {}^{\lambda>2}$ such that for every ν satisfying $\eta^* \trianglelefteq \nu \in {}^{\lambda>2}$ there are $\rho_0, \rho_1 \in {}^{\lambda>2}$ such that: $\nu \triangleleft \rho_0, \nu \trianglelefteq \rho_1$, and M_{ρ_0}, M_{ρ_1} cannot be amalgamated over M_{η^*} ?

We can find a function $h : {}^{\lambda>2} \rightarrow {}^{\lambda>2}$, such that:

- (a) the function h is one-to-one, preserving $\triangleleft$ and $(h(\nu))^{\wedge \langle \ell \rangle} \trianglelefteq h(\nu^{\wedge \langle \ell \rangle})$
- (b)_{yes} when the answer to the question is yes, it is exemplified by $\eta^* = h(\langle \rangle)$ and $M_{h(\nu^{\wedge \langle 0 \rangle})}, M_{h(\nu^{\wedge \langle 1 \rangle})}$ cannot be amalgamated over M_{η^*} (for every $\nu \in {}^{\lambda>2}$)
- (b)_{no} when the answer to the question above is no, $h(\langle \rangle) = \langle \rangle$ and if $\nu^{\wedge \langle 0 \rangle} \trianglelefteq \rho_0, \nu^{\wedge \langle 1 \rangle} \triangleleft \rho_1$ then $M_{h(\rho_0)}, M_{h(\rho_1)}$ can be amalgamated over $M_{h(\nu)}$.

Without loss of generality h is the identity, by renaming (and we can preserve $(*)_1^d$ of 1.5(2) in the relevant case). Also clearly $M_{\eta^{\wedge \langle \ell \rangle}} \neq M_\eta$ (by the non-amalgamation assumption).

Case 1 The answer is yes. We do not use the non λ -saturation of $\text{WdId}(\lambda)$ in this case.

For any $\eta \in {}^{\lambda 2}$ and $\leq_{\aleph}$ -embedding g of $M_{\langle \rangle}$ into $M_\eta =: \bigcup_{\alpha < \lambda} M_{\eta \upharpoonright \alpha}$, let

$$A_{\eta, g} =: \{ \nu \in {}^{\lambda 2} : \text{there is a } \leq_{\aleph} \text{-embedding of } M_\nu \text{ into } M_\eta \text{ extending } g \}$$

$$A_\eta =: \{ \nu \in {}^{\lambda 2} : \text{there is a } \leq_{\aleph} \text{-embedding of } M_\nu \text{ into } M_\eta \}.$$

So: $|A_{\eta, g}| \leq 1$ and $\eta \in A_\eta$ (as if $\nu_1, \nu_2 \in A_{\eta, g}$ are distinct then for some ordinal $\alpha < \lambda$ and $\nu \in {}^{\alpha 2}$ we have $\nu =: \nu_0 \upharpoonright \alpha = \nu_1 \upharpoonright \alpha, \nu_0(\alpha) \neq \nu_1(\alpha)$ and use the choice of $h(\nu^{\wedge \langle \ell \rangle})$).

Since $A_\eta = \bigcup \{ A_{\nu, g} : g \text{ is a } \leq_{\aleph} \text{-embedding of } M_{\langle \rangle} \text{ into } M_\eta \}$, we have $|A_\eta| \leq 2^{<\lambda}$.

Hence we can choose by induction on $\zeta < 2^\lambda, \eta_\zeta \in {}^{\lambda 2} \setminus \bigcup_{\xi < \zeta} A_{\eta_\xi}$ (existing by cardinality considerations as $2^{<\lambda} < \lambda$). Then $\xi < \zeta \Rightarrow M_{\eta_\xi} \not\cong M_{\eta_\zeta}$ so we have proved clause (a).

If $(2^{<\lambda})^+ < 2^\lambda$, then use the Hajnal free subset theorem ([Ha61]) to choose distinct $\{ \eta_\zeta : \zeta < 2^\lambda \} \subseteq {}^{\lambda 2}$ such that $\zeta \neq \xi \Rightarrow \eta_\zeta \notin A_{\eta_\xi}$, so we have proved clause (b), too.

Case 2: The answer is no.

Again, without loss of generality M_η has as universe the ordinal γ_η .

Let $\langle S_i : i < \lambda \rangle$ be a partition of λ to sets, none of which is in $\text{WdId}(\lambda)$. For each i we define a function F_i as follows:

if $\delta \in S_i, \eta, \nu \in {}^\delta 2, \gamma_\eta = \gamma_\nu = \delta$, and $f : \delta \rightarrow \delta$ then

$$F_i(\eta, \nu, f) = 0 \text{ if } f \text{ can be extended to an embedding of } M_{\nu \hat{\ } \langle 0 \rangle} \text{ into some } M_\rho \text{ with } \eta \hat{\ } \langle 0 \rangle \leq \rho.$$

$$F_i(\eta, \nu, f) = 1 \text{ otherwise.}$$

So as $S_i \notin \text{WdId}(\lambda)$, for some $\eta_i^* \in {}^\lambda 2$ we have:

- (*) for every $\eta \in {}^\lambda 2, \nu \in {}^\lambda 2$ and $f \in {}^\lambda \lambda$ the following set of ordinals $i < \lambda$ is stationary:

$$\{\delta \in S_i : F_i(\eta \upharpoonright \delta, \nu \upharpoonright \delta, f \upharpoonright \delta) = \eta_i^*(\delta)\}.$$

Now for any $X \subseteq \lambda$ let $\eta_X, \rho_X \in {}^\lambda 2$ be defined by:

$$\begin{aligned} \text{if } \alpha \in S_i \text{ then } i \in X \Rightarrow \eta_X(\alpha) = 1 - \eta_i^*(\alpha), i \notin X \Rightarrow \eta_X(\alpha) = 0 \text{ and} \\ \rho_X = \eta_{\{2i : i \in X\} \cup \{2i+1 : i \notin X\}}. \end{aligned}$$

Now we show

- (*) if $X, Y \subseteq \lambda$, and $X \neq Y$ then M_{ρ_X} cannot be $\leq_{\mathfrak{K}}$ -embedded into M_{ρ_Y} .

Clearly (*) will suffice for finishing the proof.

Assume toward a contradiction that f is a $\leq_{\mathfrak{K}}$ -embedding of M_{ρ_X} into M_{ρ_Y} ; as $X \neq Y$ there is i such that $i \in X \Leftrightarrow i \notin Y$ so there is $j \in \{2i, 2i+1\}$ such that $\rho_X \upharpoonright S_j = \langle 1 - \eta_j^*(\alpha) : \alpha \in S_j \rangle$ and $\rho_Y \upharpoonright S_j$ is identically zero. Clearly $E = \{\delta : f \text{ maps } \delta \text{ into } \delta\}$ is a club of λ and hence $S_j \cap E \neq \emptyset$.

So if $\delta \in S_j \cap E$ then $f \upharpoonright M_{(\rho_Y \upharpoonright \delta) \hat{\ } \langle 0 \rangle} = f \upharpoonright M_{(\rho_X \upharpoonright \delta) \hat{\ } \langle 0 \rangle}$ extend $f \upharpoonright M_{\rho_Y \upharpoonright \delta}$ and is a $\leq_{\mathfrak{K}}$ -embedding of it into some $M_{(\rho_X \upharpoonright \alpha)}$, $\alpha < \lambda$ large enough.

Now by the choice of h we get

$$\delta \in S_j \cap E \Rightarrow F(\rho_X \upharpoonright \delta, \rho_Y \upharpoonright \delta, f \upharpoonright \delta) = \rho_X(\delta) = 1 - \eta_j^*(\delta).$$

But this contradicts the choice of η_j^* .

2) This corresponds to Case I, where we do not need the weak diamond, but just $2^{<\lambda} < 2^\lambda$.

3), 4), 5). Check, similarly. □_{1.6}

1.7 Conclusion. 1) Assume

- (*)₁ for $\eta \in {}^{\lambda > 2}$, $M_\eta \in K_{<\lambda}$ and $\langle M_{\eta \upharpoonright \alpha} : \alpha \leq \ell g(\eta) \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous and $M_{\eta \hat{\ } \langle 1 \rangle}$ cannot be $\leq_{\mathfrak{K}}$ -embedded into M_ν over M_η when $\eta \hat{\ } \langle 0 \rangle \leq \nu \in {}^{\lambda > 2}$ and $\langle M_\eta : \eta \in {}^{\lambda > 2} \rangle$ is definable in $\mathfrak{B}$
- (*)₂^d $\text{WdId}^{\text{Def}}(\lambda)$ or $\text{DfWD}^+(\lambda)$ is not λ -saturated (which holds if there is no normal λ -saturated ideals on λ (which holds for non Mahlo λ) and holds if for some θ , $\{\delta < \lambda : \text{cf}(\delta) = \theta\}$ is not in the ideal).

Then the conclusion of 1.6 holds.

From the Definition below, we use here mainly “superlimit”

1.8 Definition. 1) $M \in \mathfrak{K}_\lambda$ is a *superlimit* if

- (a) for every $N \in \mathfrak{K}_\lambda$ such that $M \leq_{\mathfrak{K}} N$ there is $M' \in K_\lambda$ such that $N \leq_{\mathfrak{K}} M', N \neq M'$, and $M \cong M'$;
- (b) if $\delta < \lambda^+$ is limit, $\langle M_i : i < \delta \rangle$ is $\leq_{\mathfrak{K}}$ -increasing and $M_i \cong M$ (for $i < \delta$) then $\bigcup_{i < \delta} M_i \cong M$.

2) For $\Theta \subseteq \{\mu : \aleph_0 \leq \mu \leq \lambda, \mu \text{ regular}\}$ we say $M \in \mathfrak{K}_\lambda$ is a (λ, Θ) -superlimit if:

- (a) clause (a) from part (1) holds and
- (b) if $M_i \cong M$ is $(\leq_{\mathfrak{K}})$ -increasing for $i < \mu \in \Theta$ then $\bigcup_{i < \mu} M_i \cong M$.

3) For $S \subset \lambda^+$ we say $M \in \mathfrak{K}_\lambda$ is a (λ, S) -strong limit if:

- (a) clause (a) from part (1) holds
- (b) there is a function F from $\bigcup_{\alpha < \kappa} {}^\alpha(K_\lambda)$ to K_λ such that:
 - (α) for any sequence $\langle M_i : i < \alpha \rangle$ if $\alpha < \kappa, M_0 = M, M_i$ is $\leq_{\mathfrak{K}}$ -increasing, and $M_i \in \mathfrak{K}_\lambda$, then $j < \alpha \Rightarrow M_j \leq_{\mathfrak{K}} F(\langle M_i : i < \alpha \rangle)$
 - (β) if $\langle M_i : i < \lambda^+ \rangle$ is $\leq_{\mathfrak{K}}$ -increasing, $M_0 = M, M_i \in \mathfrak{K}_\lambda$, and for $i < \kappa, M_{i+1} \leq_{\mathfrak{K}} F(\langle M_j : j \leq i+1 \rangle) \leq_{\mathfrak{K}} M_{i+2}$ then $\{\delta \in S \mid \bigcup_{i < \delta} M_i \not\cong M\}$ is not stationary.

4) M is a (λ, κ) -limit if there is a function F as in 3b(α) such that:

- (a) if $\langle M_i : i < \kappa \rangle$ is a $<_{\mathfrak{K}}$ -increasing continuous sequence in K_λ , $F(\bar{M} \upharpoonright (i+1)) \leq_{\mathfrak{K}} M_{i+1}$ then $\bigcup_{i < \kappa} M_i \cong M$
- (b) there is at least one such sequence.

5) M is a (λ, κ) -superlimit is defined similarly, but with F omitted and $M_{i+1} \cong M$.

1.9 Claim. 1) In 1.4(1) we can replace the categoricity of $\mathfrak{K}$ in χ by “ $\mathfrak{K}$ has a super limit model in χ ” which is not an amalgamation base (see Definition 1.8). In this case the assumption of 1.4(2), and of 1.6(1) holds.

2) We can weaken (in 1.6(5) the existence of superlimit to) “for some $\kappa = \text{cf}(\kappa) \leq \chi$ there is a $(\chi, \{\kappa\})$ -super limit model which is not an amalgamation base”; provided that we add $\{\delta < \lambda : \text{cf}(\delta) = \kappa\} \notin \text{WDmId}_{<\mu}(\lambda)$ (but for 2.9(1) we need now “ $\text{WDmId}_{<\mu}(\lambda) + \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$ is not λ -saturated”). If there is $S \subseteq \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$, which belongs to $I[\lambda]$ but not to $\text{WDmId}_{<\mu}(\lambda)$ we can weaken the model theoretic requirement to: there is a $(\chi, \{\kappa\})$ -medium limit (see [Sh 88, Definition §3]) but not used here.

1.10 Claim. Assume $2^\lambda < 2^{\lambda^+}$.

0) If $\mathfrak{K}$ (an abstract elementary class) is categorical in λ , $LS(\mathfrak{K}) \leq \lambda$, $I(\lambda^+, \mathfrak{K}) < 2^{\lambda^+}$, then $\mathfrak{K}_\lambda$ has amalgamation.

1) If $\mathfrak{K}$ (an abstract elementary class) is categorical in λ and $1 \leq IE(\lambda^+, K) < 2^{\lambda^+}$ but $K_{\lambda^{++}} = \emptyset$ then $\mathfrak{K}$ has amalgamation in λ .

2) Assume $\mathfrak{K}$ has amalgamation in λ , $LS(\mathfrak{K}) \leq \lambda$, $K_{\lambda^+} \neq \emptyset$ and $K_{\lambda^{++}} = \emptyset$. Then there is $M \in K_{\lambda^+}$ saturated above λ .

3) If M is μ -saturated above λ , $LS(\mathfrak{K}) \leq \lambda_0 < \lambda$ and $\mathfrak{K}$ has amalgamation in every $\lambda'_0 \in [\lambda_0, \lambda)$ then M is μ -saturated above λ_0 .

1.11 Remark. If $I(\lambda^+, \mathfrak{K}) < 2^{\lambda^+}$, then the assumption $\mathfrak{K}_{\lambda^{++}} = \emptyset$ is not used in part (1) of 1.10; this is 1.6(1) + (2). Also if $(2^\lambda)^+ < 2^{\lambda^+}$ then the assumption $K_{\lambda^{++}} = \emptyset$ is not needed in part (1) of 1.10; by 1.6(1) + (2) (note (b) of 1.6(1)).

Proof. 0) By 1.6(1) applied to λ^+ .

1) If not, we can choose for $\eta \in \lambda^+ > 2$ a model $M_\eta \in \mathfrak{K}_\lambda$ such that $[\nu \triangleleft \eta \Rightarrow M_\nu \leq_{\mathfrak{K}} M_\eta]$, and $M_{\eta \cdot \langle 0 \rangle}, M_{\eta \cdot \langle 1 \rangle}$ cannot be amalgamated over M_η . If

$(2^\lambda)^+ < 2^{\lambda^+}$ we are done by 1.6(1), so assume $(2^\lambda)^+ = 2^{\lambda^+}$. For each

$\eta \in \lambda^+ > 2$ let $M_\eta =: \bigcup_{\alpha < \lambda^+} M_{\eta \upharpoonright \alpha}$, and let $N_\eta \in \mathfrak{K}_{\lambda^+}$ be such that

$M_\eta \leq_{\mathfrak{K}} N_\eta$, N_η is $\leq_{\mathfrak{K}}$ -maximal (exists as $\mathfrak{K}_{\lambda^{++}} = \emptyset$). Now we choose by induction $\zeta < (2^\lambda)^+$, $\eta_\zeta \in \lambda^+ > 2$ such that M_{η_ζ} is not $\leq_{\mathfrak{K}}$ -embeddable into N_{η_ξ} for $\xi < \zeta$ (exists by 1.4(2)). So necessarily for $\xi < \zeta$, N_{η_ζ} is not $\leq_{\mathfrak{K}}$ -embeddable into N_{η_ξ} (as $M_{\eta_\zeta} \leq N_{\eta_\zeta}$). Also, for $\xi < \zeta$, N_{η_ξ} is not $\leq_{\mathfrak{K}}$ -embeddable into N_{η_ζ} as otherwise by the maximality of N_ξ this implies $N_\xi \cong N_\zeta$. So $\{N_\zeta : \zeta < 2^{\lambda^+}\}$ exemplifies $IE(\lambda^+, \mathfrak{K}) = 2^{\lambda^+}$, contradicting an assumption.

2) A maximal model in K_{λ^+} will do.

3) Easy. □_{1.10}

* * *

1.12 Discussion Instead of Weak Diamond we now discuss Definable Weak Diamond, which is weaker but suffices.

Compare [MkSh 313], where many Cohen subsets are added to λ and a combinatorial principle about amalgamation of configurations $\langle M_s : s \subseteq n, s \neq n \rangle$ is gotten.

We are interested here in the case $n = 1$ (ordinary amalgamation); in §3, also $n = 2$. Even more definability can be required.

This is particularly interesting when we look at results under some other set theory, when combining $2^\lambda = 2^{\lambda^+}$ with definable weak diamond on λ^+ is helpful. This played a major role in the preliminary form of this work.

1.13 Definition. 1) In Definition 1.1 we add the superscript $\mathcal{F}$ if we restrict ourselves to functions $F \in \mathcal{F}$.

2) Fix a model $\mathfrak{B}$ whose universe includes λ and has a definable pairing function on λ , and a logic $\mathcal{L}$ closed under first order operations and substitution; also allow “ $M \in \mathfrak{K}$ ” and “ $M \leq_{\mathfrak{K}} N$ ” in the formulas, if it is not said otherwise. Let

$$\mathcal{F}_{\mathfrak{B}, \mathcal{L}}^{\text{Def}} = \left\{ F : \text{for some } g \in {}^\lambda \lambda \text{ and } \bar{h} = \langle h_\eta : \eta \in {}^{\lambda > 2} \rangle \text{ where} \right. \\
\begin{aligned}
& h_\eta : \ell g(\eta) \rightarrow \lambda \text{ and } h_{\eta \restriction \beta} \subseteq h_\eta \text{ for } \beta < \ell g(\eta) \\
& \text{and for some sequence } \bar{\psi} = \langle \psi_\alpha : \alpha < \lambda \rangle, \text{ with} \\
& \psi_\alpha \in \mathcal{L} \text{ for } \alpha < \lambda \text{ the following holds for every } \alpha < \lambda \text{ and} \\
& f \in {}^\alpha (2^{< \lambda}) : F(f) = 1 \text{ iff } (\mathfrak{B}, \alpha, g, h_f) \models \psi_\alpha \text{ iff } F(f) \neq 0 \Big\}.
\end{aligned}$$

3) The version of weak diamond from 1.1, restricted to the class $\mathcal{F}$ of 1.13(2) is called the $(\mathfrak{B}, \mathcal{L})$ -definitional version. If $\mathcal{L}$ is $\mathcal{L}_{\lambda, \lambda}$ we may omit it. If $\mathfrak{B}$ has the form $(\mathcal{H}(\chi), \in, <_\chi^*, \lambda)$ we write $\mathcal{F}_{\mathcal{L}}^{\text{Def}[\chi]}$ or $\mathcal{F}^{\text{Def}(\chi)}$ instead of $\mathcal{F}_{\mathfrak{B}, \mathcal{L}}^{\text{Def}}$ or $\mathcal{F}_{\mathfrak{B}}^{\text{Def}}$ respectively. If we omit χ , we mean $\chi = (2^\lambda)^+$ and we may put $\text{Def}(\chi)$ or Def instead of $\mathcal{F}^{\text{Def}(\chi)}$ or $\mathcal{F}^{\text{Def}}$ in the superscript. Having the definitional version or the definable weak diamond for λ means $\lambda \notin \text{WdId}^{\text{Def}}(\lambda)$.

4) Let $\text{DfWD}_{<\mu}(\lambda)$ mean that with $\mathfrak{B} = (\mathcal{H}(\chi), \in, <_\chi^*, \lambda, \mu)$ we have $\lambda \notin \text{WdId}_{<\lambda}^{\text{Def}}(\lambda)$.

Instead of “ $< \mu^+$ ” we write “ μ ” and instead of “ $< 2^{< \lambda^+}$ ” we may write nothing.

5) Let $\text{DfWD}_{<\mu}^+(\lambda)$ mean $\text{DfWD}_{<\mu}(\lambda)$ together with the principle $\otimes_\lambda$ below; we adopt the same conventions as in (4) concerning μ :

$$\otimes_\lambda \text{ if for } \eta \in {}^{\lambda > 2}, M_\eta \text{ is a } \tau_{\mathfrak{R}}\text{-model of cardinality } < \lambda, \langle M_{\eta \restriction \alpha} : \alpha \leq \ell g(\eta) \rangle \text{ is} \\
\leq\text{-increasing continuous, for } \eta \in {}^{\lambda > 2} \text{ we let } M_\eta = \bigcup_{\alpha < \lambda} M_{\eta \restriction \alpha} \text{ and for} \\
\eta \neq \nu \in {}^{\lambda > 2}, M_\eta \text{ and } M_\nu \text{ are not isomorphic over } M_{\langle \rangle}, \text{ then } \{M_\eta / \cong : \eta \in {}^{\lambda > 2}\} \\
\text{has cardinality } 2^\lambda \\
\text{(note that } 2^{< \lambda} < 2^\lambda \text{ implies that)}.$$

1.14 Claim.

1) Assume $\mathcal{L}$ first order or at least definable enriching first order.

In the definition of $\mathcal{F}_{\mathfrak{B}, \mathcal{L}}^{\text{Def}}$, we can replace “for every $\alpha < \lambda$ ” by “for a club of $\alpha < \lambda$ ”. In the definition of $\mathcal{F}^{\text{Def}}$ we can let $g \in {}^\lambda ({}^{\lambda > 2})$ and $h_\eta : \ell g(\eta) \rightarrow {}^{\lambda > 2}$. In any case $\text{WdId}_{<\mu}^{\mathcal{F}}(\lambda)$ increases with $\mathcal{F}$ and is $\subseteq \text{WdId}_{<\mu}(\lambda)$, similarly for $\text{WdId}_{<\mu}^{\mathcal{F}}(\lambda)$.

2) If $\mathcal{F} = \mathcal{F}_{\mathfrak{B}}^{\text{Def}}$ and $\text{cf}(\mu) > \lambda$ then $\text{WdId}_{<\mu}(\lambda)$ is a normal ideal (but possibly is equal to $\mathcal{P}(\lambda)$).

3) Assume $V \models “\lambda = \chi^+, \chi^{<\chi} = \chi, \mu > \lambda”$ and P is the forcing notion of adding μ Cohen subsets to χ (i.e.

$\{g : g \text{ a partial function from } \mu \text{ to } \{0\} \text{ with domain of cardinality } < \chi\}$).

Then in V^P we have $\text{WdId}_{<\mu}^{\text{Def}}(\lambda)$ is the ideal of non-stationary subsets of λ ; i.e. with

$\mathfrak{B} = (\mathcal{H}(\chi), \in, <_\mu^*)^{V^P}$ for any χ . Also $\otimes_\lambda$ of Definition 1.13(5) holds.

Remark. In 1.14(1) we use the assumption on $\mathcal{L}$; anyhow not serious: reread the definition 1.1(1).

Proof. 1); 2) By manipulating the h 's (using the pairing function on λ).
 3) See [MkSh 313] or think (the point being that we can break the forcing, first adding $\bar{\psi}$ and g (or the $< \mu$ ones) and then (read 1.1(1)) choose $\eta \in {}^\lambda 2$ as $g \restriction [\gamma, \gamma + \lambda)$ not “used before”. Now for any candidate $f \in {}^\lambda({}^{\lambda > 2})$ for a club of $\tilde{\delta} < \lambda$, $\eta(\delta) = g(\gamma + \delta)$ is not used in the definition of $f \restriction \delta$, $h_f \restriction \delta$ so stationarily often $\eta(\delta)$ “guesses” rightly.) $\square_{1.14}$

1.15 Claim. 1) If $2^\theta = 2^{<\lambda} < 2^\lambda$ then $DfWD^+(\lambda)$.
 2) $DfWD_{<\mu}(\lambda)$ holds when $\lambda \notin WDMId_{<\mu}(\lambda)$ (see 1.2 for sufficient conditions).

1.16 Discussion: We hope to get successful “guessing” not just on a stationary set, but on a positive set for the same ideal for which have guessed; i.e. there is I a normal ideal on λ such that for $A \in I^+$ there is $\eta \in {}^\lambda 2$ guessing I -positively; this is connected to questions on λ^+ -saturation. For more see [Sh 638].

We phrased the following notion originally in the hope of later eliminating $\mu_{wd}(\lambda)$ (i.e. using 2^λ instead of $\mu_{wd}(\lambda)$).

1.17 Definition. 1)

$$\begin{aligned} \text{UDmId}_{<\mu}^{\mathcal{F}}(\lambda) = \Big\{ S \subseteq \lambda : & \text{for some } i^* < \mu \text{ and } F_i \in \mathcal{F} \text{ (for } i < i^*) \\ & \text{for every } \eta \in {}^S 2 \text{ there is } f \in {}^\lambda(2^{<\lambda}) \text{ and } i < i^* \\ & \text{and club } E \text{ of } \lambda \text{ such that:} \\ & \text{for every } \delta \in E \text{ we have:} \\ & \delta \in S \Rightarrow \eta(\delta) = F_i(f \restriction \delta) \\ & \delta \in \lambda \setminus S \Rightarrow 0 = F_i(f \restriction \delta) \Big\}. \end{aligned}$$

2) We omit μ if $\mu = 1$.

3) $BA^{\mathcal{F}}(\lambda)$ is defined as the family of $S \subseteq \lambda$ such that for some $F \in \mathcal{F}$ and $\eta = O_S$ the condition above holds.

1.18 Claim. Assume $\mathcal{F} = \mathcal{F}_{\mathfrak{B}}^{\text{Def}}$.

- 1) In the definition 1.17(1) we can replace $f \in {}^\lambda(2^{<\lambda})$ by $f \in {}^\lambda 2$ or $f \in {}^\lambda({}^{\lambda > 2})$.
- 2) $\text{UDmId}^{\mathcal{F}}(\lambda)$ is a normal ideal on λ (but possibly is $\mathcal{P}(\lambda)$).
- 3) $BA^{\mathcal{F}}(\lambda)$ is a Boolean algebra of subsets of λ including all non-stationary subsets of λ and even $\text{UDmId}^{\mathcal{F}}(\lambda)$, and is closed under unions of $< \lambda$ sets and even under diagonal union.
- 4) If $S \in BA^{\mathcal{F}}(\lambda)$ and $F \in \mathcal{F}$ then for some $\eta \in {}^S 2$ we have:

(*) for every $f \in {}^\lambda({}^{\lambda > 2})$ we have

$$\{\delta \in S : \eta(\delta) = F(f \restriction \delta)\} \neq \emptyset \text{ mod } \text{UDmId}^{\mathcal{F}}(\lambda).$$

5) $UDmId_{<\mu}^{\mathcal{F}}(\lambda) \subseteq WDmId_{<\mu}^{\mathcal{F}}(\lambda) \subseteq WDmId_{<\mu}(\lambda)$ and they increase with $\mathcal{F}$ and $\lambda \in UDmId_{<\mu}^{\mathcal{F}}(\lambda) \Leftrightarrow \lambda \in WDmId_{<\mu}^{\mathcal{F}}(\lambda)$.

Proof. Straightforward. $\square_{1.18}$

1.19 Discussion Remember

- (*)₁ If $V = “\chi = \chi^{<\chi} \ \& \ 2^\chi = \chi^+”$, P is the forcing notion of adding $\mu > \chi^+$ Cohen subsets to χ then in V^χ , any equivalence relation on $\mathcal{P}(\lambda)$ definable with parameters $X \subseteq \chi$ and ordinals which has at least χ^{++} equivalence classes has at least μ equivalence classes

and (see [Sh:f, XVI,§2])

- (*)₂ ZFC is consistent with $CH+$ for some stationary, costationary $S \subseteq \omega_1$ we have

- (a) $WDmId(\aleph_1) = \{A \subseteq \omega_1 : A \setminus S \text{ is not stationary}\}$
- (b) $\mathcal{D}_{\omega_1} + S$ is $\aleph_2$ -saturated.

- (*)₃ $ZFC + GCH$ is consistent with $\{\delta < \aleph_2 : cf(\delta) = \aleph_1\} \in WDmId(\aleph_2)$

- (*)₄ $ZFC + 2^{\aleph_1} < 2^{\aleph_2}$ is consistent with $\{\delta < \aleph_2 : cf(\delta) = \aleph_0\} \in WDmId(\aleph_2)$.

§2 FIRST ATTEMPTS

Given amalgamation in K_λ (cf. 2.2) we try to define and analyze types $p \in \mathcal{S}(M)$ for $M \in K_\lambda$. But types here (as in [Sh 300]) are not sets of formulas. They may instead be represented by triples (M, N, a) with $M \leq_{\mathfrak{K}} N$ and $a \in N \setminus M$. We look for nice types (i.e. triples) and try to prove mainly the density of the set of minimal types).

To simplify matters we allow uses of stronger assumptions than are ultimately desired (e.g. $2^{\lambda^+} > \lambda^{++}$ and/or $K_{\lambda^{++}} = \emptyset$). These will later be eliminated. However the first extra assumption is still a “mild set theoretic assumption”, and the second is harmless if we think only of proving our main theorem 0.2 and not on subsequent continuations.

So the aim of this section is to show that we can start to analyze such classes.

2.1 Hypothesis. $\mathfrak{K}$ is an abstract elementary class.

2.2 Claim. *Assume*

$(*)_\lambda^2$ K is categorical in λ ; $1 \leq I(\lambda^+, K) < 2^{\lambda^+}$; $LS(\mathfrak{K}) \leq \lambda$ and: $2^\lambda < 2^{\lambda^+}$, or at least the definable weak diamond holds for λ^+ holds.

Then

- 1) $\mathfrak{K}_\lambda$ has amalgamation.
- 2) If $I(\lambda^{++}, K) = 0$ then $\mathfrak{K}$ has a model in λ^+ which is universal homogeneous above λ , hence saturated above λ (see 0.22(2)).
- 3) If $I(\lambda^{++}, K) = 0$ then $M \in K_\lambda \Rightarrow |\mathcal{S}(M)| \leq \lambda^+$.

Proof. 1) If amalgamation fails in K_λ and $2^\lambda < 2^{\lambda^+}$, then the assumptions of 1.4(1) hold with λ^+ in place of λ . Hence by 1.6(2) the statement $(*)_1$ of 1.6(1) (see there) holds and easily also $(*)_2$ of 1.6(1), hence by 1.6(4) we have $I(\lambda^+, K) = 2^{\lambda^+}$, a contradiction. If $2^\lambda = 2^{\lambda^+}$, we are using the variants from 1.13.

2) As $I(\lambda^{++}, K) = 0 < I(\lambda^+, K)$, there is $M \in K_{\lambda^+}$ which is maximal. If M is not universal homogeneous above λ then there are $N_0, N_1 \in K_\lambda$ with $N_0 \leq_{\mathfrak{K}} M$ and $N_0 \leq_{\mathfrak{K}} N_1$ such that N_1 cannot be $\leq_{\mathfrak{K}}$ -embedded into M over N_0 . Use 0.20(2) to get a contradiction.

3) Follows from (2). □_{2.2}

2.3 Definition.

1)(a) $K_\lambda^3 = \{(M_0, M_1, a) : M_0 \leq_{\mathfrak{K}} M_1 \text{ are both in } K_\lambda \text{ and } a \in M_1 \setminus M_0\}$.

(b) $(M_0, M_1, a) \leq (M'_0, M'_1, a')$ iff $a = a'$, $M_0 \leq_{\mathfrak{K}} M'_0$, $M_1 \leq_{\mathfrak{K}} M'_1$

(c) $(M_0, M_1, a) \leq_h (M'_0, M'_1, a')$ iff $h(a) = a'$, and for $\ell = 0, 1$ we have:
 $h \upharpoonright M_\ell$ is a $\leq_{\mathfrak{K}}$ -embedding of M_ℓ into M'_ℓ .

$$(d) \quad (M_0, M_1, a) < (M'_0, M'_1, a') \text{ if } (M_0, M_1, a) \leq (M'_0, M'_1, a) \text{ and } M_0 \neq M'_0.$$

$$(e) \quad \text{similarly } <_h$$

- 2) $(M_0, M_1, a) \in K_\lambda^3$ has the weak extension property if there is $(M'_0, M'_1, a) \in K_\lambda^3$ such that $(M_0, M_1, a) \leq (M'_0, M'_1, a)$ and $M_0 \neq M'_0$.
- 3) $(M_0, M_1, a) \in K_\lambda^3$ has the extension property if: for every $N_0 \in \mathfrak{K}_\lambda$ and $\leq_{\mathfrak{K}}$ -embedding f of M_0 into N_0 there are N_1, b and g such that: $(M_0, M_1, a) \leq_g (N_0, N_1, b) \in K_\lambda^3$ and $g \supseteq f$ (so $g(a) = b$ and g is a $\leq_{\mathfrak{K}}$ -embedding of M_1 into N_1).

2.4 Claim. Assume

$$(*)_\lambda^3 \quad LS(\mathfrak{K}) \leq \lambda, K \text{ is categorical in } \lambda \text{ and in } \lambda^+, \text{ and } 1 \leq I(\lambda^{++}, K).$$

Then every $(M_0, M_1, a) \in K_\lambda^3$ has the weak extension property that is:
If $M_0 \leq_{\mathfrak{K}} M_1$ are in K_λ and $a \in M_1 \setminus M_0$, then we can find M'_1 in $\mathfrak{K}_\lambda$ such that: $M_0 <_{\mathfrak{K}} M'_1$ hence $M_0 \neq M'_0$ and $(M_0, M_1, a) \leq (M'_0, M'_1, a)$.

Proof. We can choose $\langle N_i, a_i : i < \lambda^+ \rangle$ such that:

- (a) $N_i \in K_\lambda$ is $\leq_{\mathfrak{K}}$ -increasing continuous in i ;
- (b) h_i is an isomorphism from M_1 onto N_{i+1} such that $h_i(M_0) = N_i, h_i(a) = a_i$.

Now as $a \in M_1 \setminus M_0$ clearly $i < j < \lambda^+ \Rightarrow a_i \in N_{i+1} \leq_{\mathfrak{K}} N_j$ & $a_j \notin N_{i+1}$ hence $\bigcup_{i < \lambda^+} N_i \in K_{\lambda^+}$.

By 0.20(1) applied to λ^+ there are $M'_0 \leq_{\mathfrak{K}} M'_1$ in K_{λ^+} , $M'_0 \neq M'_1$, and there is $b \in M'_1 \setminus M'_0$. As K is categorical in λ^+ , without loss of generality $M'_0 = \bigcup_{i < \lambda^+} N'_i$.

Let χ be large enough and $\mathfrak{B} \prec (\mathcal{H}(\chi) \in, <_\chi^*)$ be such that $\lambda \subseteq \mathfrak{B}, \|\mathfrak{B}\| = \lambda$ and $\langle N_i, a_i : i < \lambda^+ \rangle, M'_0, M'_1, b$ and the definition of $\mathfrak{K}$ belong to $\mathfrak{B}$.

Let $\delta = \mathfrak{B} \cap \lambda^+$, so $\delta \in (\lambda, \lambda^+)$ is a limit ordinal and

$$N_\delta \leq_{\mathfrak{K}} N_{\delta+1} \leq_{\mathfrak{K}} M'_1,$$

$$N_\delta \leq_{\mathfrak{K}} (M'_1 \cap \mathfrak{B}) \leq_{\mathfrak{K}} M'_1$$

$$\mathfrak{B} \cap M'_0 = N_\delta$$

$$N_{\delta+1} \cap (M'_1 \cap \mathfrak{B}) = N_\delta,$$

so for some N we have:

$$N \in \mathfrak{K}_\lambda, N \leq_{\mathfrak{K}} M'_1, \text{ and } (N_{\delta+1} \cup (M'_1 \cap \mathfrak{B})) \subseteq N$$

so (see Definition 2.3(1) above)

$$(N_\delta, N_{\delta+1}, a_\delta) \leq_{\mathfrak{K}} (M'_1 \cap \mathfrak{B}, N, a_\delta),$$

and b witnesses $N_\delta \neq M'_1 \cap \mathfrak{B}$.

As $(M_0, M_1, a) \cong (N_\delta, N_{\delta+1}, a_\delta)$, the result follows. $\square_{2.4}$

2.5 Definition. 1) $(M_0, M_1, a) \in K_\lambda^3$ is minimal when:

$$\text{if } (M_0, M_1, a) \leq_{h_\ell} (M'_0, M'_1, a_\ell) \in K_\lambda^3 \text{ for } \ell = 1, 2,$$

$$\text{then } \text{tp}(a_1, M'_0, M'_1) = \text{tp}(a_2, M'_0, M'_1).$$

2) $(M_0, M_1, a) \in K_\lambda^3$ is reduced when:

$$\text{if } (M_0, M_1, a) \leq (M'_0, M'_1, a) \in \mathfrak{K}_\lambda^3 \text{ then } M'_0 \cap M_1 = M_0.$$

3) We say $p \in \mathcal{S}(M_0)$ is minimal, where $M_0 \in K_\lambda$, if for some a, M_1 we have: $p = \text{tp}(a, M_0, M_1)$ and $(M_0, M_1, a) \in K_\lambda^3$ is minimal.

4) We say $p \in \mathcal{S}(M_0)$ is reduced where $M_0 \in K_\lambda$, if for some a, M_1 we have $p = \text{tp}(a, M_0, M_1)$ and $(M_0, M_1, a) \in K_\lambda^3$ is reduced.

2.6 Fact. 1) For every $(M_0, M_1, a) \in K_\lambda^3$ there is a reduced (M'_0, M'_1, a) such that: $(M_0, M_1, a) \leq (M'_0, M'_1, a) \in K_\lambda^3$.

2) Assume $\langle (M_{0,\alpha}, M_{1,\alpha}, a) : \alpha < \delta \rangle$ is an increasing sequence of members of K_λ^3

$$(a) \text{ if } \delta < \lambda^+ \text{ then } (M_{0,\alpha}, M_{1,\alpha}, a) \leq (\bigcup_{\beta < \delta} M_{0,\beta}, \bigcup_{\beta < \delta} M_{1,\beta}, a) \in K_\lambda^3.$$

(b) If $\delta = \lambda^+$ the result may be in $K_{\lambda^+}^3$: if $\{\alpha < \delta : M_{0,\alpha} \neq M_{0,\alpha+1}\}$ is cofinal, this holds.

$$(c) \text{ If } \delta < \lambda^+ \text{ and each } (M_{0,\alpha}, M_{1,\alpha}, a) \text{ is reduced then so is } (\bigcup_{\beta < \delta} M_{0,\beta}, \bigcup_{\beta < \delta} M_{1,\beta}, a).$$

3) If $(M_0, M_1, a) \leq (M'_0, M'_1, a)$ are in K_λ^3 and the first triple is minimal then so is the second.

4) If $(M_0, M_1, a) \leq (M'_0, M'_1, a)$ are in K_λ^3 then $\text{tp}(a, M_0, M_1) \leq \text{tp}(a, M'_0, M'_1)$; (Definition 0.24(1)).

5) If K_λ has amalgamation, then: $(M_0, M_1, a) \in K_\lambda^3$ is minimal if and only if:

$$(*) \text{ If } (M_0, M_1, a) \leq_{h_\ell} (M'_0, M'_1, a_\ell) \in K_\lambda^3 \text{ for } \ell = 1, 2 \text{ and } h_1 \upharpoonright M_0 = h_2 \upharpoonright M_0 \text{ then } \text{tp}(a_1, M'_0, M'_1) = \text{tp}(a_2, M'_0, M'_1).$$

- 6) If there is no maximal member² of K_λ^3 and there are $N_0 <_{\mathfrak{K}} N_1$ in K_λ , then there are $N^0 <_{\mathfrak{K}} N^1$ in $\mathfrak{K}_{\lambda^+}$.
- 7) If every triple in K_λ^3 has the weak extension property, and there are $N_0 <_{\mathfrak{K}} N_1$ in $\mathfrak{K}_\lambda$ then there are $N^0 <_{\mathfrak{K}} N^1$ in $\mathfrak{K}_{\lambda^+}$.
- 8) If $LS(\mathfrak{K}) \leq \lambda$ and every triple in K_λ^3 has the extension property and $K_\lambda^3 \neq \emptyset$ then no $M \in K_{\lambda^+}$ is $<_{\mathfrak{K}}$ -maximal hence $K_{\lambda^{++}} \neq \emptyset$.
- 9) If $LS(\mathfrak{K}) \leq \lambda$ and $K_{\lambda^+} \neq \emptyset$, then $K_\lambda^3 \neq \emptyset$.

Proof. Easy. Note that part (7) is 2.4.

2.7 Claim. Assume $(*)_{\lambda^+}^2 + (*)_{\lambda}^3$ (i.e. the hypothesis of 2.2 and 2.4) and $2^{\lambda^+} > \lambda^{++}$, and $K_{\lambda^{+3}} = \emptyset$.

Then in $\mathfrak{K}_\lambda^3$ the minimal triples are dense (i.e. above every triple in K_λ^3 there is a minimal one).

Remark. We do not intend to adopt the hypotheses “ $2^{\lambda^+} > \lambda^{++}$ ”, $K_{\lambda^{+3}} = \emptyset$ indefinitely. They will be eliminated in §3.

Proof. If not, we can choose by induction on $\alpha < \lambda^+$, for $\eta \in {}^\alpha 2$ a triple $(M_\eta^0, M_\eta^1, a_\eta)$ and $h_{\eta, \nu}$ for $\nu \trianglelefteq \eta$ such that:

- (i) $(M_\eta^0, M_\eta^1, a_\eta) \in K_\lambda^3$
- (ii) $\nu \triangleleft \eta \Rightarrow (M_\nu^0, M_\nu^1, a_\nu) \leq_{h_{\eta, \nu}} (M_\eta^0, M_\eta^1, a_\eta)$
- (iii) $\nu_0 \triangleleft \nu_1 \triangleleft \nu_2 \Rightarrow h_{\nu_2, \nu_0} = h_{\nu_2, \nu_1} \circ h_{\nu_1, \nu_0}$
- (iv) $(M_{\eta^\wedge \langle \ell \rangle}^0, M_{\eta^\wedge \langle \ell \rangle}^1, h_{\eta^\wedge \langle \ell \rangle, \eta} \upharpoonright M_\eta^0)$ for $\ell = 0, 1$ are equal
- (v) $\text{tp}(a_{\eta^\wedge \langle 0 \rangle}, M_{\eta^\wedge \langle 0 \rangle}^0, M_{\eta^\wedge \langle 0 \rangle}^1) \neq \text{tp}(a_{\eta^\wedge \langle 1 \rangle}, M_{\eta^\wedge \langle 1 \rangle}^0, M_{\eta^\wedge \langle 1 \rangle}^1)$;
this makes sense as $M_{\eta^\wedge \langle 0 \rangle}^0 = M_{\eta^\wedge \langle 1 \rangle}^0$
- (vi) if $\eta \in {}^\delta 2$ and $\delta < \lambda^+$ is a limit ordinal, then $M_\eta^\ell = \bigcup_{\alpha < \delta} h_{\eta, \eta \upharpoonright \alpha}(M_{\eta \upharpoonright \alpha}^\ell)$
for $\ell = 0, 1$,
- (vii) $(M_{< >}^0, M_{< >}^1, a_{< >}) \in K_\lambda^3$ is a triple above which there is no minimal one.

This is straightforward: for $\alpha = 0$ choose a triple in K_λ^3 above which supposedly there is no minimal triple; in limit α take limits of diagrams (chasing the h 's); in successor α , use non-minimality and 2.6(5).

Let $M^* \in \mathfrak{K}_{\lambda^{++}}$ be saturated above λ (exists by 2.2(2) so it is necessarily homogeneous universal above λ^+ , hence above λ ; note: λ there stands for λ^+ here).

We choose by induction on $\alpha < \lambda^+$ for $\eta \in {}^\alpha 2$, a $\leq_{\mathfrak{K}}$ -embedding g_η of M_η^0 into M^* such that:

$$\nu \triangleleft \eta \Rightarrow g_\nu = g_\eta \circ h_{\eta, \nu}.$$

$$g_{\eta^\wedge \langle 0 \rangle} = g_{\eta^\wedge \langle 1 \rangle}.$$

²will be applied for λ^+

This is clearly possible. Let $N_\eta^0 = M^* \upharpoonright \text{Rang}(g_\eta)$. For $\eta \in {}^{\lambda^+}2$ let $N_\eta^0 = M^* \upharpoonright \bigcup_{\alpha < \lambda^+} \text{Rang}(g_{\eta \upharpoonright \alpha})$ and let $g_\eta = \bigcup_{\alpha < \lambda^+} g_{\eta \upharpoonright \alpha}$. Chasing arrows we can find for $\eta \in {}^{\lambda^+}2$ a limit to $(\langle M_{\eta \upharpoonright \alpha}^0, M_{\eta \upharpoonright \alpha}^1, a_{\eta \upharpoonright \alpha} \rangle, h_{\eta \upharpoonright \beta, \eta \upharpoonright \alpha} : \alpha < \beta < \lambda^+)$ say $(M_\eta^0, M_\eta^1, a_\eta) \in K_{\lambda^+}^3$ and $h_{\eta, \nu}$ for $\nu \triangleleft \eta$ as usual. Let f_η be the function from M_η^0 into M^* such that for $\alpha < \lambda^+$ we have $f_\eta \circ h_{\eta, \eta \upharpoonright \alpha} = g_\eta$. So f_η is a $\leq_{\mathfrak{K}}$ -embedding of M_η^0 into M^* . So we can extend f_η to f_η^+ , a $\leq_{\mathfrak{K}}$ -embedding of M_η^1 into M^* .

Let $a_\eta^* = f_\eta^+(a_\eta)$ for $\eta \in {}^{\lambda^+}2$.

As $2^{\lambda^+} > \lambda^{++}$ for some $\eta_0 \neq \eta_1$ we have $a_{\eta_0}^* = a_{\eta_1}^*$. So for some $\alpha < \lambda^+$, $\eta_0 \upharpoonright \alpha = \eta_1 \upharpoonright \alpha$ but $\eta_0(\alpha) \neq \eta_1(\alpha)$, without loss of generality $\eta_\ell(\alpha) = \ell$ and by clause (v) above we get a contradiction. $\square_{2.7}$

2.8 Claim. 1) Assume $(*)_\lambda^2$ or just

$(*)_\lambda^{2^-}$ $\mathfrak{K}$ has amalgamation in λ and $LS(\mathfrak{K}) \leq \lambda$.

If $M_0 \leq_{\mathfrak{K}} N_0 \in K_{\lambda^+}$ and $(M_0, M_1, a) \in K_\lambda^3$ then there is $N \in K_{\leq \lambda^+}$ such that: $N_0 \leq_{\mathfrak{K}} N$ and for every $c \in N$ satisfying $tp(c, M_0, N) = tp(a, M_0, M_1)$, there is a $\leq_{\mathfrak{K}}$ -embedding h of M_1 into N extending id_{M_0} such that $h(a) = c$ and $N \notin K_{\lambda^+} \Rightarrow N$ is a $<_{\mathfrak{K}}$ -maximal member of K_λ .

2) Assume $M_0 \leq_{\mathfrak{K}} N_0 \in K_\lambda$ and $(M_0, M_1, a) \in K_\lambda^3$ has the weak extension property. Then there is $N \in K_{\lambda^+}$ such that: $N_0 \leq_{\mathfrak{K}} N$ and for every $c \in N'$ either for some $N' \in K_\lambda$ we have $N_0 \cup \{c\} \subseteq N \leq_{\mathfrak{K}} N$ and c does not strongly realize $tp(a, M_0, M_1)$ or there is an $\leq_{\mathfrak{K}}$ -embedding h of M_1 into N extending id_{M_0} such that $h(a) = c$.

Proof. 1) We choose by induction on $\alpha < \lambda^+$, a model $N_\alpha \in K_\lambda$ increasing (by $\leq_{\mathfrak{K}}$) continuous such that: for α even $N_\alpha \neq N_{\alpha+1}$ if N_α is not $\leq_{\mathfrak{K}}$ -maximal, and for α odd let $\beta_\alpha = \text{Min}\{\beta : \beta = \alpha + 1 \text{ or } \beta \leq \alpha \text{ and there is } c \in N_\beta \text{ such that there is no } \leq_{\mathfrak{K}}\text{-embedding } h \text{ of } M_1 \text{ into } N_\alpha \text{ extending } id_{M_0} \text{ such that } h(a) = c \text{ but for some } N \in K_\lambda, N_\alpha \leq_{\mathfrak{K}} N \text{ and there is a } \leq_{\mathfrak{K}}\text{-embedding } h \text{ of } M_1 \text{ into } N \text{ extending } id_{M_0} \text{ such that } h(a) = c\}$, and if $\beta_\alpha \leq \alpha$ then choose N exemplifying this and let $N_{\alpha+1} = N$. By the definition of type we are done.

2) Same proof, note that the non $\leq_{\mathfrak{K}}$ -maximality of N_α (and hence N) follows by a weak extension property.

2.9 Claim. Assume $(*)_\lambda^2$ or just:

$(*)_\lambda^{2^-}$ $\mathfrak{K}$ has amalgamation for λ and $LS(\mathfrak{K}) \leq \lambda$.

1) Assume that above $(M_0, M_1, a) \in K_\lambda^3$ there is no minimal member of K_λ^3 then (M_0, M_1, a) itself has the extension property.

2) If $(M_0, M_1, a) \in K_\lambda^3, M_0 \leq_{\mathfrak{K}} N \in K$ and the number of $c \in N$ such that $tp(c, M_0, N) = tp(a, M_0, M_1)$ is $> \lambda$ then (M_0, M_1, a) has the extension property.

3) Assume above $(M_0, M_1, a) \in K_\lambda^3$ there is no minimal member of K_λ^3 then

(*) for some N we have: $M_0 \leq_{\mathfrak{K}} N$ and N is as required in part (2).

2.10 Remark. 1) See 3.25.

2) Note that $(*)_\lambda^2$ is from 2.2 and $(*)_\lambda^2 \Rightarrow (*)_\lambda^{2^-}$ by 2.2.

Proof. 1) Follows by part (2) and (3) (part (b)).

2) Without loss of generality N has cardinality λ^+ and also is as in 2.8.

By 0.20(2) for any M'_0 such that $M_0 \leq_{\mathfrak{K}} M'_0 \in K_\lambda$ there is $N_1, N \leq_{\mathfrak{K}} N_1 \in K_{\lambda^+}$ and a $\leq_{\mathfrak{K}}$ -embedding h of M'_0 into N_1 extending id_{M_0} with $h_c(a) = c$. Now some $c \in N \setminus h(M'_0)$ realizes $\text{tp}(a, M_0, M_1)$ and there is an embedding h_c of M_1 into N extending id_{M_0} such that $h_c(a) = c$. Lastly let $N'_1 \leq_{\mathfrak{K}} N_1$ be of cardinality λ and include $\text{Rang}(h_c) \cup \text{Rang}(h)$ (send a to c via h_c).

3) We first prove

$(*)_0$ For some $M_0^+, M_0 \leq_{\mathfrak{K}} M_0^+ \in K_\lambda$ and $\text{tp}(a, M_0, M_1)$ has $> \lambda$ extensions in $\mathcal{S}(M_0^+)$ (in fact $\geq \min\{2^\mu : 2^\mu > \lambda\}$).

Proof of (a). Let $M_\eta^0, M_\eta^1, a_\eta, h_{\eta, \nu}$ ($\eta \in {}^{\lambda^+}2$ and $\nu \leq \eta$) be as in the proof of 2.7 (i.e. satisfy (i) – (vi) there) and $M_{<\zeta}^0 = M^0, M_{<\zeta}^1 = M^1$. Let $\mu = \min\{\mu : 2^\mu > \lambda\}$, so $\mu > 2$ has cardinality $\leq \lambda$ and $\mu \leq \lambda$. Let ${}^{\mu > 2} = \{\eta_\zeta : \zeta < \zeta^*\}$ be such that $\eta_\xi \triangleleft \eta_\zeta \Rightarrow \xi < \zeta$ and so $\zeta^* < \lambda^+$ and without loss of generality is a limit ordinal. Now we can choose by induction on $\zeta \leq \zeta^*$ a model $M_\zeta^* \in K_\lambda$ and, if $\zeta < \zeta^*$ also a function g_{η_ζ} such that:

- (α) M_ζ^* is $\leq_{\mathfrak{K}}$ -increasing continuous in ζ
- (β) $M_0^* = M_\emptyset$
- (γ) g_{η_ζ} is a $\leq_{\mathfrak{K}}$ -embedding of $M_{\eta_\zeta}^0$ into $M_{\zeta+1}^*$
- (δ) if $\eta_\xi \triangleleft \eta_\zeta$ then $g_{\eta_\zeta} \circ h_{\eta_\zeta, \eta_\xi} = g_{\eta_\xi}$
- (ε) if $\xi < \xi_0, \xi < \xi_1, \eta_{\xi_0} = \eta_\xi \hat{\ } \langle 0 \rangle, \eta_{\xi_1} = \eta_\xi \hat{\ } \langle 1 \rangle$ then $(M_{\xi_0}^* = M_{\xi_1}^* \text{ and } g_{\eta_{\xi_0}} = g_{\eta_{\xi_1}})$.

So for $\eta \in {}^\mu 2$ we can find g_η , a $\leq_{\mathfrak{K}}$ -embedding of M_η^0 into $M_{\zeta^*}^*$, such that

$g_\eta \circ h_{\eta, \eta \upharpoonright \alpha} = g_{\eta \upharpoonright \alpha}$ for every $\alpha < \mu$. We also can let

$p_\eta^0 = g_\eta[\text{tp}(a_\eta, M_\eta^0, M_\eta^1)] \in \mathcal{S}(M_{\zeta^*}^* \upharpoonright \text{Rang}(g_\eta))$, and find p_η such that $p_\eta^0 \leq p_\eta \in \mathcal{S}(M_{\zeta^*}^*)$ (possible as $\mathfrak{K}_\lambda$ has amalgamation by 2.2(1) if $(*)_\lambda^2$ holds and by $(*)_\lambda^{2^-}$ otherwise).

For $\eta \in {}^\mu 2$ and $\alpha \leq \mu$ let $N_{\eta \upharpoonright \alpha}^0 = M_{\zeta^*}^* \upharpoonright \text{Rang}(g_{\eta \upharpoonright \alpha})$. Clearly for $\eta \in {}^{\mu \geq 2} 2, N_\eta^0$ is well defined; $\eta \triangleleft \nu \in {}^{\mu \geq 2} 2 \Rightarrow N_\eta^0 \leq_{\mathfrak{K}} N_\nu^0$; and $N_{\eta \hat{\ } \langle 0 \rangle}^0 = N_{\eta \hat{\ } \langle 1 \rangle}^0$. Also letting for $\eta \in {}^\mu 2$, and $\alpha \leq \mu$, the type $p_{\eta \upharpoonright \alpha}^0$ be $p_\eta^0 \upharpoonright N_{\eta \upharpoonright \alpha}^0$ we have: $p_\eta^0 \in \mathcal{S}(N_\eta^0)$ is well defined, $\eta \triangleleft \nu \in {}^{\mu \geq 2} 2 \Rightarrow p_\eta^0 \leq p_\nu^0$ and $p_{\eta \hat{\ } \langle 0 \rangle}^0 \neq p_{\eta \hat{\ } \langle 1 \rangle}^0$. Hence for $\eta_0 \neq \eta_1$ from ${}^\mu 2$ we have $p_{\eta_0} \neq p_{\eta_1}$. So $|\mathcal{S}(M_{\zeta^*}^*)| \geq 2^\mu > \lambda$.

Proof of ().* We choose by induction on $i < \lambda^+, N_i \in K_\lambda$ which is $\leq_{\mathfrak{K}}$ -increasing continuous, $N_0 = M_0^+$ (M_0^+ is from $(*)_0$ above) and for each i some $c_i \in N_{i+1}$ realizes over N_0 a (complete) extension of $p = \text{tp}(a, M_0, M_1)$ not realized in N_i . There is such type by clause (*) above and there is such an N_{i+1} as $\mathfrak{K}$ has amalgamation in λ . Clearly $c_i \notin N_i$ and so $\bigcup_{i < \lambda^+} N_i$ is as required.

4) By using 0.20(2) repeatedly λ^+ times. □_{2.9}

2.11 Claim. Assume $(*)_{\lambda^+}^{2^-}$ (from 2.9; that is $\mathfrak{K}$ has amalgamation in λ and $LS(\mathfrak{K}) \leq \lambda$).

If $(M_0, M_1, a) \leq (M'_0, M'_1, a)$ are from K_λ^3 , and the second has the extension property, then so does the first.

Proof. Use amalgamation over M_0 : if $M_0 \leq_{\mathfrak{K}} N_0 \in K_\lambda$ we can find N'_0 such that $M'_0 \leq_{\mathfrak{K}} N'_0 \in K_\lambda$ and there is a $\leq_{\mathfrak{K}}$ -embedding of N_0 into N'_0 over M_0 . Now use “ (M'_0, M'_1, a) has the extension property” for N'_0 . $\square_{2.11}$

Now we introduce

2.12 Definition. For any models $M, M_0 \in K_\lambda$, any type $p \in \mathcal{S}(M_0)$ and $f_0 : M_0 \xrightarrow[\text{iso}]{\text{onto}} M$ we let $\mathcal{S}_p(M) = \mathcal{S}_M^p = \{f_0(f(p)) : f \in \text{AUT}(M_0)\}$. Note: $\mathcal{S}_M^p$ does not depend on f_0 . If K is categorical in λ , $\mathcal{S}_p(M)$ is well defined for every $M \in K_\lambda$. We write also $\mathcal{S}_{\text{tp}(a, M_0, M_1)}(M)$ or $\mathcal{S}_{(M_0, M_1, a)}(M)$ when $(M_0, M_1, a) \in K_\lambda^3$.

2.13 Claim. Assume $(*)_{\lambda^+}^{2^-} + (*)_{\lambda^+}^2 + (*)_{\lambda^+}^3 + 2^{\lambda^+} < 2^{\lambda^{++}} + K_{\lambda^{++}} = \emptyset$.
If $(M_0, M_1, a) \in K_\lambda^3$ is minimal then it has the extension property.

Remark. Instead of $2^{\lambda^+} < 2^{\lambda^{++}}$, we can just demand the definable weak diamond.

Proof. Assume not. By the previous two claims (2.9(1), 2.11) we may assume that (M_0, M_1, a) is minimal. As $\mathfrak{K}$ has amalgamation in λ^+ by 2.2(1), there is $M^* \in K_{\lambda^{++}}$ which is saturated above λ^+ (as $K_{\lambda^{++}} = \emptyset$) hence M^* is saturated above λ (by 1.10(3)). By 2.6(1) + 2.11, without loss of generality

$$\otimes_0 (M_0, M_1, a) \text{ is reduced.}$$

Let $h : M_0 \rightarrow M^* \in K_{\lambda^+}$ be a $\leq_{\mathfrak{K}}$ -embedding and let $p = \text{tp}(a, M_0, M_1)$. If $h(p)$ is realized in M^* by $\geq \lambda^+$ elements we are done by 2.9(2). So assume

$$\otimes_1 h(p) \text{ is realized by } \leq \lambda \text{ members of } M^*.$$

Similarly

$$\otimes_1^+ g \text{ is realized by } \leq \lambda \text{ members of } M^* \text{ for } q = g(\text{tp}(a, M'_0, M'_1)) \text{ if } g \text{ is a } \leq_{\mathfrak{K}}\text{-embedding of } M'_0 \text{ into } M^* \text{ and } (M'_0, M'_1, a) \geq (M_0, M_1, a).$$

Next we prove

$$\otimes_2 \text{ for some reduced } (M'_0, M'_1, a) \geq (M_0, M_1, a) \text{ from } K_\lambda^3 \text{ we have } |\mathcal{S}_{(M'_0, M'_1, a)}(M'_0)| > \lambda^+.$$

Proof of $\otimes_2$. If not, we build two non-isomorphic members of K_{λ^+} as follows.

First: Choose by induction on $i < \lambda^+$, $(N_{0,i}, N_{1,i}, a) \in K_\lambda^3$ reduced (see 2.6(1)), increasing continuously (see 2.6(2)), with $N_{0,i} \neq N_{0,i+1}$, $(N_{0,0}, N_{1,0}, a) = (M_0, M_1, a)$;

this is possible as $(N_{0,i}, N_{1,i}, a) \in K_\lambda^3$ has the weak extension property (by 2.4 see 2.3(1)). Let $N^1 = \bigcup_{i < \lambda^+} N_{0,i}$.

Second: Choose by induction on $i < \lambda^+$, $N_i^0 \leq_{\mathfrak{K}} M^*$, $\|N_i^0\| = \lambda$, N_i^0 strictly increasing continuous such that:

- (*) for every $\beta < \lambda^+$, and $q \in \mathcal{S}_{(M_0', M_1', a)}(N_\beta^0)$ for some $\gamma \in (\beta, \lambda^+)$ there are no $N', N_\gamma^0 \leq_{\mathfrak{K}} N' \in K_\lambda$ and $c \in N' \setminus N_\gamma^0$ such that c realize q .

This is straightforward by $\otimes_1 + \otimes_2$ and bookkeeping. Let $N^0 = \bigcup_{i < \lambda^+} N_i^0$. By

categoricity of K_{λ^+} there is an isomorphism g from N^1 onto N^0 , so $E = \{\delta < \lambda^+ : g \text{ maps } N_{0,\delta} \text{ onto } N_\delta^0\}$ is a club of λ^+ . Now let $\delta^* \in E$, and apply (*) for $\beta = \delta^*$, $q = g(\text{tp}(a, N_{0,\delta^*}, N_{1,\delta^*}))$ to get γ . Choose $\delta \in E$ which is $> \gamma$. Now $N_{1,\delta}$ gives a contradiction. $\square_{\otimes_2}$

Without loss of generality

$$\otimes_3 \quad |\mathcal{S}_{(M_0, M_1, a)}(M_0)| > \lambda^+ \text{ and } p = \text{tp}(a, M_0, M_1).$$

Next we claim

$$\otimes_4 \quad \text{If } M \in K_\lambda, M \leq_{\mathfrak{K}} M^*, \Gamma \subseteq \bigcup \{\mathcal{S}_p(M') : M' \leq_{\mathfrak{K}} M, \|M'\| = \lambda\}, \\ |\Gamma| \leq \lambda^+, \text{ then}$$

$$\Gamma^* =: \{q \in \mathcal{S}_p(M) : \text{there is } M', M \leq_{\mathfrak{K}} M', \|M'\| = \lambda, \\ M' \text{ realizes } q \text{ but no } r \in \Gamma\}$$

has cardinality λ^{++} , in fact $|\mathcal{S}_p(M) \setminus \Gamma^*| \leq \lambda^+$.

Proof of $\otimes_4$. Without loss of generality $|M^*| = \lambda^{++}$ (i.e. the universe of M^* is λ^{++}). For every $q \in \mathcal{S}_p(M)$ there is a triple $(M_0, M_{1,q}, a_q)$ isomorphic to (M_0, M_1, a) (hence reduced) such that $\text{tp}(a_q, M_0, M_{1,q}) = q$. As M^* is saturated above λ , by 0.26 without loss of generality $M_{1,q} \leq_{\mathfrak{K}} M^*$.

Without loss of generality $\delta < \lambda^{++}$ & $(\lambda^+ \text{ divides } \delta) \Rightarrow M_\delta =: M^* \restriction \delta \leq_{\mathfrak{K}} M^*$. Now

- (*)₀ $q_1 \neq q_2 \Rightarrow a_{q_1} \neq a_{q_2}$ and
 (*)₁ $a_q \notin \delta$ & $\delta < \lambda^{++}$ & λ^+ divides $\delta \Rightarrow M_{1,q} \cap M_\delta = M$.

[Why? As $(M, M_{1,q}, a_q)$ is reduced].

Now if $r \in \Gamma$, say $r \in \mathcal{S}_p(M'')$ then by $\otimes_1$ we know $A_r = \{c \in M^* : c \text{ realizes } r\}$ has cardinality $\leq \lambda$ and hence $A = \bigcup \{A_r : r \in \Gamma\}$ has cardinality $\leq \lambda^+$, so we can find $\delta < \lambda^{++}$ divisible by λ^+ such that $A \subseteq \delta$. So as (by $\otimes_3$) we have $|\mathcal{S}_p(M)| > \lambda^+$ hence we can find $q[\delta] \in \mathcal{S}_p(M)$ such that $a_{q[\delta]} \notin \delta$ hence $(M, M_{1,q[\delta]}, a_{q[\delta]})$, exemplifies the conclusion of $\otimes_4$. $\square_{\otimes_4}$

Final contradiction: By $\otimes_4$ we can construct 2^{λ^+} non-isomorphic members of K_{λ^+} using 1.6(1) as follows. We choose by induction on $\alpha < \lambda^+$, for every $\eta \in {}^\alpha 2$, $M_\eta, p_\eta^0, p_\eta^1$ such that:

- (a) $M_{<\alpha} = M_0$
- (b) $M_\eta \in K_\lambda$
- (c) $\langle M_{\eta \upharpoonright \beta} : \beta \leq \ell g(\eta) \rangle$ is $<_{\mathfrak{K}}$ -increasing continuous
- (d) $p_\eta \in \mathcal{S}_p(M_{\eta \upharpoonright \beta})$
- (e) for $\beta \leq \alpha$, we have $p_\eta^0, p_\eta^1 \in \mathcal{S}(M_\eta)$ and: M_η realizes $p_{\eta \upharpoonright \beta}^\ell$ iff $\beta < \alpha$ & $\ell = \eta(\beta)$.

If $\alpha = 0$ or α is a limit, there is no problem to define M_η for $\eta \in {}^\alpha 2$. If M_η is defined, we can choose by induction on $i < \lambda^{++}$, $(N_{\eta,i}, a_{\eta,i})$ such that $(M_\eta, N_{\eta,i}, a_{\eta,i}) \in K_\lambda^3$, $\text{tp}(a_{\eta,i}, M_\eta, N_{\eta,i}) \in \mathcal{S}_p(M_\eta)$ and $N_{\eta,i}$ omits any $q \in \{p_{\eta \upharpoonright \beta}^\ell : \beta < \ell g(\eta), \ell \neq \eta(\beta)\} \cup \{\text{tp}(b, M_\eta, N_{\eta,j}) : j < i \text{ and } b \in N_{\eta,j} \text{ and } \text{tp}(b, M_\eta, N_{\eta,j}) \in \mathcal{S}_p(M_\eta)\}$. By $\otimes_4$ we can choose $(N_{\eta,j}, a_{\eta,j})$. Hence $|W_{\eta,i}| \leq \lambda$ where $W_{\eta,i} = \{j < \lambda^{++} : \text{for some } b \in N_{\eta,i} \text{ we have } \text{tp}(b, M_\eta, N_{\eta,i}) = \text{tp}(a_{\eta,j}, M_\eta, N_{\eta,j})\}$. Hence we can find $i < j < \lambda^{++}$ such that $i \notin W_{\eta,j}$ & $j \notin W_{\eta,i}$. Let $M_{\eta \upharpoonright \langle 0 \rangle} = N_{\eta,i}$, $M_{\eta \upharpoonright \langle 1 \rangle} = N_{\eta,j}$, $p_\eta^0 = \text{tp}(a_{\eta,i}, M_\eta, N_{\eta,i})$, $p_\eta^1 = \text{tp}(a_{\eta,j}, M_\eta, N_{\eta,j})$. Let for $\eta \in {}^{\lambda^+} 2$, $M_\eta = \bigcup_{\alpha < \lambda^+} M_{\eta \upharpoonright \alpha}$, and apply 1.6(1). $\square_{2.12}$

2.14 Conclusion. $[(*)_\lambda^{2^-} + (*)_{\lambda^+}^2 + (*)_\lambda^3 + 2^{\lambda^+} < 2^{\lambda^{++}} + K_{\lambda^{++}} = \emptyset]$.

Every $(M_0, M_1, a) \in K_\lambda^3$ has the extension property.

Proof. By 2.11 and 2.9 + 2.12. $\square_{2.14}$

2.15 Remark. Conclusion 2.14 says in other words:
if

- (a) $LS(\mathfrak{K}) \leq \lambda$
- (b) K is categorical in λ and in λ^+
- (c) $1 \leq I(\lambda^{++}, K) < 2^{\lambda^{++}}$
- (d) $K_{\lambda^{++}}$ is empty
- (e) $2^{\lambda^+} < 2^{\lambda^{++}}$ (or just definable weak diamond)

then every triple (M_0, M_1, a) in K_λ^3 has the extension property.

2.16 Claim. $[(*)_\lambda^3]$, in other words $LS(\mathfrak{K}) \leq \lambda$; $\mathfrak{K}$ categorical in λ and in λ^+ ; and $1 \leq I(\lambda^{++}, K) < 2^{\lambda^{++}}$.

If $M_0 \leq_{\mathfrak{K}} M_1$ are in K_λ then we can find $\alpha < \lambda^+$ and $\langle N_i : i \leq \alpha \rangle$ which is $\leq_{\mathfrak{K}}$ -increasing continuous, $N_i \in K_\lambda$, $(N_i, N_{i+1}, a_i) \in K_\lambda^3$ is reduced, $M_0 = N_0$, and $M_1 \leq_{\mathfrak{K}} N_\alpha$.

Proof. If not, we can contradict categoricity in K_{λ^+} (similar to the proof of $\otimes_2$ during the proof of 2.12).

Without loss of generality $M_0 \neq M_1$. We choose by induction on $i < \lambda^+$, $N_i^0 \in K_\lambda, \leq_{\aleph}$ -increasing continuous such that $(N_i^0, N_{i+1}^0) \cong (M_0, M_1)$ (possible by 2.6(9) and the categoricity of K in λ). Let $N^0 = \bigcup_{i < \lambda^+} N_i^0$.

We choose by induction on $i < \lambda^+$, $N_i^1 \in K_\lambda, \leq_{\aleph}$ -increasing continuous and a_i such that $(N_i^1, N_{i+1}^1, a_i) \in K_\lambda^3$ is reduced and let $N^1 = \bigcup_{i < \lambda^+} N_i^1$ (possible by 2.6(1) and the categoricity of K in λ). So by the categoricity in λ^+ without loss of generality $N^1 = N^0$, hence for some $\delta_1 < \delta_2 < \lambda^+$ we have

$$N_{\delta_1}^0 = N_{\delta_1}^1, N_{\delta_2}^0 = N_{\delta_2}^1.$$

By changing names $(N_{\delta_1}^0, N_{\delta_1+1}^0) = (M_0, M_1)$ and so $\langle N_{\delta_2+i} : i \leq \delta_2 - \delta_1 \rangle$ is as required. $\square_{2.16}$

2.17 Conclusion. $[(*)_{\lambda}^{2^-} + (*)_{\lambda^+}^2 + (*)_{\lambda}^3 + 2^{\lambda^+} < 2^{\lambda^{++}} + K_{\lambda^+3} = \emptyset$, i.e. the assumption of 2.14].

K_λ has disjoint amalgamation (M_2, M_1 are in disjoint amalgamation over M_0 in M_3 if $M_0 \leq_{\aleph} M_\ell \leq_{\aleph} M_3, M_1 \cap M_2 = M_0$).

Proof. By 2.16 and iterated applications of 2.14. $\square_{2.17}$

§3 NON-STRUCTURE

The first major aim of this section is to prove the density of minimal types using as set theoretic assumptions only $2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}}$ from cardinal arithmetic. The second aim is to prepare for a proof of a weak form of uniqueness of amalgamation in $\mathfrak{K}_\lambda$. Our aim is also to explain various methods. The proofs are similar to the ones in [Sh 87b, §6].

The immediate role of this section is to get many models in λ^{++} from the assumption “the minimal triples in K_λ^3 are not dense”: in 3.25 we get this under some additional assumptions, and in 3.28 we get it using only the additional assumption $I(\lambda, K^{+3}) = 0$, which suffices for our main theorem (this does not suffice for the theorem of [Sh 600], hence is eliminated there).

But the section is done in a more general fashion, so let us first explain two general results concerning the construction of many models based on repeated “failures of amalgamation” or “nonminimality of types”.

In 3.19, we give a construction assuming the ideal of small subsets of λ^+ (that is $\text{WdId}(\lambda^+)$) is not λ^{++} -saturated, as exemplified by $\langle S_\alpha : \alpha < \lambda^{++} \rangle$. We build for $\eta \in (\lambda^{+2})^{>2}$ models $M_\eta \in K_{\lambda^+}$ such that $M_\eta = \bigcup_{\alpha < \lambda^+} M_{\eta, \alpha}$, $|M_\eta| = \lambda \times (1 + \ell g \eta)$ and $\nu \triangleleft \eta \Rightarrow M_\nu \leq_{\mathfrak{K}} M_\eta$. Building $M_{\eta \smallfrown \langle \ell \rangle}$ manufacture $M_{\eta \smallfrown \langle \ell \rangle, \alpha+1}$ as a limit of models $\langle M_{\eta \smallfrown \langle \ell \rangle, \alpha} : \alpha < \lambda^+ \rangle$, a representation of $M_{\eta \smallfrown \langle \ell \rangle}$, usually in a way predetermined simply, except when $\alpha \in S_{\ell g(\eta)}$ and $\ell = 1$, and then we consult a weak diamond sequence. This is like 1.6(1), but there we use our understanding of models in K_λ to build many models in K_{λ^+} while here we build models in $K_{\lambda^{++}}$, thus getting $2^{\lambda^{+2}}$ models in λ^{+2} . We even get $2^{(\lambda^{+2})}$ models in $K_{\lambda^{+2}}$ with none $\leq_{\mathfrak{K}}$ -embeddable into any other.

A second proof 3.23 is like 1.4(1) in the sense that we get only close to $2^{\lambda^{++}}$ models. It is similar to [Sh 87b, 6.4], and the parallel to [Sh 87b, 6.3] holds here. So we have to find an analog of [Sh 87b, definition 6.5, 6.7]. But there we use fullness on the side (meaning: $M \in K_\lambda$ is full over $N \in K_\lambda$ if $N \leq_{\mathfrak{K}} M$, and $(M, c)_{c \in N}$ is saturated), but we do not have this yet.

We still have not explained the framework of this section. In 3.1 - 3.5 we present construction frameworks **C**, which involve sequences of models of length $\leq \lambda$ each of cardinality $< \lambda$ and in particular, define local and nice **C**. In our applications here λ^+ plays the role of λ (and $< \lambda^+$ is specialized to λ).

Then in 3.6 - 3.8 we present examples of such frameworks. Our intention is to use the limit of a sequence $\langle M_\alpha : \alpha < \lambda \rangle$ as an approximation to a model of cardinality λ^+ . For this we define in 3.10 - 3.11 a successor relation (next approximation), modulo a “ $< \lambda$ -amalgamation choice function”; this is denoted $\bar{M}^1 \leq_{F_1}^{\text{at}} \bar{M}^2$. Iterating it we get the quasi order $\leq_F$ (see 3.13). In 3.14 we define the key coding properties, (of an amalgamation choice function F for the framework **C**). The intention is that these coding properties suffice to build many non-isomorphic models in λ^+ . In 3.17 we give the “atomic step” for this construction.

In 3.18 we prove the existence of 2^{λ^+} non-isomorphic models, using the λ -coding property. As we do not have this in some applications we have in mind, we next turn to the weak λ -coding property in 3.19 as well as the weak (local) λ -coding property and corresponding properties of F (all in Definition 3.20, 3.22), connect

them 3.21, and prove there are many models in 3.23.

Lastly, 3.25, 3.28 deal with our concrete case: if the minimal triples in $\mathfrak{K}_\lambda^3$ are not dense then in most cases failures of amalgamation lead to the λ^+ -coding property and hence to many models in cardinality λ^{++} .

Note generally that we mainly axiomatize the construction of models in λ^+ , not how we get $\bar{M}', \bar{M} \leq_{F,a}^{\text{at}} \bar{M}' \in \mathbf{Seq}_\lambda$ that is coding properties; for the last point, see the examples just cited.

Later, in 6.10, we shall need again to use the machinery from this section, in trying to prove that there are enough cases of disjoint amalgamation in $\mathfrak{K}_\lambda$.

We may want to turn the framework presented here into a more general one; see more in [Sh 600].

3.1 Context.

- 1) $\mathfrak{K}$ is an abstract elementary class.
- 2) But $=_M$ or $=_{\mathfrak{K}}$ is just an equivalence relation, i.e. for $M \in K$, $=^M$ is an equivalence relation on $|M|$, moreover a congruence relation relative to all relations and functions in $\tau(M)$ that is for $R \in \tau(M)$ an n -ary relation, we have

$$\bigwedge_{\ell < n} a_i =^M b_i \Rightarrow \langle a_0, \dots, a_{n-1} \rangle \in R^M \equiv \langle b_0, \dots, b_{n-1} \rangle \in R^M.$$

We let $\|M\| = |(M)/=^M|$ and

$$K_{\lambda,\mu} = \{M : |M| \text{ has } \mu \text{ elements and } \|M\| \text{ has } \lambda \text{ elements}\}.$$

- 3) Now the meaning of $\leq_{\mathfrak{K}}$ should be clear but $M <_{\mathfrak{K}} N$ means $(M \in K, N \in K$ and) $M \subseteq N$ and $M/ =^M \leq_{\mathfrak{K}} N/ =^N$ and $\exists a \in N[a/ =^N \notin (M/ =^N)]$ i.e. $(\exists a \in N)(\forall b \in M)(\neg a =^N b)$.
- 4) $K_\lambda^3 = \{(M, N, a) : M \leq_{\mathfrak{K}} N \text{ are from } K_{\lambda,\lambda} \text{ and } a \in N, (a/ =^N \notin (M/ =^N))\}$.
- 5) In this content “ R is an isomorphism relation from M_1 onto M_2 if

- (a) $R \subseteq M_1 \times M_2$
- (b) $a_1 =^{M_1} b_1 \ \& \ a_2 =^{M_2} b_2 \Rightarrow a_1 R a_2 \leftrightarrow b_1 R b_2$
- (c) $(\forall x \in M_1)(\exists y \in M_2) x R y$
- (d) $(\forall y \in M_2)(\exists x \in M_1) x R y$
- (e) if $Q \in \tau(M_1) = \tau(M_2)$ is an n -place relation and $a_\ell Q b_\ell$ for $\ell = 0, \dots, n-1$ then $(a_0, \dots, a_{n-1}) \in Q^{M_1} \rightarrow (b_0, \dots, b_{n-1}) \in Q^{M_2}$.

3.2 Explanation. The need of 3.1(2) is just to deal with amalgamations which are not necessarily disjoint. If we use disjoint amalgamation, we can omit 3.1(2) below on Definition 3.10, a disappears so F is four place and use K_λ instead of $K_{\lambda,\lambda}$. This is continued in [Sh 600, 2.17]. May be better understood after reading 3.10, after clause (c).

3.3 Definition. Let λ be regular uncountable and $\mathfrak{K}$ an abstract elementary class.

1) A λ -construction framework $\mathbf{C} = \mathbf{C}(\mathfrak{K}^+, \mathbf{Seq}, \leq^*)$ means (we shall use it below with λ^+ playing the role of λ):

- (a) $\tau^+ = \tau^+(\mathfrak{K}^+)$ is a vocabulary extending τ . $\mathfrak{K}^+$ is an abstract elementary class satisfying axioms I,II,III from 0.6 and $M \leq_{\mathfrak{K}^+} N \Rightarrow M \upharpoonright \tau \leq_{\mathfrak{K}} N \upharpoonright \tau$. Furthermore $\mathfrak{K}^+ = \mathfrak{K}_{<\lambda}^+$. As above, equality (in τ) is a congruence relation.
- (b) $\mathbf{Seq} = \bigcup_{\alpha \leq \lambda} \mathbf{Seq}_\alpha$ where for $\alpha \leq \lambda$, $\mathbf{Seq}_\alpha$ is a subset of $\{\bar{M} : \bar{M} = \langle M_i : i < \alpha \rangle, M_i \in \mathfrak{K}^+ \text{ is } \leq_{\mathfrak{K}^+}\text{-increasing continuous}\}$. For $\alpha = \lambda$ we require further that $M/\equiv^M$ has cardinality λ , where $M = \bigcup_{i < \lambda} M_i$.
- (c) $\leq^*$ is a relation on triples x, y, t written $x \leq_t^* y$ for $x, y \in \mathbf{Seq}$ and t a set of pairwise disjoint closed intervals of $\ell g(x)$.

We require:

- (d) $\mathbf{Seq}$ is closed under isomorphism and initial segments.
- (e) If $\bar{M}^1 \leq_t^* \bar{M}^2$ and $\gamma \in \cup t$ then $M_\gamma^1 \leq_{\mathfrak{K}^+} M_\gamma^2$ and hence $M_\gamma^1 \upharpoonright \tau \leq_{\mathfrak{K}} M_\gamma^2 \upharpoonright \tau$.
- (f) If $\bar{M}^1 \leq_t^* \bar{M}^2$, $s \subseteq t$, and $\bar{M}^2 \leq \bar{M}^3 \in \mathbf{Seq}$ then $\bar{M}^1 \leq_s^* \bar{M}^3$.
- (g) If t is a set of closed pairwise disjoint intervals of $\ell g(\bar{M})$ and $\bar{M} \in \mathbf{Seq}$ then $\bar{M} \leq_t^* \bar{M}$.

3.4 Convention/Definition:

- 1) From now on $\mathbf{C}$ will be a λ -construction framework.
- 2) If $\bar{M} \in \mathbf{Seq}_\lambda$ then we let $\bar{M} = \langle M_i : i < \lambda \rangle$ and $M =: \bigcup_{i < \lambda} M_i$; similarly with $\bar{M}^x = \langle M_i^x : i < \lambda \rangle$.
- 3) $K_\lambda^{\text{qr}} = \{(\bar{M}, \mathbf{f}) : \bar{M} \in \mathbf{Seq}_\lambda \text{ and } \mathbf{f} : \lambda \rightarrow \lambda\}$.
- 4) If $(\bar{M}^\ell, \mathbf{f}^\ell) \in K_\lambda^{\text{qr}}$ for $\ell = 1, 2$ then $(\bar{M}^1, \mathbf{f}^1) \leq (\bar{M}^2, \mathbf{f}^2)$ means that: for some club E of λ , we have

- (a) $\delta \in E \Rightarrow \mathbf{f}^1(\delta) \leq \mathbf{f}^2(\delta)$ and
- (b) $\bar{M}^1 \leq_t^* \bar{M}^2$ where $t = t_{E, \mathbf{f}^1} = \{[\delta, \delta + \mathbf{f}^1(\delta)] : \delta \in E\}$.

5) $\mathbf{Seq}^s = \{\bar{M} \in \mathbf{Seq} : \bigcup_i |M_i| \text{ is a set of ordinals } < \lambda^+\}$; similarly for $\mathbf{Seq}_\alpha$.

6) $K_\lambda^{\text{qs}} = \{(\bar{M}, \mathbf{f}) \in K_\lambda^{\text{qr}} : \bar{M} \in \mathbf{Seq}_\lambda^s\}$.

7) $\mathbf{C}$ is local (respectively, revised local) if (a-c) following hold:

- (a) $\bar{M} = \langle M_i : i < \alpha \rangle \in \mathbf{Seq}_\alpha$ iff:
 - (α) $\bar{M}$ is $\leq_{\mathfrak{K}^+}$ -increasing continuous in $\mathfrak{K}_{<\lambda}^+$
 - (β) $i + 1 < \alpha \Rightarrow \langle M_i, M_{i+1} \rangle \in \mathbf{Seq}_2$
 - (γ) if $\alpha = \lambda$ then $|M/\equiv^M| = \lambda$, (recall $M = \bigcup_{i < \lambda} M_i$)

- (b) for $\bar{M}^1, \bar{M}^2 \in \mathbf{Seq}$ and t a set of pairwise disjoint closed intervals contained in $\ell g(\bar{M}^1)$ we have:

$$\bar{M}^1 \leq_t^* \bar{M}^2 \text{ iff } [\gamma_1, \gamma_2] \in t \text{ implies}$$

- (α) $\gamma \in [\gamma_1, \gamma_2] \Rightarrow M_\gamma^1 \leq_{\bar{\kappa}^+} \bar{M}_\gamma^2$
 (β) in the local case: $\gamma \in [\gamma_1, \gamma_2] \Rightarrow \langle M_\gamma^1, M_{\gamma+1}^1 \rangle \leq_{\{[0,1]\}}^* \langle M_\gamma^2, M_{\gamma+1}^2 \rangle$;
 in the revised local case: if $\ell g(\bar{M}^1), \ell g(\bar{M}^2) < \lambda$ then
 $\gamma \in [\gamma_1, \gamma_2] \Rightarrow \langle M_\gamma^1, M_\gamma^1 \rangle \leq_{\{[0,1]\}}^* \langle M_\gamma^2, M_\gamma^2 \rangle$, and generally for some club E of λ , $\gamma \in [\gamma_1, \gamma_2] \ \& \ \gamma < \delta \in E \Rightarrow \langle M_\gamma^1, \bigcup_{\beta < \delta} M_\beta^1 \rangle \leq_{\{[0,1]\}}^* \langle M_\gamma^2, \bigcup_{\beta < \delta} M_\beta^2 \rangle$ (and if $\ell g(\bar{M}^\ell) = \alpha_\ell < \delta$, $\bigcup_{\beta < \delta} M_\beta^\ell$ means $\bigcup_{\beta < \alpha_\ell} M_\beta^\ell$).

- (c) if $\langle M_0^\zeta, M_1^\zeta \rangle \in \mathbf{Seq}_2$ for $\zeta < \zeta^* < \lambda$, $\langle M_0^\zeta : \zeta \leq \zeta^* \rangle$ and $\langle M_1^\zeta : \zeta \leq \zeta^* \rangle$ are $\leq_{\bar{\kappa}^+}$ -increasing continuous, and $\zeta < \zeta^* \Rightarrow \langle M_0^\zeta, M_1^\zeta \rangle \leq_{\{[0,1]\}}^* \langle M_0^{\zeta+1}, M_1^{\zeta+1} \rangle$
then $\langle M_0^0, M_1^0 \rangle \leq_{\{[0,1]\}}^* \langle M_0^{\zeta^*}, M_1^{\zeta^*} \rangle \in \mathbf{Seq}$.

So intervals $[\alpha, \alpha] \in t$ are irrelevant for the local version. In the revised local version it is natural to add monotonicity for $\leq_{\{[0,1]\}}$.

8) For $\alpha \leq \lambda$ we say $\mathbf{C}$ is closed for α if:

- (α) $\bar{M} = \langle M_i : i < \alpha \rangle \in \mathbf{Seq}$ iff $\beta < \alpha \Rightarrow \bar{M} \upharpoonright (\beta + 1) \in \mathbf{Seq}$
 (β) if $\bar{M}^\ell = \langle M_i^\ell : i < \alpha_\ell \rangle \in \mathbf{Seq}$ for $\ell = 1, 2$ and $\alpha = \alpha_1 < \alpha_2$, then

$$\bar{M}^1 \leq_t^* \bar{M}^2 \Leftrightarrow \bar{M}^1 \leq_t^* \bar{M}^2 \upharpoonright \alpha.$$

9) $\mathbf{C}$ is disjoint if: $\bar{M}^1 \leq_t^* \bar{M}^2$, $[\gamma_1, \gamma_2] \in t$, $\gamma \in [\gamma_1, \gamma_2]$ implies $M_\gamma^1 = M_{\gamma+1}^1 \cap M_\gamma^2$.

$\mathbf{C}$ is truly disjoint if: $\bar{M}^1 \leq_t^* \bar{M}^2$, $[\gamma_1, \gamma_2] \in t$, $\gamma \in [\gamma_1, \gamma_2]$ implies $M_\gamma^1 = M^1 \cap M_\gamma^2$.

10) In K_λ^{qr} , we say $(\bar{M}, \mathbf{f})$ is a m.u.b. (minimal upper bound) of $\langle (\bar{M}^\xi, \mathbf{f}^\xi) : \xi < \delta \rangle$ if

- (a) $\xi < \delta \Rightarrow (\bar{M}^\xi, \mathbf{f}^\xi) \leq (\bar{M}, \mathbf{f})$ and
 (b) for any $(\bar{M}, \mathbf{f}')$ satisfying (a), for some club E of λ we have: if $\alpha \in E$ and $j \leq \mathbf{f}(\alpha)$ then $\mathbf{f}(\alpha) \leq \mathbf{f}'(\alpha)$ and $M_{\alpha+j} \leq_{\bar{\kappa}^+} M'_{\alpha+j}$.

When we require an increasing sequence in K_λ^{qr} to be continuous we mean that a m.u.b. is used at limits.

11) We say $\mathbf{C}$ is explicitly local if it is local and

- (d) if $\zeta^* < \lambda$ is a limit ordinal, $\langle M_0^\zeta, M_1^\zeta \rangle \in \mathbf{Seq}_2$ for $\zeta \leq \zeta^*$ and for $\ell = 0, 1$ the sequence $\langle M_\ell^\zeta : \zeta < \zeta^* \rangle$ is $\leq_{\bar{\kappa}}$ -increasing continuous, $M_\ell^\zeta \leq_{\bar{\kappa}^+} M_\ell^{\zeta^*}$, and $\zeta < \xi \leq \zeta^* \Rightarrow \langle M_0^\zeta, M_1^\zeta \rangle \leq_{\{[0,1]\}}^* \langle M_0^\xi, M_1^\xi \rangle$ then $\langle \bigcup_{\zeta < \zeta^*} M_0^\zeta, \bigcup_{\zeta < \zeta^*} M_1^\zeta \rangle \leq_{\{[0,1]\}}^* \langle M_0^{\zeta^*}, M_1^{\zeta^*} \rangle$.

12) $\mathbf{C}$ is closed if it is closed for every ordinal $\leq \lambda$.

13) $\mathbf{C}$ is semi (respectively almost) closed, as witnessed by G , if:

- (a) $\mathbf{C}$ is closed for every limit ordinal $\delta < \lambda$;
- (b) G is a function from $\mathbf{Seq}_{<\lambda}$ to $\mathbf{Seq}_{<\lambda}$ such that $\bar{M} \triangleleft G(\bar{M})$;
- (c) $\bar{M} = \langle M_\alpha : \alpha < \lambda \rangle$ belongs to $\mathbf{Seq}_\lambda$ if $\bar{M}$ obeys G , which means: $\beta < \lambda \Rightarrow \bar{M} \restriction \beta \in \mathbf{Seq}_\beta$ and $\{\alpha < \lambda : G(\bar{M} \restriction \alpha) \triangleleft \bar{M}\}$ is unbounded in λ
- (d) in the almost closed version, we add: $G(\bar{M})$ depends on $\bigcup \bar{M}$ only.

14) $\mathbf{C}$ is λ -nice if

- (a) $\leq$ is a transitive on K_λ^{qr} ;
- (b) any increasing continuous sequence in K_λ^{qr} of length $< \lambda^+$ has a m.u.b.; (see part (10)) (not necessarily unique)
- (c) $\mathbf{C}$ is closed (see part (12)).

15) $\mathbf{C}$ is almost λ -nice (as witnessed by G) is defined similarly, replacing “closed” by “almost closed” (witnessed by G).

3.5 Claim. *Let $\mathbf{C}$ be a local (or revised local) λ -construction framework.*

1) *If $(\bar{M}^\ell, \mathbf{f}^\ell) \in K_\lambda^{qr}$ for $\ell = 1, 2$ and E is a club of λ^+ and $(*)$ below then $(\bar{M}^1, \mathbf{f}^1) \leq (\bar{M}^2, \mathbf{f}^2)$ when*

- (*) if $\delta \in E$ then $M_{\delta+i}^1 = M_{\delta+i}^2$ for $i \leq \mathbf{f}^1(\delta)$ and $\mathbf{f}^1(\delta) \leq \mathbf{f}^2(\delta)$; in the “revised local” version assume in addition that $M^1 = M^2$.*

2) *$\leq$ is transitive and a reflexive relation on K_λ^{qr} .*

3) *Any increasing continuous sequence of pairs from K_λ^{qr} of length $< \lambda^+$ has a minimal upper bound.*

4) *If in addition $\mathbf{C}$ is explicitly local (see Definition 3.4(11)) then any increasing sequence in K_λ^{qr} of length $< \lambda^+$ has a lub.*

5) *$\mathbf{C}$ is λ -nice (hence in particular, λ -closed).*

Proof. 1) Check clause (b) of Definition 3.4(7) and Definition 3.4(a).

2) Use clauses (b,c) of Definition 3.4(7). In (c) take $\zeta^* = 2$

3) Without loss of generality the elements of the sequence are $(\bar{M}^\xi, \mathbf{f}^\xi) \in K_\lambda^{qr}$ for $\xi < \mu$, where μ is a regular cardinal $\leq \lambda$. For $\xi < \zeta < \mu$, let $E_{\xi, \zeta}$ be a closed unbounded subset of λ exemplifying Definition 3.4(4) for $(\bar{M}^\xi, \mathbf{f}^\xi) \leq (\bar{M}^\zeta, \mathbf{f}^\zeta)$. First assume $\mu < \lambda$. Let $E \subseteq \bigcap_{\xi < \zeta < \mu} E_{\xi, \zeta} \subset \lambda$ be a closed unbounded subset of λ , such

that: $\alpha \in E \Rightarrow \alpha + (\sup_{\xi < \mu} \mathbf{f}^\xi(\alpha)) + 1 < \text{Min}(E \setminus (\alpha + 1))$. Let $E = \{\alpha_i : i < \lambda\}$ with

α_i increasing continuously with i . Notice that for every $i, \xi < \zeta < \mu \Rightarrow \mathbf{f}^\xi(\alpha_i) \leq \mathbf{f}^\zeta(\alpha_i)$. Let $E^* = \{i : \alpha_i = i\}$.

We now define $\bar{M} = \langle M_j : j < \lambda \rangle$ by defining M_j by induction on j . If $j = \alpha_j \in E^*$ let $M_j = \bigcup_{\xi < \mu} M_j^\xi$. If $\alpha < j \leq \alpha + \mathbf{f}^\xi(\alpha)$ for some $\alpha \in E^*$ and some $\xi < \mu$, let

$\xi' = \sup(\xi : \alpha + \mathbf{f}^\xi(\alpha) \leq j)$ and set $M_j = \bigcup_{\xi > \xi'} M_j^\xi$. If $j = \sup_{\xi}(\alpha + \mathbf{f}^\xi(\alpha))$ for some $\alpha \in E$ and $j > \alpha + \mathbf{f}^\xi(\alpha)$ for each $\xi < \mu$. Let $M_j = \bigcup_{\beta < j} M_\beta$. Finally, if $j < \lambda$ does not fall under any of the previous cases, let $M_j = \bigcup_{\xi < \mu} M_{\alpha_j}^\xi$.

We claim that $\bar{M} \in \mathbf{Seq}_\lambda$. One checks that $\bar{M}$ is continuous and increasing, the main point being that if $\alpha \in E^*$ and $\alpha < j_1 < \alpha + f^{\xi_1}(\alpha) \leq j_2 < \alpha + f^{\xi_2}(\alpha)$ with $\xi_1 < \xi_2 < \mu$, then $M_{j_1}^{\xi_1} \leq M_{j_1}^{\xi_2} \leq M_{j_2}^{\xi_2}$. One must also check that $\langle M_j, M_{j+1} \rangle \in \mathbf{Seq}_2$ for all j . This follows from clause (c) of Definition 3.4(7).

Let $\mathbf{f}$ be defined by $\mathbf{f}(\alpha_i) = \sup\{\mathbf{f}^\xi(\alpha_i) : \xi < \mu\}$ if $i \in E^*$ and $\mathbf{f}(\alpha_i) = 0$ otherwise. Clearly $(\bar{M}^\xi, \mathbf{f}^\xi) \leq (\bar{M}, \mathbf{f})$ for $\xi < \mu$.

What about being a $\leq$ -m.u.b.? Assume that $(\bar{M}', \mathbf{f}') \in K_\lambda^{qr}$ and $\xi < \mu \Rightarrow (\bar{M}^\xi, \mathbf{f}^\xi) \leq (\bar{M}', \mathbf{f}')$. So for each $\xi < \mu$ some club E'_ξ of λ exemplifies Definition 3.4(4), and let $E' =: \bigcap_{\xi < \mu} E'_\xi \cap E^*$, a club of λ .

Now for $\delta \in E'$ we have $(\forall \xi < \mu)(\mathbf{f}^\xi(\delta) \leq \mathbf{f}'(\delta))$ hence $\mathbf{f}(\delta) = \sup_{\xi < \mu} \mathbf{f}^\xi(\delta) \leq \mathbf{f}'(\delta)$, so $\delta \in E' \Rightarrow \mathbf{f}(\delta) \leq \mathbf{f}'(\delta)$. Similarly $\delta \in E' \ \& \ j \leq \mathbf{f}(\delta) \Rightarrow M_{\delta+j} \leq_{\mathfrak{K}} M'_{\delta+j}$. So clearly $(\bar{M}, \mathbf{f}) \leq (\bar{M}', \mathbf{f}')$ and $(\bar{M}, \mathbf{f})$ is a minimal u.b. (see Definition 3.4(10)).

4) As in the proof of part (3), let $\langle (\bar{M}^\xi, \mathbf{f}^\xi) : \xi < \mu \rangle$ be as therein and let $(\bar{M}, \mathbf{f})$ be constructed as above. For proving it is a lub, let $(M^\xi, \mathbf{f}^\xi) \leq (\bar{M}', \mathbf{f}')$ for $\xi < \mu$, and define E' as there. For $\delta \in E', j < \mathbf{f}(\delta)$ we have $\langle (M_{\delta+j}^\xi, M_{\delta+j+1}^\xi) : \xi \in (\xi_{\delta, j+1}, \mu) \rangle$ is $\leq_{\{[0,1]\}}^*$ -increasing continuous and $\langle M_{\delta+j}^\xi, M_{\delta+j+1}^\xi \rangle \leq_{\{[0,1]\}}^* (M'_{\delta+j}, M'_{\delta+j+1})$ for $\xi \in (\xi_{\delta, j+1}, \mu)$, so as $\mathbf{C}$ is explicitly local by clause (d) in Definition 3.4(11) we have

$$\langle M_{\delta+j}, M_{\delta+j+1} \rangle = \left\langle \bigcup_{\xi \in (\xi_{\delta, j+1}, \mu)} M_{\delta+j}^\xi, \bigcup_{\xi \in (\xi_{\delta, j+1}, \mu)} M_{\delta+j+1}^\xi \right\rangle \leq_{\{[0,1]\}} (M'_{\delta+j}, M'_{\delta+j+1})$$

is as required.

The proof for the case $\mu = \lambda$ is similar, using diagonal intersection.

5) Left to the reader. □_{3.5}

* * *

It may clarify matters if we introduce some natural cases of $\mathbf{C}$. We shall use the forthcoming $\mathbf{C}^0$ in our construction of many models in $\mathfrak{K}_{\lambda+2}$.

3.6 Definition. For $\ell \in \{0, 1, 2\}$ and $\lambda = \text{cf}(\lambda) > LS(\mathfrak{K})$, let $\mathbf{C} = \mathbf{C}_{\mathfrak{K}, \lambda}^\ell$ consist of

- (a) $\tau^+ = \tau$,
 $\mathfrak{K}^+ = \{M \in \mathfrak{K}_{<\lambda} : \text{if } \lambda \text{ is a successor cardinal then } \|M\|^+ = \lambda\}$
(with $=^M$ being equality)
- (b) $\mathbf{Seq}_\alpha = \{\bar{M} : \bar{M} = \langle M_i : i < \alpha \rangle \text{ is a } \leq_{\mathfrak{K}}\text{-increasing continuous sequence of members of } \mathfrak{K}_{<\lambda} \text{ and if } \alpha = \lambda \text{ then } \bigcup_{i < \alpha} M_i \text{ has cardinality } \lambda\}$
- (c) $\bar{M} <^*_t \bar{N}$ when:
 - (α) $\bar{M} = \langle M_i : i < \alpha^* \rangle, \bar{N} = \langle N_i : i < \beta^* \rangle$ are from $\mathbf{Seq}$

- (β) if $[\gamma_1, \gamma_2] \in t$ then:
- (i) $\gamma \in [\gamma_1, \gamma_2] \Rightarrow M_\gamma \leq_{\mathfrak{K}} N_\gamma$
 - (ii) if $\ell = 1$, then in addition $\gamma \in [\gamma_1, \gamma_2] \Rightarrow M_\gamma = M_{\gamma+1} \cap N_\gamma$
 - (iii) if $\ell = 2$, then in addition
 $\gamma \in [\gamma_1, \gamma_2] \Rightarrow M_\gamma = M \cap N_\gamma$ where $M = \bigcup_{i < \alpha^*} M_i$.

Note that $\mathbf{C}_{\mathfrak{K}, \lambda}^1$ is interesting when we have disjoint amalgamation in the appropriate cases.

3.7 Fact.

- 1) If $\ell = 0$ or 1 , then $\mathbf{C}_{\mathfrak{K}, \lambda}^\ell$ is an explicitly local λ -construction framework, (hence λ -nice by 3.5(5)) and $\mathfrak{K}^+$ satisfies axioms I-VI.
- 2) If $\ell = 0$ or 2 , then $\mathbf{C}_{\mathfrak{K}, \lambda}^\ell$ is an explicitly revised local λ -construction framework (hence λ -nice by 3.5(5)) and $\mathfrak{K}^+$ satisfies axioms I-VI.

3.8 Definition. If $\lambda = \text{cf}(\lambda) > LS(\mathfrak{K})$ then $\mathbf{C}_{\mathfrak{K}, \lambda}^3$ consists of

- (a) $\tau^+ = \tau \cup \{P, <\}$, $\mathfrak{K}^+$ is the set of $(M, P^M, <^M)$ where $M \in \mathfrak{K}_{<\lambda}$, $P^M \subseteq M$, $<^M$ a linear ordering of P^M (but $=^M$ may be as in 3.1(2)) and $M_1 \leq_{\mathfrak{K}^+} M_2$ iff $(M_1 \upharpoonright \tau) \leq_{\mathfrak{K}} (M_2 \upharpoonright \tau)$ and $M_1 \subseteq M_2$
- (b) $\text{Seq}_\alpha = \{\bar{M} : \bar{M} = \langle M_i : i \leq \alpha \rangle$ is an increasing continuous sequence of members of $\mathfrak{K}^+$ and $\langle M_i \upharpoonright \tau : i \leq \alpha \rangle$ is $\leq_{\mathfrak{K}}$ -increasing, and for $i < j < \alpha : P^{M_i}$ is a proper initial segment of $(P^{M_j}, <^{M_j})$ and there is a first element in the difference}
we denote the $<^{M_{i+1}}$ -first element of $P^{M_{i+1}} \setminus P^{M_i}$, by $a_i[\bar{M}]$
- (d) $\bar{M} <_t^* \bar{N}$ iff
 $\bar{M} = \langle M_i : i < \alpha^* \rangle, \bar{N} = \langle N_i : i < \alpha^{**} \rangle$ are from Seq , t is a set of pairwise disjoint closed intervals of α^* and for any $[\alpha, \beta] \in t$ we have ($\beta < \alpha^*$ and):
 $\gamma \in [\alpha, \beta] \Rightarrow M_\gamma \leq_{\mathfrak{K}} N_\gamma$ & $a_\gamma[\bar{M}] \notin N_\gamma$, moreover
 $a_\gamma[\bar{M}] = a_\gamma[\bar{N}]$.

3.9 Fact. $\mathbf{C}_{\mathfrak{K}, \lambda}^3$ is an explicitly local and revised local λ -construction framework (hence λ -nice by 3.5(5)) and $\mathfrak{K}^+$ satisfies axioms I-VI.

We now introduce amalgamation choice functions. The use of “ F a λ -amalgamation choice function” is to help use the weak diamond, by taking out most of the freedom in choosing amalgams. This gives possibilities for coding (3.14, 3.17).

3.10 Definition. 1) We say that F is a λ -amalgamation choice function for the construction framework $\mathbf{C}$ iff F is a five place function satisfying:

- (a) if $M_\ell \in K_{<\lambda}^+$ for $\ell < 3$, $M_0 \leq_{\mathfrak{K}^+} M_1, M_0 \leq_{\mathfrak{K}^+} M_2, M_1 \cap M_2 = M_0$ (before dividing by $=^{M_\ell}$); $a \in M_2$ and $(\forall b \in M_0)[\neg a =^{M_1} b]$; and A is a set such that $A \cup |M_1| \cup |M_2|$ is a set of ordinals then
 $F(M_0, M_1, M_2, A, a)$, if defined, is a member N of $\mathfrak{K}^+$ with universe
 $A \cup |M_1| \cup |M_2|$, which $\leq_{\mathfrak{K}^+}$ -extends M_1 and M_2 and $(a / =^N) \notin (M_1 / =^N)$;

- (b) [uniqueness]
 if (M_0, M_1, M_2, A, a) and $(M'_0, M'_1, M'_2, A', a')$ are as above and in the domain of F , f is an order preserving mapping from $A \cup |M_1| \cup |M_2|$ onto $A' \cup |M'_1| \cup |M'_2|$ such that $f \upharpoonright M_\ell$ is an isomorphism from M_ℓ onto M'_ℓ for $\ell = 0, 1, 2$ (so preserving $=^{M_\ell}$, and its negation) and $f(a) = a'$ then f is an isomorphism from $F(M_0, M_1, M_2, A, a)$ onto $F(M'_0, M'_1, M'_2, A', a')$;
- (c) if $F(x_0, x_1, x_2, x_3, x_4)$ is well defined then x_0, x_1, x_2, x_3, x_4 are as in part (a).

Observe that as $M_1 \cap M_2 = M_0$ in (a), if we do not have disjoint amalgamation then we are *forced* to allow $=_N$ to be a nontrivial congruence.

2) If F is defined whenever the conditions in part (a) hold and $A \setminus M_1 \setminus M_2$ has large enough cardinality then we say F is full (if $\lambda = \mu^+$, it suffices to demand $A \setminus M_1 \setminus M_2$ has cardinality μ).

3) We say F has strong uniqueness if

- (d) if (M_0, M_1, M_2, A, a) and $(M'_0, M'_1, M'_2, A', a')$ are as above and in the domain of F and for $\ell = 0, 1, 2$ we have R_ℓ is a isomorphism relation R_ℓ from M_ℓ onto M'_ℓ such that $R_0 = R_1 \cap (M_0 \times M'_0) = R_2 \cap (M_0 \times M'_0)$ and $|A| = |A'|$, then there is an isomorphism relation R from $M = F(M_0, M_1, M_2, A, a)$ onto $M' = F(M'_0, M'_1, M'_2, A', a)$ such that $R_\ell = R \cap (M_\ell \times M'_\ell)$.

3.11 Definition. Assume $\mathbf{C}$ is a λ -construction framework and F is a λ -amalgamation choice function for $\mathbf{C}$. Let $(\bar{M}^\ell, \mathbf{f}^\ell) \in K_\lambda^{\text{qr}}$ for $\ell = 1, 2$.

1) $(\bar{M}^1, \mathbf{f}^1) <_{F,a}^{\text{at}} (\bar{M}^2, \mathbf{f}^2)$ (if we omit a , this means for some a ; “at” stands for *atomic* extension; we may write $\leq_{F,a}^{\text{at}}$ instead of $<_{F,a}^{\text{at}}$) means that:

- (a) $(\bar{M}^1, \mathbf{f}^1) \leq (\bar{M}^2, \mathbf{f}^2)$
- (b) for some club E of λ , for every $\delta \in E$ taking $e_\delta =: [\delta, \delta + \mathbf{f}^1(\delta)]$ we have
- (*) if $\beta < \gamma$ are successive members of e_δ then:
 $M_\gamma^2 = F(M_\beta^1, M_\gamma^1, M_\beta^2, |M_\gamma^2|, a)$
- (**) $|M_\gamma^2| = |M_\gamma^1| \cup |M_\beta^2| \cup \{i : i \text{ an ordinal not in } |M_\gamma^1| \cup |M_\beta^2| \text{ and } \text{otp}(|M_\gamma^2| \cap i \setminus |M_\gamma^1| \cup |M_\beta^2|) < \|M_\gamma^2\|\}$
- (***) $\|M_\gamma^2\| = \text{Min}\{\|N\| : N = F(M_\beta^1, M_\delta^1, M_\beta^2, |N|, a)\}$.

A suitable club E may be called a *witness* for the relation. Implicit in clause (b) in $a \in M^2$ and $\neg(\exists b)(b \in M^1 \ \& \ a =^{M^2} b)$.

2) $(\bar{M}', \mathbf{f}') \leq_F (\bar{M}'', \mathbf{f}'')$ means that: there is a sequence $\langle (\bar{M}^\zeta, \mathbf{f}^\zeta) : \zeta \leq \xi \rangle$ such that:

- (a) $\xi < \lambda$
- (b) $(\bar{M}^\zeta, \mathbf{f}^\zeta) \in K_\lambda^{\text{qr}}$ is $\leq$ -increasing continuous in ζ
 (remember Definition 3.4(10))
- (c) for each $\zeta < \xi$ we have $(\bar{M}^\zeta, \mathbf{f}^\zeta) \leq_F^{\text{at}} (\bar{M}^{\zeta+1}, \mathbf{f}^{\zeta+1})$
- (d) $(\bar{M}', \mathbf{f}') = (\bar{M}, \mathbf{f}^0)$ and $(\bar{M}'', \mathbf{f}'') = (\bar{M}^\xi, \mathbf{f}^\xi)$.

A club E which witnesses all the relations in (c) is called a witness for the relation $\leq_F$.

3) $<_{F,a}^{\text{at},*}$ is defined similarly to part (1) but we demand in clause (b) only that $e_\delta \subseteq [\delta, \delta + \mathbf{f}^2(\delta)]$ is closed and $\{\delta, \delta + \mathbf{f}^2(\delta)\} \subseteq e_\delta$; the requirement from clause (a) is unchanged and we require also:

$$\text{if } \beta \in [\delta, \delta + \mathbf{f}^2(\delta)] \text{ then } M_\beta^2 = M_{\max(e_\delta \cap (\beta+1))}^2.$$

Then define $\leq_F^*$ by iterating $\leq_F^{\text{at},*}$.

4) We may replace F by $\mathbf{F}$, a family of such functions. Then in each case in 3.11(2)(c) we use one such that F 's. $\mathbf{F}^*$ is the family of all such F 's.

5) $(\bar{M}^1, \mathbf{f}^1) <_{F,a} (\bar{M}^2, \mathbf{f}^2)$ means that for some $(\bar{M}, \mathbf{f})$ we have $(\bar{M}^1, \mathbf{f}^1) \leq_{F,a}^{\text{at}} (\bar{M}, \bar{\mathbf{f}}) \leq_F (\bar{M}^2, \mathbf{f}^2)$.

6) We define mub as in 3.4(10).

3.12 Remark. 1) What we prove below on $<_{F,a}^{\text{at}}, \leq_F$ also holds for $<_{F,a}^{\text{at},*}, \leq_F^*$.

2) Note: using $\mathbf{F}$ rather than F may help in proving cases of Definition 3.20, but we can use one F which codes all members of $\mathbf{F}$ by asking on $A \setminus M_1 \setminus M_2$, though artificially.

3) We can replace F by $\langle F_\eta : \eta \in \lambda \text{ a sequence of ordinals of length } < \lambda, \eta(1+i) < 2, \eta(0) < 2^{<\lambda} \rangle$, each F_η with uniqueness 3.10(3) and $(**)$ of 3.11(1)(b) is replaced by $M_\gamma^2 = f_{\eta \upharpoonright \delta}(M_\beta^1, M_\gamma^1, M_\beta^2, |M_\gamma^2|, a)$, and omit $(***)$ there.

3.13 Claim. *If $\mathbf{C}$ is nice, then on $K_\lambda^{\text{qr}}, \leq_F$ is a quasi-order, and every increasing continuous sequence of length less than λ^+ has a mub.*

Proof. Check. □_{3.13}

3.14 Definition. 1) We say a λ -amalgamation choice function F for $\mathfrak{R}^+$ has the λ -coding property for $\mathbf{C}$ if: $\text{Seq}_\lambda \neq \emptyset$, and for every $\bar{M}^1 \in \text{Seq}_\lambda$, function $\mathbf{f}^1 : \lambda \rightarrow \lambda$, and $S \subseteq \lambda$ we can find $\bar{M}^{2,\eta} \in \text{Seq}_\lambda$ for $\eta \in {}^\lambda 2$ with η extending $0_{\lambda \setminus S}$ that is $\eta \upharpoonright (\lambda \setminus S)$ being constantly zero, a function $\mathbf{f}^2 : \lambda \rightarrow \lambda$ such that $\mathbf{f}^1 \leq_{\mathcal{D}_\lambda} \mathbf{f}^2$, $\mathbf{f}^2 \upharpoonright (\lambda \setminus S) = \mathbf{f}^1 \upharpoonright (\lambda \setminus S)$ and an element a_η of $M^{2,\eta}$ (usually $a_\eta = a$) such that:

$(*)_1$ $(\bar{M}^1, \mathbf{f}^1) \leq_{F,a}^{\text{at}} (\bar{M}^{2,\eta}, \mathbf{f}^1)$ for all η extending $0_{\lambda \setminus S}$, and

$\eta \upharpoonright \alpha = \nu \upharpoonright \alpha \Rightarrow \bar{M}^{2,\eta} \upharpoonright \alpha = \bar{M}^{2,\nu} \upharpoonright \alpha$;

$(*)_2$ for some club E of λ the following is impossible: for some $\eta_3, \eta_4 \in {}^\lambda 2$ extending $0_{\lambda \setminus S}$, for $\ell = 3, 4$ we have $(\bar{M}^1, \mathbf{f}^2) \leq_{F,a^\ell} (\bar{M}^\ell, \mathbf{f}^\ell)$ witnessed by a club E_ℓ we have (abusing our notation we are dividing by the equality congruence) f_ℓ a $\leq_{\mathfrak{R}}$ -embedding of $M^{2,\eta_\ell} \upharpoonright \tau$ into $M^\ell \upharpoonright \tau$ over M_λ^1 , and for some $\delta \in E \cap E_3 \cap E_4 \cap S$ we have $\bar{M}^3 \upharpoonright \delta = \bar{M}^4 \upharpoonright \delta, a^3 \in M_\delta^{2,\eta_3}, a^4 \in M_\delta^{2,\eta_4}, f_3(a^3) = f_4(a^4), \mathbf{f}^3 \upharpoonright \delta = \mathbf{f}^4 \upharpoonright \delta, f_3 \upharpoonright M_\delta^{2,\eta_3} = f_4 \upharpoonright M_\delta^{2,\eta_4}$, and $\ell \in \{3, 4\} \Rightarrow \text{Rang}(f_\ell \upharpoonright M^{2,\eta_\ell}) \subseteq M_\delta^\ell, \eta_3 \upharpoonright \delta = \eta_4 \upharpoonright \delta, \eta_3(\delta) \neq \eta_4(\delta)$.

2) We say that F has the weak λ -coding property if above we restrict ourselves to the cases $\mathbf{f}^1 \upharpoonright S = 0_S$. We can even restrict ourselves to the cases $\mathbf{f}^1 \in \mathcal{F} \subseteq {}^\lambda \lambda$ provided that $0_\lambda \in \mathcal{F}$ and we demand $\mathbf{f}^2 \in \mathcal{F}$.

- 3) If we replace F by $\mathbf{F}$, a family of such functions it means we use Definition 3.11(4).
 4) We say $\mathfrak{K}$ has a coding property if some λ -amalgamation choice function F has this property. Typically the actual choice of F is irrelevant as long as its domain is sufficiently rich.

3.15 Observation. The restriction above to η such that η extends $0_{\lambda \setminus S}$, S is natural but inessential, as we can extend the definition of $M^{2,\eta}$ to all η in ${}^\lambda 2$ by defining $M^{2,\eta} = M^{2,\eta'}$ where $\eta' \restriction S = \eta \restriction S$ and $\eta' \restriction (\lambda \setminus S)$ is constantly zero. Then the same properties will hold.

3.16 Remark.

- 1) For a local construction framework $\mathbf{C}$ in 3.14(1) the conditions $(*)_1$ and $(*)_2$ can be replaced by local requirements. For example, in condition $(*)_1$, we may take 3.4(7b) into account.
 2) In 3.14 $(*)_2$ a sufficient condition for the impossibility of the stated conditions on E, η^3, η^4 and δ , where $\eta^3 \restriction \delta = \eta^4 \restriction \delta = \eta$, say, is that there is $\bar{a} \in M_{\delta+\mathbf{f}^2(\delta)}^{2,\eta}$ so that:

$$(*)_3 \quad \text{tp}(\bar{a}, M_{\delta+\mathbf{f}^2(\delta)}^1, M^{2,\eta^3}) \neq \text{tp}(\bar{a}, M_{\delta+\mathbf{f}^2(\delta)}^1, M^{2,\eta^4}).$$

We can even allow $\bar{a}$ to be infinite here, say a full listing of $M_{\delta}^{2,\eta}$.

To see that this suffices, suppose that we also have the conditions of 3.14 $(*)_2$. Then for $\ell = 3, 4$ as $M_{\delta+j+1}^{2,\eta_\ell}$ is given by

$$F(M_{\delta+j}^1, M_{\delta+j+1}^1, M_{\delta+j}^{2,\eta_\ell}, |M_{\delta+j+1}^{2,\eta_\ell}|, a^\ell)$$

where $j < \mathbf{f}^2(\delta)$, we find $M_{\delta+\mathbf{f}^2(\delta)}^{2,\eta_3} = M_{\delta+\mathbf{f}^2(\delta)}^{2,\eta_4} = M^*$, say.

Since M^{2,η_3} and M^{2,η_4} can be amalgamated over M^* , we have

$$\text{tp}(f_3(\bar{a}), M_{\delta+\mathbf{f}^2(\delta)}^1, M^{3,\eta_3}) = \text{tp}(f_4(\bar{a}), M_{\delta+\mathbf{f}^2(\delta)}^1, M^{4,\eta_4}).$$

On the other hand by $(*)_2$ we have

$$\text{tp}(\bar{a}, M_{\delta+\mathbf{f}^2(\delta)}^1, M^{2,\eta_\ell}) = \text{tp}(f_\ell(\bar{a}), M_{\delta+\mathbf{f}^2(\delta)}^1, M^{\ell,\eta_\ell})$$

and this gives a contradiction.

3) To understand Definitions 3.14(1,2) you may look at the places they are verified, such as 3.25 and 3.27. Also see 3.21(3,4).

4) The next lemma deduces from the criterion in 3.14(1) another one which is natural for use in a non-structure theorem.

5) Note that 3.14(1) implies: for every $(\bar{M}^1, \mathbf{f}^1)$ there is $(\bar{M}^2, \mathbf{f}^2)$ such that $(\bar{M}^1, \mathbf{f}^1) \leq_{F,a}^{at} (\bar{M}^2, \mathbf{f}^2)$ and $M_\lambda^1 \neq M_\lambda^2$.

6) In Definition 3.14(1) we can replace $<_{F,a}^{at}$ by $<_F$ or $<_{F,a}$ with no harm as $<_{F,a}$ satisfies the requirement on $<_{F,a}^{at}$ and starting from it we again get $<_F$.

7) In $(*)_2$ of Definition 3.14(1) for some function H depending on $(\bar{M}^1, f^1)$ we may add the further restriction $\ell \in \{3, 4\}$ and $\alpha < \delta$ implies $\delta \in H(\bar{M}^\ell \restriction \alpha, f_\ell \restriction M_\alpha^\ell)$. I.e. this weakening of the demand does not change the desired conclusions.

8) We can weaken the demand in $(*)_2$ of 3.14(1) to extensions which actually arise but this seems more cumbersome. While the adaptation is straightforward, we have no application in mind.

9) In 3.14(1), $(*)_2$ we may strengthen the requirement by excluding the case where the club E is allowed to depend on η . That is, we consider quadruples $(E^\eta, \bar{M}^\eta, \mathbf{f}^\eta, f^\eta)$ for $\eta \in {}^\lambda 2$ such that $(\bar{M}^1, \mathbf{f}^2) \leq_F (\bar{M}^\eta, \mathbf{f}^\eta)$ is witnessed by a club E^η in λ and f^η is a $\leq_{\mathfrak{K}}$ -embedding of $M^{2,\eta} \restriction \tau$ into $M^\eta \restriction \tau$ over M^1 . We require:

$(*)'_2$ for no $\eta^3, \eta^4 \in {}^\lambda 2$ and
 $\delta \in E^{\eta^3} \cap E^{\eta^4} \cap E \cap S$ do we have:

$$\eta_3 \restriction \delta = \eta_4 \restriction \delta, \eta_3(\delta) \neq \eta_4(\delta), \mathbf{f}^3 \restriction \delta = \mathbf{f}^4 \restriction \delta;$$

$$\bar{M}^{2,\eta_3} \restriction [\delta, \delta + \mathbf{f}^2(\delta)] = \bar{M}^{2,\eta_4} \restriction [\delta, \delta + \mathbf{f}^3(\delta)];$$

$$f^{\eta_3} \restriction M_\delta^{2,\eta_3} = f^{\eta_4} \restriction M_\delta^{2,\eta_4};$$

$$f^{\eta_\ell} [M_\delta^{2,\eta_\ell}] \subseteq M_\delta^{\eta_\ell} \text{ for } \ell = 3, 4.$$

10) In 3.14 we can also require the models $M^1, M^{2,\eta}$ to have universes $\lambda(1+\alpha)$ and $\lambda(1+\alpha+1)$ respectively for some α , with $\lambda(1+\alpha) \in M_0^{2,\eta}$. This will not change much.

3.17 Lemma. Assume $(\exists \mu)(\lambda = \mu^+ \ \& \ 2^\mu < 2^{\mu^+})$ or at least the definitional weak diamond on λ holds. Assume $\mathbf{C}$ is λ -nice, $J = \text{WDmId}^{\text{Def}}(\lambda)$.

If the λ -amalgamation choice function F has the λ -coding property, then it has the explicit (λ, J) -coding property, which means: if $(\bar{M}^1, \bar{\mathbf{f}}^1) \in K_\lambda^{\text{qs}}$ and $S \subseteq \lambda$ satisfies $S \notin J$ then we can find $(\bar{M}^2, \mathbf{f}^2) \in K_\lambda^{\text{qs}}$ such that:

- (a) $(\bar{M}^1, \mathbf{f}^1) \leq_F^{\text{at}} (\bar{M}^2, \mathbf{f}^1)$ and $\mathbf{f}^1 \restriction (\lambda \setminus S) = \mathbf{f}^2 \restriction (\lambda \setminus S)$
- (b) if $(\bar{M}^1, \mathbf{f}^2) \leq_F (\bar{M}^3, \mathbf{f}^3) \in K_\lambda^{\text{qs}}$ then $M^2 \restriction \tau$ cannot be $\leq_{\mathfrak{K}}$ -embedded into $M^3 \restriction \tau$ over M^1 .

Proof. The proof is straight forward once you digest the meaning of weak diamond.

Let $S = \lambda$. Suppose $\langle \bar{M}^{2,\eta} : \eta \in {}^\lambda 2 \rangle$ and $\mathbf{f}^2$ are as in 3.14(1), taking note of 3.15. Then we claim there is $\nu \in {}^\lambda 2$ for which 3.17 holds on taking $\bar{M}^2$ to be $\bar{M}^{2,\nu}$. Assume toward a contradiction that this fails for each ν . Then clause (b) fails, and for each $\nu \in {}^\lambda 2$ we have some $\bar{M}^{3,\nu}$ and $\mathbf{f}^{3,\nu}$ with:

$$(\bar{M}^1, \mathbf{f}^2) \leq_F (\bar{M}^{3,\nu}, \mathbf{f}^{3,\nu}) \text{ witnessed by a club } E^\nu;$$

$$f_\nu : M^{2,\nu} \restriction \tau \rightarrow M^{3,\nu} \restriction \tau \text{ over } M^1 \restriction \tau \text{ a } \leq_{\mathfrak{K}}\text{-embedding.}$$

Now by the definition of $\text{WDmId}^{\text{Def}}(\lambda)$ we can find $\eta_3 \neq \eta_4$ in ${}^\lambda 2$ and $\delta \in E^{\eta_3} \cap E^{\eta_4} \cap E^* \cap S$ with $E^* = \{\alpha < \lambda \text{ limit: } \beta < \alpha \text{ implies } \beta + \mathbf{f}^2(\delta), \beta + \mathbf{f}^1(\delta) < \alpha\}$, as forbidden in $(*)_2$ of 3.14(1). $\square_{3.17}$

Now we can give a reasonable non-structure theorem.

3.18 Theorem. Assume $\mathbf{C}$ is λ -nice, $(\exists \mu)(\lambda = \mu^+ \ \& \ 2^\mu < 2^{\mu^+}) \ \& \ 2^\lambda < 2^{\lambda^+}$, or at least $DWD(\lambda)$, and $DWD^+(\lambda^+)$. Let $J = WDMId^{\text{Def}}(\lambda)$.

If F has the (λ, J) -coding property, then $I(\lambda^+, \mathfrak{K}) \geq 2^{\lambda^+}$; also if $2^{\lambda^+} > (2^\lambda)^+$ then $IE(\lambda^+, \mathfrak{K}) \geq 2^{\lambda^+}$.

Proof. We choose by induction on $\alpha < \lambda^+$, for every $\eta \in {}^\alpha 2$, a pair $(\bar{M}^\eta, \mathbf{f}^\eta)$ such that:

- (a) $(\bar{M}^\eta, \mathbf{f}^\eta) \in K_\lambda^{\text{qs}}$
- (b) if $\nu \triangleleft \eta$ then $(\bar{M}^\nu, \mathbf{f}^\nu) \leq_F (\bar{M}^\eta, \mathbf{f}^\eta)$
- (c) $(\bar{M}^\eta, \mathbf{f}^\eta) \leq_{F, a_{\eta^\wedge \langle 0 \rangle}}^{\text{at}} (\bar{M}^{\eta^\wedge \langle 0 \rangle}, \mathbf{f}^{\eta^\wedge \langle 0 \rangle})$
- (d) $(\bar{M}^\eta, \mathbf{f}^\eta) \leq_{F, a_{\eta^\wedge \langle 1 \rangle}}^{\text{at}} (\bar{M}^{\eta^\wedge \langle 1 \rangle}, \mathbf{f}^{\eta^\wedge \langle 1 \rangle})$
- (e) if $\ell g(\eta)$ is a limit ordinal and $(\bar{M}^\eta, \mathbf{f}^{\eta^\wedge \langle 0 \rangle}) \leq_F (\bar{M}', \mathbf{f}')$ then
 $M^{\eta^\wedge \langle 1 \rangle} \upharpoonright \tau$ cannot be $\leq_{\mathfrak{K}}$ -embedded into $M' \upharpoonright \tau$ over M^η
- (f) if α is limit ordinal, then $(\bar{M}^\eta, \mathbf{f}^\eta)$ is a $\leq_F$ -mub of $\langle (\bar{M}^{\eta \upharpoonright i}, \mathbf{f}^{\eta \upharpoonright i}) : i < \alpha \rangle$.

For $\alpha = 0$ note that as $\text{Seq}_\lambda \neq \emptyset$ also $\text{Seq}_\lambda^s \neq \emptyset$ hence $K_\lambda^{\text{qs}} \neq \emptyset$. For α limit use 3.6, 3.10. For $\alpha = \beta + 1, \beta$ a limit ordinal and $\nu \in {}^\beta 2$, define $(\bar{M}^{\nu^\wedge \langle \ell \rangle}, \mathbf{f}^{\nu^\wedge \langle \ell \rangle})$ for $\ell = 0, 1$ by 3.17. If $\alpha = \beta + 1, \beta$ non-limit use 3.16(3). Let $M^\eta = \bigcup_{\alpha < \lambda^+} M^{\eta \upharpoonright \alpha} \upharpoonright \tau$ for

$\eta \in (\lambda^+)^2$. Now note $\{a_{\eta \upharpoonright (i+1)} / =^{M^\eta} : i < \lambda^+\} \subseteq M^\eta / =^{M^\eta}$ are pairwise distinct so $M^\eta \in K_{\lambda^+}$. Now we can apply 1.6 (with λ^+ here standing for λ there). $\square_{3.18}$

* * *

Unfortunately, in some interesting cases we get only weak coding.

3.19 Theorem. Assume $\mathbf{C}$ is λ -nice, $(\exists \mu)(\lambda = \mu^+ \text{ and } 2^\mu < 2^\lambda < 2^{\lambda^+} \text{ and } WDMId(\lambda^+) \text{ is not } \lambda^+ \text{-saturated (or at least } DfWD(\lambda), DfWD^+(\lambda^+) \text{ and } WDMId^{\text{Def}}(\lambda) \text{ is not } \lambda^+ \text{-saturated}))$.

If F has the weak λ -coding property (see Definition 3.14(2)), or at least the parallel of the conclusion of 3.17, then $I(\lambda^+, K) \geq 2^{\lambda^+}$.

Proof. We can find $\langle S_\alpha^* : \alpha < \lambda^+ \rangle$ such that:

$$S_\alpha^* \subseteq \lambda$$

$$[\alpha < \beta \Rightarrow |S_\alpha^* \setminus S_\beta^*| < \lambda]$$

and

$$S'_\alpha =: S_{\alpha+1}^* \setminus S_\alpha^* \notin WDMId^{\text{def}}(\lambda).$$

We again choose by induction on $\alpha < \lambda^+$ for every $\eta \in {}^\alpha 2$ a pair $(\bar{M}^\eta, \mathbf{f}^\eta)$ such that:

- (a) $(\bar{M}^\eta, \mathbf{f}^\eta) \in K_\lambda^{\text{qs}}$
- (b) if $\nu \triangleleft \eta$, then $(\bar{M}^\nu, \mathbf{f}^\nu) \leq_F (\bar{M}^\eta, \mathbf{f}^\eta)$
- (c) $f^\eta \upharpoonright (\lambda \setminus S_{\ell g(\eta)}) = 0_{\lambda \setminus S_{\ell g(\eta)}}$
- (d) $(\bar{M}^\eta, \mathbf{f}^\eta) <_{F, a_{\eta^\wedge \langle 0 \rangle}}^{\text{at}} (\bar{M}^{\eta^\wedge \langle 0 \rangle}, \mathbf{f}^{\eta^\wedge \langle 0 \rangle})$
- (e) $(\bar{M}^\eta, \mathbf{f}^\eta) <_{F, a_{\eta^\wedge \langle 1 \rangle}}^{\text{at}} (\bar{M}^{\eta^\wedge \langle 1 \rangle}, \mathbf{f}^\eta)$
- (f) if $(\bar{M}^\eta, \mathbf{f}^{\eta^\wedge \langle 0 \rangle}) \leq_F (\bar{M}^\eta, \mathbf{f}^\eta)$ then $M^{\eta^\wedge \langle 1 \rangle} \upharpoonright \tau$ cannot be $\leq_{\mathfrak{K}}$ -embedded into $M' \upharpoonright \tau$ over M^η
- (g) if α is a limit ordinal, then $(\bar{M}^\eta, \mathbf{f}^\eta)$ is a $\leq_F$ -mub of $\langle (\bar{M}^{\eta \upharpoonright i}, \mathbf{f}^{\eta \upharpoonright i}) : i < \alpha \rangle$.

Again there are no problems (the difference is in clause (c)).

Then we apply 1.6(1) (or 1.7(1)).

□_{3.19}

3.20 Definition. 1) We say a λ -amalgamation choice function F for $\mathbf{C}$ has the weak³ (or weakly) local λ -coding property for $\mathbf{C}$ if:

- (*)₁ Assume $\langle M_0, M_1 \rangle \in \mathbf{Seq}_2$, $M_0 \leq_{\mathfrak{K}^+} N_0 \in \mathfrak{K}_{<\lambda}^+$, $M_1 \cup N_0 \subseteq \lambda^+$, and $|M_1| \cap |N_0| = |M_0|$ and $a \in N_0, a / =^{N_0} \notin (M_0 / =^{N_0})$ (i.e. $(\forall b \in M_0)(\neg a =^{N_0} b)$). Then we can find $N^1, N^2 \in \mathfrak{K}_{<\lambda}^+$ such that:
 - (a) $N^1 = F(M_0, M_1, N_0, |N^1|, a)$
 - (b) $\langle N_0, N^1 \rangle \in \mathbf{Seq}_2$ and $\langle N_0, N^2 \rangle \in \mathbf{Seq}_2$
 - (c) $M_1 \leq_{\mathfrak{K}^+} N^1$ and $M_1 \leq_{\mathfrak{K}^+} N^2$
 - (d) $N^1 \upharpoonright \tau, N^2 \upharpoonright \tau$ are contradicting⁴ amalgamations of $M_1 \upharpoonright \tau, N_0 \upharpoonright \tau$ over $M_0 \upharpoonright \tau$; i.e. for no N', h do we have: $(N^1 \upharpoonright \tau) \leq_{\mathfrak{K}} N' \in \mathfrak{K}_{<\lambda}$ and h is a $\leq_{\mathfrak{K}}$ -embedding of N^2 into N' over $M_1 \cup N_0$; or at least
 - (d)⁻ (N^1, N^2) is a τ -contradicting pair of amalgamations of M_1, N_0 over M_0 which just says: if $N^1 \leq_{\mathfrak{K}^+} N \in \mathfrak{K}_{<\lambda}^1$ then there is no $\leq_{\mathfrak{K}}$ -embedding h of $N^2 \upharpoonright \tau$ into $N \upharpoonright \tau$ over $M_1 \cup N_0$ (i.e. is the identity on M_1 and on N_0)

(note: this is not necessarily symmetric; and we use just the τ -reducts of N^2, M_0, M_1, N_0 so we can replace them by $N^2 \upharpoonright \tau, M_0 \upharpoonright \tau, M_1 \upharpoonright \tau, N_0 \upharpoonright \tau$ respectively).

2) We say that a λ -amalgamation choice function F for $\mathbf{C}$ has the *local* λ -coding property if:

- (*)₂ if $\bar{M} = \langle M_j : j < \lambda \rangle \in \mathbf{Seq}_\lambda, \bar{N} = \langle N_j : j \leq \delta + i \rangle \in \mathbf{Seq}_{\delta+i}, a \in N_0, (a / =^{N_{\delta+i}}) \notin M_{\delta+i} / =^{M_{\delta+i}}$ and $\bar{M} \upharpoonright (\delta + i + 1) \leq_{\{\delta, \delta+i\}}^* \bar{N}$, and $N_{\delta+j+1} = F(M_{\delta+j}, M_{\delta+j+1}, N_{\delta+j}, |N_{\delta+j+1}|, a)$ for $j < i$ then for some $i_1, i_2 \in (i, \lambda)$ and $\bar{N}^\ell = \langle N_\alpha^\ell : \alpha < \delta + i_\ell \rangle \in \mathbf{Seq}_{\delta+i_\ell}$ for $\ell = 1, 2$ we have:
 - (a) $\bar{N}^\ell \upharpoonright (\delta + i + 1) = \bar{N}$ for $\ell = 1, 2$

³there is no clear relation between “weak local” and “local” λ -coding inspite of the name

⁴Of course, we may consider only ones in “legal” extensions. We can also note that for the intended use, the disjointness is automatic (so 3.1(2) not needed)

- (b) for $j \in [\delta + i, \delta + i_1)$ we have
 $N_{j+1}^1 = F(M_j, M_{j+1}, N_j^1, |N_{j+1}^1|, a)$
- (c) $M_{\delta+i_2} \leq_{\mathfrak{R}^+} N_{\delta+i_2}^2$
- (d) $N_{\delta+i_1}^1 \upharpoonright \tau, N_{\delta+i_2}^2 \upharpoonright \tau$ are contradictory amalgamations of
 $M_{\delta+i_1} \upharpoonright \tau$ and $N_\delta \upharpoonright \tau$ over $M_\delta \upharpoonright \tau$ or at least
- (d)⁻ $(N_{\delta+i_1}^1, N_{\delta+i_2}^2)$ are τ -contradictory amalgamations of $M_{\delta+i(*)}$ and N_δ
over M_δ where $i(*) = \min\{i_1, i_2\}$.

(So if $i = 0$, this gives us a possibility to amalgamate, helpful for
 $i \in \lambda \setminus \bigcup_{\delta \in E} [\delta, \delta + \mathbf{f}(\delta))$).

3) We say that a λ -amalgamation choice function F for $\mathbf{C}$ has the *weaker* local λ -coding property for $\mathbf{C}$ if:

(*)₃ as in part (2) but $i = 0$.

4) In 1), 2), 3) above we say $\mathbf{C}$ has weak local or the local or the weaker local coding property respectively, if we omit the mention of F meaning for some F (clause (a) in (*))₁, clause (b) in (*))₂.

3.21 Claim. 1) If F has the weak local λ -coding property for $\mathbf{C}$ or F has the local λ -coding property for $\mathbf{C}$ then F has the weaker local λ -coding property for $\mathbf{C}$.

2) Assume

- (a) $\mathbf{C}$ has a local λ -construction framework
- (b) F has the λ -coding (or weaker λ -coding) (or the weak λ -coding) property for $\mathbf{C}$.

Then for some F' we have:

- (α) F' , too, is a λ -amalgamation choice function for $\mathbf{C}$
- (β) if $F(N_0, N_1, N_2, A, a)$ is well defined and its τ -reduct is $<_{\mathfrak{R}} M \in K_{<\lambda}$ and $A \subseteq A' \subseteq \lambda^+, |A'| < \lambda$, then for some $A'', A' \subseteq A'' \subseteq \lambda^+, |A''| < \lambda$, and $F'(N_0, N_1, N_2, A', a)$ is well defined and $F(N_0, N_1, N_2, A, a) \leq_{\mathfrak{R}^+} F'(N_0, N_1, N_2, A'', a)$ and $M \leq_{\mathfrak{R}} F'(N_0, N_1, N_2, A'', a) \upharpoonright \tau$
- (γ) F' has the local or weak local or weaker local (respectively as in (b)) λ -coding property for $\mathbf{C}$.

3) If $\mathbf{C}$ is local (λ -construction framework), F a λ -amalgamation choice function, with the weak (or just weaker) local λ -coding property, then F has the weak λ -coding property (hence under the set theoretic assumptions of 3.19, $I(\lambda^+, \mathfrak{R}) \geq 2^{\lambda^+}$).

4) If $\mathbf{C}$ is local (λ -construction framework), F a λ -amalgamation choice function, with the local λ -coding property, then F has the λ -coding property (hence under the set theoretic assumptions of 3.18, we have $I(\lambda^+, \mathfrak{R}) \geq 2^{\lambda^+}$ and $(2^\lambda)^+ < 2^{\lambda^+} \Rightarrow IE(\lambda^+, \mathfrak{R}) \geq 2^{\lambda^+}$).

Remark. The parallel of part (2) holds for local and weaker local property if F acts on sequences. See 3.22 below.

Proof. 1) Check.

2) Here we use clause $(d)^-$ rather than (d) .

3) Similar (or read the proof of 3.25).

4) Check. $\square_{3.21}$

In this context we may consider

3.22 Definition. We say that F is a λ -amalgamation choice function for sequences, for $\mathbf{C}$ if:

- (a) if $x = F(x_1, x_2, x_3, x_4, x_5, x_6)$ is defined then for some $\alpha_1, \alpha_2, \alpha_3, \alpha < \lambda$ we have $x_\ell = \bar{M}^\ell \in \text{Seq}_{\alpha_\ell}^s$ for $\ell < 3$, $\bar{M}^1 \triangleleft \bar{M}^2$, $t = x_4$ is a set of pairwise disjoint intervals $\subseteq \alpha_i$, $\bar{M}^1 <_t^* \bar{M}^3$, $A = x_5$ a set of $< \lambda$ ordinals $< \lambda^+$, $x = \bar{M} \in \text{Seq}_\alpha^s$, M_α has universe A , $\bar{M}^1 \leq_{t \cup \{\alpha_1, \alpha_2\}}^* \bar{M}$
- (b) [uniqueness]
as in Definition 3.10.

3.23 Claim. Assume:

- (a) $(\exists \mu < \lambda)(2^\mu = 2^{<\mu} < 2^\lambda), 2^\lambda < 2^{\lambda^+}$ or at least we have the definable weak diamond for λ and λ^+ ; and $2^{<\lambda} \geq \lambda^+$.
- (b) $\mathbf{C}$ is a nice (λ -construction framework).
- (c) F is a λ -amalgamation choice function for $\mathbf{C}$.
- (d) F has the weaker λ -coding property for λ .

Then $I(\lambda^+, \mathfrak{K}) \geq \mu_{\text{wd}}(\lambda^+)$.

Proof. Straight forward using 1.4.

Let $\langle M_i : i < i^* \rangle$ list the models in $\mathfrak{K}_{\lambda^+}$ up to isomorphisms and assume toward contradiction that $i^* < \mu_{\text{wd}}(\lambda^+)$. It is enough to choose by induction on $\alpha < \lambda^+$ a sequence $\bar{M}_0^\eta$ for $\eta \in {}^\alpha 2$ such that:

- (a) $\bar{M}^\eta \in \text{Seq}_\lambda$
- (b) M_λ^η has universe $\gamma_\eta < \lambda^+$
- (c) $\nu \triangleleft \eta \Rightarrow \bar{M}^\nu \leq_F \bar{M}^\eta$
- (d) if $\delta = \ell g(\eta)$ is a limit ordinal then $\bar{M}^\eta$ is a mub of $\langle \bar{M}^\eta \restriction^\alpha : \alpha < \delta \rangle$
- (e) if $\eta \restriction^\alpha \langle 0 \rangle \leq \nu \in {}^{\lambda^+} 2$ then $M_\lambda^{\eta \restriction^\alpha \langle 1 \rangle} \restriction \tau$ cannot be $\leq_{\mathfrak{K}}$ -embedded into $M_\lambda^\nu \restriction \tau$ over $M_\lambda^\eta \restriction \tau$.

This is possible by 3.21(3).

Having the $M_\eta, \eta \in {}^{\lambda^+} 2$ we get the conclusion by 1.5. $\square_{3.23}$

3.24 Remark. 1) If we are just interested in $I(\lambda^+, K)$ rather than also in $IE(\lambda^+, K)$, then we can change the definition of τ -contradictory to:

N^1, N^2 are τ -contradictory amalgamations of M_1, N over M_0 iff $M_0 \leq_{\mathfrak{K}^+} M_1 \leq_{\mathfrak{K}^+} N^\ell, M_0 \leq_{\mathfrak{K}^+} N_0 \leq_{\mathfrak{K}^+} N^\ell, M_0 = M_1 \cap N_0$ and there are no $N_*^1, N_*^2 \in \mathfrak{K}_{<\lambda}^+$ such that: $N^\ell \leq_{\mathfrak{K}^+} N_*^\ell$ and $N_*^1 \restriction \tau, N_*^2 \restriction \tau$ are isomorphic over $M_1 \cup N_0$.

2) Note that it is unreasonable to assume that we will always use the local versions: e.g. if we have a superlimit model in $\mathfrak{K}_{\lambda^+}$ and we want to have $\bar{M} \in \mathbf{Seq}_{\lambda} \Rightarrow M_{\lambda}$ superlimit, we have to add some global condition (see [Sh 600]). Also possibly we will have in the construction (i.e. in 3.18 or 3.19) that $\bar{M}_{\eta}$ has two “contradictory” extensions $\bar{M}_{\eta \smallfrown \langle 0 \rangle}, \bar{M}_{\eta \smallfrown \langle 1 \rangle}$, (see clauses (e) and (f) in their proof) only when $\text{cf}(\ell g(\eta)) = \theta$, where $S = \{\delta : \delta < \lambda^+ \text{ and } \text{cf}(\delta) = \theta\} \notin \text{WdMId}(\lambda)$; or even $\ell g(\eta) \in S$, for a given $S \in \text{WdMId}(\lambda)^+$. We shall deal with such cases when needed.

Remark. We intend to continue this elsewhere.

3.25 Lemma. *Let $\mathfrak{K}$ be an abstract elementary class with $LS(\mathfrak{K}) \leq \lambda$ which is categorical in λ and in λ^+ , with $1 \leq I(\lambda^{++}, K) < 2^{\lambda^{++}}$. Assume that $2^{\lambda} < 2^{\lambda^+} < 2^{\lambda^{++}}$, or at least that the definitional weak diamond holds for both λ^+ and λ^{++} .*

If there is a model in K_{λ^+} which is saturated over λ , then the minimal triples are dense in K_{λ}^3 .

Proof. Let $\mathbf{C}$ be $\mathbf{C}_{\mathfrak{K}, \lambda^+}^0$ (see Definition 3.6) hence $\mathbf{C}$ is explicitly local λ -construction framework (by 3.7(1)). Suppose toward contradiction that above $(M^*, N^*, a^*) \in K_{\lambda}^3$, there is no minimal triple. We claim in this case that there is a λ^+ -amalgamation choice function F for $\mathbf{C}$ with the λ^+ -coding property, with domain the quadruples (M_0, M_1, M_2, A, b) such that: (M^*, N^*, a) embeds in (M_0, M_2, b) ; A and the universes of M_1, M_2 are contained in λ^{++} ; $|A \setminus (|M_1| \cup |M_2|)| = \lambda$; and $M_0 \leq_{\mathfrak{K}} M_1, M_2$. Then applying 3.17 and 3.18 we get a contradiction.

We first make two observations concerning triples (M, N, b) lying above (M^*, N^*, b) . Any such triple has the extension property by 2.11 (or just 2.9(1)) and hence we can manufacture a λ^+ -amalgamation choice function with the specified domain. Furthermore, there is $M' \in K_{\lambda^+}$ with $M \leq_{\mathfrak{K}} M'$ such that $\text{tp}(a, M, N)$ has more than one extension to M' , by the failure of minimality.

Let us show that any λ^+ -amalgamation choice function F with the specified domain has the λ^+ -coding property on $\mathbf{C} = \mathbf{C}_{\mathfrak{K}, \lambda^+}^0$.

Let $\bar{M}^1 \in \mathbf{Seq}_{\lambda^+}$ and let $\mathbf{f}^1 : \lambda \rightarrow \lambda$. For any set S we must find sequences $\bar{M}^{2, \eta}$ (depending on S) as in 3.21(1). The approach will be to first build suitable $M^{2, \eta}$ for all $\eta \in {}^{\lambda}2$, independent of S , then restrict appropriately given S .

Let $M^1 = \bigcup_i M_i^1$. Then $M^1 \in \mathfrak{K}_{\lambda^+}$ and by our assumptions M^1 is therefore saturated over λ . Hence we may suppose $M^* \leq_{\mathfrak{K}} M^1$. We may also suppose that the universe of M^1 is an ordinal, and we may choose a subset A^2 of λ^{++} which is the union of an increasing continuous sequence A_{α}^2 (for $\alpha < \lambda^+$) so that: $A_{\alpha}^2 \cap |M^1| = |M_{\alpha}^1|$ and $A_0^2 \setminus |M_0^1|$ and $A_{\alpha+1}^2 \setminus (A_{\alpha}^2 \cup |M_{\alpha+1}^1|)$ have cardinality λ . Let E be the club:

$$\{\delta < \lambda : \text{for } \alpha < \delta, \text{ we have } \alpha + \mathbf{f}^1(\alpha) + 1 < \delta\}.$$

We now define triples $(M_{\eta}^*, N_{\eta}^*, a^*)$ for all $\eta \in {}^i 2$, by induction on i , together with ordinals α_{η} satisfying the following conditions:

- (a) for any $\eta \in {}^{\lambda}2$, the sequence $(M_{\eta \smallfrown j}^*, N_{\eta \smallfrown j}^*, a^*)(j \leq i)$ is increasing and continuous; and similarly the α_{η} are increasing and continuous.

- (b) $(M_{<>}^*, N_{<>}^*) = (M^*, N^*)$.
- (c) $M_\eta^* = M_{\alpha_\eta}^1$ and the universe of N_η^* is $A_{\alpha_\eta}^2$.
- (d) If $\delta \in E, \eta \in {}^\delta 2$ and $\alpha_\eta = \delta$, then:
 - (d1) for $i < \mathbf{f}^1(\delta)$ and $\eta \triangleleft \nu \in {}^{\delta+i+1}2$, the model N_ν^* is given by F applied to amalgamate $M_{\delta+i+1}^1$ and $N_{\nu \upharpoonright (\delta+i)}^*$ over $M_{\delta+i}^1$, using $A_{\delta+i+1}^2$ and keeping a^* out of $M_{\delta+i+1}^*$
 - (d2) for all $\nu, \nu' \in {}^{\delta+\mathbf{f}^1(\delta)+1}2$ extending η if $\nu' \neq \nu$, then for some β $\text{tp}(a^*, M_\beta^*, N_\nu^*) \neq \text{tp}(a^*, M_\beta^*, N_{\nu'}^*)$
 - (d3) for non-zero $i \leq \mathbf{f}^1(\delta)$ and ν such that $\eta \triangleleft \nu \in {}^{\delta+i}2$ we have $M_\nu^* = M_{\alpha_\nu}^1$ and $\alpha_\nu = \alpha_{\ell g(\nu)} = \alpha_\eta + i = \alpha_\delta + i$ (so $M_\nu^* = M_{\delta+i}^1$)
 - (d4) for non-zero $i \leq \mathbf{f}^1(\delta)$ and $\nu, \rho \in {}^{\delta+i}2$ such that $\eta \triangleleft \nu$ & $\eta \triangleleft \rho$ we have $N_\nu^* = N_\rho^*$ call it N_η^1 .

During carrying the definition the main point is guaranteeing clause (d). So let $\eta \in {}^\delta 2$ and assume that $\alpha_\eta = \delta \in E^*$. First we define by induction on $i \leq \mathbf{f}^1(\delta)$, a model $N_{\eta,i}^1$ such that N_i^1 has universe $A_{\delta+i}^2$, $N_{\eta,i}^1$ is $\leq_{\aleph}$ -increasing, contradiction, $M_{\delta+i}^1 \leq_{\aleph} N_{\eta,i}^1$ and $N_{\eta,0}^1 = N_\eta^*$ and $f, i < \mathbf{f}^1(\delta)$ then

$$N_{\eta,i+1}^1 = F(M_{\delta+i}^1, M_{\delta+i+1}^1, N_{\eta,i}^1, A_{\delta+i+1}^2, a^*).$$

Next we choose by induction on $i \leq \mathbf{f}^1(\delta)$ for each $\rho \in {}^i 2$, an ordinal $\beta_{\eta,\rho} \in [\delta + \mathbf{f}^1(\delta), \lambda^+)$ and model $N_{\eta,\rho}^1$ such that:

- (i) $N_{\eta,\rho}^1 \in K_\lambda$ has universe $A_{\beta_{\eta,\rho}}^2$
- (ii) $M_{\beta_{\eta,\rho}}^1 \leq_{\aleph} N_{\eta,\rho}^1$ and $(a^* / \equiv^{N_{\eta,\rho}^1}) \notin M_{\beta_{\eta,\rho}}^1 / \equiv^{N_{\eta,\rho}^1}$
- (iii) $\beta_{\eta,<>} = \delta + \mathbf{f}^1(\delta)$
- (iv) $\rho_1 \triangleleft \rho_2 \Rightarrow \beta_{\eta,\rho_1} < \beta_{\eta,\rho_2}$ & $N_{\eta,\rho_1}^1 \leq_{\aleph} N_{\eta,\rho_2}^1$
- (v) i limit $\Rightarrow N_{\eta,\rho}^1 = \bigcup_{\zeta < i} N_{\eta,\rho \upharpoonright \zeta}^1$
- (vi) $\beta_{\eta,\rho \wedge <0>} = \beta_{\eta \wedge \rho \wedge <1>}$ and $\text{tp}(a^*, M_{\beta_{\eta,\rho \wedge <0>}}^1, N_{\eta,\rho \wedge <0>}^1) \neq \text{tp}(a^*, M_{\beta_{\eta,\rho \wedge <1>}}^1, N_{\eta,\rho \wedge <1>}^1)$.

There is no problem to carry the definition. Now let

$$\alpha_{\eta \wedge \rho} = \delta + i = \ell g(\eta \wedge \rho) \text{ if } \rho \in {}^i 2 \text{ and } i \leq \mathbf{f}^1(\delta)$$

$$\alpha_{\eta \wedge \rho} = \beta_{\eta,\rho} \text{ if } \rho \in {}^{\mathbf{f}^1(\delta)+1} 2$$

$$M_{\eta \wedge \rho}^* = M_{\alpha_{\eta \wedge \rho}}^1 \text{ if } \rho \in {}^i 2, i \leq \mathbf{f}^1(\delta) + 1$$

$$N_{\eta \wedge \rho}^* = N_{\eta,i}^1 \text{ if } \rho \in {}^i 2 \text{ and } i \leq \mathbf{f}^1(\delta)$$

$$N_{\eta \restriction \rho}^* = N_{\eta, \rho}^1 \text{ if } \rho \in {}^{\mathbf{f}^1(\delta)+1}2.$$

Now check.

Having carried the induction for $\eta \in {}^{\lambda^+}2$ we let $\bar{M}^{2,\eta} = \langle N_{\eta \restriction \alpha}^* : \alpha < \lambda^+ \rangle$ and $\mathbf{f}^2 = \mathbf{f}^1 + 1$. We have to check that the demand in Definition 3.14 holds.

Note that this is essentially the proof mentioned in 3.16(9).

By our initial remarks there is little difficulty in carrying out this induction.

We then set $\bar{M}^{2,\eta} = \langle N_{\eta \restriction i}^* : i < \lambda^+ \rangle$ for $\eta \in {}^{\lambda^+}2$. Given a set $S \subseteq \lambda^+$ we consider $\bar{M}^{2,\eta}$ for $\eta \in {}^{\lambda^+}2$ extending $0_{\lambda \setminus S}$, together with the function $\mathbf{f}^2$ equal to $\mathbf{f}^1 + 1$ on S and to $\mathbf{f}^1$ on S . We claim that the two conditions of Definition 3.21(1) are met.

The first of these is a condition on the type of construction allowed, and, of course, it has been obeyed, notably in (d1) above:

$$(*)_1 \quad (\bar{M}^1, \mathbf{f}^1) \leq_{F,a}^{\text{at}} (\bar{M}^{2,\eta}, \mathbf{f}^1); \text{ and } M^{2,\eta} \restriction \alpha \text{ is determined by } \eta \restriction \alpha.$$

The second condition referred to a club E , which can be the intersection of the club we have defined above with $\{\delta : \alpha_\delta = \delta\}$. This condition goes as follows:

- (*)₂ it is impossible to find sequences η^3, η^4 (extending $0_{\lambda \setminus S}$), extensions $(\bar{M}^1, \mathbf{f}^2) \leq_F (\bar{M}^3, \mathbf{f}^3), (\bar{M}^4, \mathbf{f}^4)$ witnessed by clubs E^3, E^4 (i.e. E^ℓ is the intersection of the clubs which witness the atomic relations $\leq_F^{\text{at}}$ implicit in $\leq_F$), and embeddings $f_\ell : M^{2,\eta^\ell} \rightarrow M^\ell$ ($\ell = 3$ or 4) over M^1 such that for some $\delta \in E \cap E^3 \cap E^4 \cap S$ we have:
- (i) $\bar{M}^3 \restriction (\delta + \mathbf{f}^2(\delta) + 1) = \bar{M}^4 \restriction (\delta + \mathbf{f}^2(\delta) + 1)$;
 - (ii) $\mathbf{f}^3 \restriction \delta = \mathbf{f}^4 \restriction \delta$;
 - (iii) $\eta^3 \restriction \delta = \eta^4 \restriction \delta$ (call the restriction η) and $\eta^3(\delta) \neq \eta^4(\delta)$;
 - (iv) f_3, f_4 are equal on $M_\delta^{2,\eta}$; and
 - (v) for $\ell = 3, 4$, f_ℓ maps $M_\delta^{2,\eta}$ into M_δ^ℓ .

Suppose on the contrary we have η^3, η^4 (extending $0_{\lambda \setminus S}$), $(\bar{M}^1, \mathbf{f}^2) \leq_F (\bar{M}^3, \mathbf{f}^3), (\bar{M}^4, \mathbf{f}^4), E^3, E^4, f_3, f_4$ and δ as above.

Let $\hat{M} = M_{\delta + \mathbf{f}^2(\delta)}^3$. It follows from condition (i) and the fact that δ belongs to the witnessing clubs E^3, E^4 that $\hat{M} = M_{\delta + \mathbf{f}^2(\delta)}^4$. Then f_3, f_4 provide embeddings of $N_{\eta^3 \restriction (\delta + \mathbf{f}^1(\delta) + 1)}^*$ and $N_{\eta^4 \restriction (\delta + \mathbf{f}^1(\delta) + 1)}^*$ into $\hat{M}$ which agrees on N_η^* (hence on a^*) and on M_η^1 . By 3.16(2) we are done. $\square_{3.25}$

3.26 Lemma. *Let $\mathfrak{K}$ be an abstract elementary class with $LS(\mathfrak{K}) \leq \lambda$ which is categorical in λ and in λ^+ , with $1 \leq I(\lambda^{++}, K) < 2^{\lambda^{++}}$. Assume that $2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}}$, or at least that the definitional weak diamond holds for both λ^+ and λ^{++} . Then:*

- (*) *for any $M \in K_{\lambda^+}$ and any triple (M^0, N^0, a^0) in K_λ^3 with $M^0 \leq_{\mathfrak{K}} M$, we can find sequences $\bar{M} = \langle M_i : i < \lambda^+ \rangle, \bar{N} = \langle N_i : i < \lambda^+ \rangle$ such that*

- (a) $(M^0, N^0, a^0) = (M_0, N_0, a^0)$;
- (b) (M_i, N_i, a) is increasing and continuous in K_λ^3 ;
- (c) the union of the M_i is M ;
- (d) the set $S(\bar{M}, \bar{N}, a)$ of $\delta < \lambda^+$ such that for some $j > \delta$ for all $i \geq j$

if we have $(M_j, N_j, a) \leq_\ell (M_i, N^\ell, a)$ for $\ell = 1, 2$ then we can amalgamate (M_i, N^1, a) and (M_i, N^2, a) over (M_i, N_δ, a)

is stationary in λ^+ .

Proof. Otherwise, we claim that any full λ^+ -amalgamation choice function will have the λ^+ -coding property.

Let $\bar{M} \in \mathbf{Seq}_{\lambda^+}$, $\mathbf{f} : \lambda^+ \rightarrow \lambda^+$ and $S \subseteq \lambda^+$ be given. Then as K is categorical in λ^+ , we may suppose that $\bar{M}$ is the union of the M_i . If (c) fails, then (d) fails on a club of δ , providing enough failures of amalgamation to carry the proof as in 3.12.

$\square_{3.12}$

3.27 Lemma. *Let $\mathfrak{K}$ be an abstract elementary class with $LS(\mathfrak{K}) \leq \lambda$ which is categorical in λ, λ^+ and λ^{++} and with no model in cardinality λ^{+3} .*

Suppose that there is no model in K_{λ^+} saturated above λ and that:

(*) *for any $M \in K_{\lambda^+}$ and any triple (M^0, N^0, a^0) in K_λ^3 with $M^0 \leq_{\mathfrak{K}} M$, we can find sequences $\bar{M} = \langle M_i : i < \lambda^+ \rangle, \bar{N} = \langle N_i : i < \lambda^+ \rangle$ such that*

- (a) $(M^0, N^0, a^0) = (M_0, N_0, a^0)$;
- (b) (M_i, N_i, a) is increasing and continuous in K_λ^3 ;
- (c) the union of the M_i is M ;
- (d) the set $S(\bar{M}, \bar{N}, a)$ of $\delta < \lambda^+$ such that for some $j > \delta$ for all $i \geq j$

if we have $(M_j, N_j, a) \leq_{h_\ell} (M_i, N^\ell, a)$ for $\ell = 1, 2$ then we can amalgamate (M_i, N^1, a) and (M_i, N^2, a) over (M_i, N_δ, a)

is stationary in λ^+ .

Then the minimal triples are dense in K_λ^3 .

Proof. Suppose that there is no minimal triple above (M^*, N^*, a^*) . It suffices to show that there is no maximal model in $K_{\lambda^{++}}$ and as $K_{\lambda^{++}}$ is categorical, this will follow from the existence of a single pair of models (M', N') in $K_{\lambda^{++}}$ with $M' <_{\mathfrak{K}} N'$. So it suffices to show:

every triple (M, N, a) in $K_{\lambda^+}^3$ has a proper extension in $K_{\lambda^+}^3$

as the desired pair (M', N') can then be built as the limit of an increasing continuous chain.

Fix (M, N, a) in $K_{\lambda^+}^3$. As there is no model saturated over λ in K_{λ^+} , there is some M_0 in K_λ over which there are more than λ^+ types. By λ -categoricity we may suppose $M_0 \leq_{\mathfrak{K}} M$. Fix a triple (M_0, N_0, b) in K_λ^3 for which $\text{tp}(b, M_0, N_0)$ is not realized in M .

Apply $(*)$ to M and (M_0, N_0, b) to get sequences $\bar{M}, \bar{N}$ of length λ^+ as in $(*)$. Let $S = S(\bar{M}, \bar{N}, b)$. In particular $M = \bigcup M_i$. Let $M^1 = \bigcup N_i$. As $\mathfrak{K}$ has amalgamation in λ^+ we may suppose $M^1, N \leq_{\mathfrak{K}} N^1$ with $N^1 \in \mathfrak{K}_{\lambda^+}$.

We can also choose an increasing continuous sequence (M'_i, N'_i, a^*) beginning with (M^*, N^*, a^*) such that each (M'_i, N'_i, a^*) is reduced and $\text{tp}(a^*, M'_i, N'_i)$ has more than one extension in $\mathcal{S}(M'_{i+1})$, using the failure of minimality and 2.7(1). By categoricity we may suppose $M = \bigcup M'_i$. Set $M^2 = \bigcup N'_i$. By amalgamation we may suppose $M^2, N^1 \leq_{\mathfrak{K}} N^2 \in K_{\lambda^+}$.

We claim that one of the triples (M^1, N^1, a) or (M^2, N^2, a) is a proper extension of (M, N, a) . Suppose on the contrary that a belongs to both M^1 and M^2 .

Represent N^2 as the union of a continuous $\leq_{\mathfrak{K}}$ -increasing chain $\langle N_i^* : i < \lambda^+ \rangle$ of models in K_λ .

Let E be

$$\{i < \lambda^+ : M_i = M'_i; N_i^* \cap M = M_i; N_i^* \cap M^1 = N_i; N_i^* \cap M^2 = N_i^*\},$$

a club in λ^+ .

Fix $\delta \in E \cap S$ such that a is in N_δ and N'_δ . We show now that $a^* \in N_\delta$. Now $(M'_\delta, N'_\delta, a^*)$ is reduced. If $a^* \notin N_\delta$ then (N_δ, N^2, a^*) lies over $(M'_\delta, N'_\delta, a^*)$ and hence $N_\delta \cap N'_\delta \leq M'_\delta$; but the element a witnesses the failure of this condition. So $a^* \in N_\delta$.

Let $j > \delta$ be chosen in accordance with the definition of $S(\bar{M}, \bar{N}, b)$ and let $i > j' > j_a, k, j' \in E$. As $\text{tp}(a^*, M'_{j'}, N'_{j'})$ has more than one extension to $M'_{j'+1}$, the same applies to M'_i . However, $M'_i = M_i$ and $M'_{j'} = M_{j'}$ and thus $\text{tp}(a^*, M_{j'}, N_{j'}^2) = \text{tp}(a^*, M_{j'}, N_{j'}^*) = \text{tp}(a^*, M_{j'}, N_{j'})$ has more than one extension over M_i . Thus, M_i and $N_{j'}$ may be amalgamated in two incompatible ways over $M_{j'}$, getting N^+ and N^- , say (i.e. the N^+ and N^- cannot be amalgamated over M_i preserving the images of a^*). Furthermore $(M_i, N^\pm, b)$ lies above (M_j, N_j, b) in $K_{\lambda^+}^3$, that is, b is not mapped into M_i , because M does not realize $\text{tp}(b, M_0, N_0)$. However, this contradicts the definition of S , as the triples (M_i, N^+, b) and (M_i, N^-, b) ; cannot be amalgamated over (M_i, N_δ) since a^* belongs to N_δ . $\square_{3.27}$

3.28 Theorem. *Let $\mathfrak{K}$ be an abstract elementary class with $LS(\mathfrak{K}) \leq \lambda$ which is categorical in λ and in λ^+ with $1 \leq I(\lambda^{++}, K) < 2^{\lambda^{++}}$. Assume that $2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}}$, or at least that the definitional weak diamond holds for both λ^+ and λ^{++} .*

Then under either of the following assumptions, the minimal triples are dense in K_λ^3 :

- (A) *K is categorical in λ^{++} and has no model in cardinality λ^{+3} ;*
- (B) *there is a model saturated above λ in cardinality λ^+ .*

Proof. By the previous lemmas 3.25, 3.26, 3.27. Note that $(*)$ is exactly the negation of $(*)$. $\square_{3.28}$

3.29 Remark. This will be proved without the additional assumptions (A, B) in [Sh 600]. In any case this does not affect the proof of Theorem 0.5

3.30 Claim. *Let $\mathfrak{K}$ be an abstract elementary class with $LS(\mathfrak{K}) \leq \lambda$ which is categorical in λ and in λ^+ , with $1 \leq I(\lambda^{++}, K) < 2^{\lambda^{++}}$, and with no model in cardinality λ^{+3} . Assume that $2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}}$.*

Then the minimal triples are dense in K_λ^3 .

Proof. If $2^{\lambda^+} > \lambda^{++}$ we get the conclusion by 2.7. If $2^{\lambda^+} = \lambda^{++}$ then as $2^\lambda < 2^{\lambda^+}$ we have $2^\lambda = \lambda^+$. Thus, there is a model in K_{λ^+} which is saturated above λ , and Lemma 3.28 applies. $\square_{3.30}$

§4 MINIMAL TYPES

We return to the analysis of minimal types initiated in 2.12.

We use from §2 only 2.1, 2.6, 2.9 so there are repetitions.

4.1 Hypothesis.

- (a) $\mathfrak{K}$ is an abstract elementary class with $LS(\mathfrak{K}) \leq \lambda$ (for simplicity $K_{<\lambda} = \emptyset$).
- (b) $\mathfrak{K}$ is categorical in λ, λ^+ with $K_{\lambda+2} \neq \emptyset$ (note: $(a) + (b) = (*)_\lambda^3$ of 2.4).
- (c) $\mathfrak{K}$ has amalgamation in λ (2.2(1)),
so by (a) + (c), we have $(*)_\lambda^2$ from 2.9 hence $\mathfrak{K}$ satisfies the model theoretic properties which were deduced in 2.4-2.6, 2.9 in particular:
 - (i) every $(M, N, a) \in K_\lambda^3$ has the weak extension property 2.4;
 - (ii) criteria for the extension property 2.9;
 - (iii) basic Definitions and properties 2.3, 2.6.

4.2 Definition. 1) If $p \in \mathcal{S}(N)$, $N \in K_\lambda$ and $N' \in K_\lambda$ remember (from Definition 2.13)

$$\mathcal{S}_p(N') = \{f(p) : f \text{ is an isomorphism from } N \text{ onto } N'\}$$

and let

$$\mathcal{S}_{\geq p}(N') = \left\{ q \in \mathcal{S}(N') : q \text{ not algebraic (i.e. not realized by any } c \in N') \text{ and, for some } N'' \in K_\lambda, N'' \leq_{\mathfrak{K}} N', \text{ we have } q \upharpoonright N'' \in \mathcal{S}_p(N'') \right\}.$$

2) We say the type $p \in \mathcal{S}(N)$ is λ -algebraic if $\|N\| \leq \lambda$ and for every M such that $N \leq_{\mathfrak{K}} M$ we have: $\lambda \geq |\{c \in M : \text{tp}(c, N, M) = p\}|$.

4.3 Claim. If $(M_0, M_1, a) \in K_\lambda^3$ is minimal, then it has the extension property.

Proof. Let $p^* = \text{tp}(a, M_0, M_1)$, and assume it is a counter-example. We note:

- ⊗₁ for some M^* we have $M_0 \leq_{\mathfrak{K}} M^* \in K_\lambda$ but for no M^+ and b do we have $M^* \leq_{\mathfrak{K}} M^+ \in K_\lambda, b \in M^+ \setminus M^*$ and b realizes p^* .
[Why? If not for every $N, M_0 \leq_{\mathfrak{K}} N \in K_\lambda$, we can find $N_1, N \leq_{\mathfrak{K}} N_1 \in K_\lambda$ and $b \in N_1 \setminus N$ which realizes p^* . Hence (as $\mathfrak{K}_\lambda$ has amalgamation in λ) we can find N_2 such that $N_1 \leq_{\mathfrak{K}} N_2 \in K_\lambda$, and g a $\leq_{\mathfrak{K}}$ -embedding of M_1 into N_2 extending id_{M_0} such that $g(a) = b$. This proves the extension property].

- $\otimes_2$ if $p \in \mathcal{S}_{\geq p^*}(N)$ and $N \in K_\lambda$ and $N \leq_{\mathfrak{K}} N^* \in K$, then the set of elements of $b \in N^*$ realizing p has cardinality $\leq \lambda$.
 [Why? by 4.1(c)(ii); so indirectly 2.9(2)].
- $\otimes_3$ if $N \in K_\lambda$, then $|\mathcal{S}_{\geq p^*}(N)| > \lambda^+$.

Proof of $\otimes_3$. If N forms a counterexample, as K is categorical in λ and using $\otimes_2$ we can find $\langle N_i : i < \lambda^+ \rangle$, $\leq_{\mathfrak{K}}$ -increasing continuous sequence of members of K_λ such that:

- (*) for every $\alpha < \lambda^+$ and $q \in \mathcal{S}_{\geq p^*}(N_\alpha)$, for some $\beta = \beta_q < \lambda^+$ we have: for no N', b do we have $N_\beta \leq_{\mathfrak{K}} N' \in K_\lambda, b \in N' \setminus N_\beta$ and b realizes q .

So $N_{\lambda^+} = \bigcup_{i < \lambda^+} N_i$ has the property

- (**) if $\bar{N}' = \langle N'_\alpha : \alpha < \lambda^+ \rangle$ is a representation of N_{λ^+} then for a club of $\delta < \lambda^+$ for every $q \in \mathcal{S}_{\geq p^*}(N'_\delta)$ for a club of $\beta \in (\delta, \lambda^+)$, for no N', b do we have: $N_\beta \leq_{\mathfrak{K}} N' \in K_\lambda, b \in N' \setminus N_\beta$ and b realizes q .

On the other hand, we can choose by induction on $\alpha < \lambda^+$ a triple $(N_{0,\alpha}, N_{1,\alpha}, a) \in K_\lambda^3$ increasing continuous in α such that $(N_{0,0}, N_{1,0}, a) = (M_0, M_1, a)$ and $N_{0,\alpha} \neq N_{0,\alpha+1}$ (existence by the weak extension property; i.e. 2.4 = 4.1(c)(i)).

Now $N_{0,\lambda^+} = \bigcup_{\alpha < \lambda^+} N_{0,\alpha} \in K_{\lambda^+}$ does not satisfy the statement (**):

$\langle N_{0,\alpha} : \alpha < \lambda^+ \rangle$ is a representation of N_{0,λ^+} , and for every α , $\text{tp}(a, N_{0,\alpha}, N_{1,\alpha})$ extend $\text{tp}(a, N_{0,0}, N_{1,0}) = \text{tp}(a, M_0, M_1) = p^*$ hence $\text{tp}(a, N_{0,\alpha}, N_{1,\alpha}) \in \mathcal{S}_{\geq p^*}(N_{0,\alpha})$ satisfies: for every $\beta \in (\alpha, \lambda^+)$, there is $N', N_{0,\beta} \leq_{\mathfrak{K}} N'$ and some $b \in N' \setminus N_{0,\beta}$ realizes $\text{tp}(a, N_{0,\alpha}, N_{1,\alpha})$; simply choose $(N', b) = (N_{1,\beta}, a)$. So $N_{0,\lambda^+}, N_{\lambda^+}$ cannot be isomorphic (as one satisfies (**) the other not). But both are in K_{λ^+} , contradicting the categoricity of $\mathfrak{K}$ in λ^+ . $\square_{\otimes_3}$

To finish the proof of 4.3 it is enough to prove

4.4 Claim. If $p^* \in \mathcal{S}(M_0)$ is minimal, $M_0 \in K_\lambda$, then $N \in K_\lambda \Rightarrow |\mathcal{S}_{\geq p^*}(N)| \leq \lambda^+$.

Proof. By 4.6, 4.7 below. Note that $\mathcal{S}_{\geq p^*}(N)$ has the same cardinality for every $N \in K_\lambda$.

4.5 Claim. 1) If $N_1 \leq_{\mathfrak{K}} N_2$ are in K_λ and $p_1 \in \mathcal{S}(N_1)$ is minimal and is omitted by N_2 then p_1 has a unique extension in $\mathcal{S}(N_2)$, call it p_2 , and $p_1 \in \mathcal{S}_{\geq p^*}(N_1) \Rightarrow [p_2 \in \mathcal{S}_{\geq p^*}(N_2) \text{ and } p_2 \text{ is minimal}]$.
 2) If $N_1 \leq_{\mathfrak{K}} N_2$ are in $K_\lambda, p_1 \in \mathcal{S}(N_1)$ minimal, then p_1 has a unique non-algebraic extension in $\mathcal{S}(N_2)$ called p_2 , it is minimal and $p_1 \in \mathcal{S}_{\geq p^*}(N_1) \Rightarrow p_2 \in \mathcal{S}_{\geq p^*}(N_2)$.
 3) (Continuity) if $\langle N_i : i \leq \alpha \rangle$ is a $\leq_{\mathfrak{K}}$ -increasing continuous sequence of members

of $K_\lambda, p_i \in \mathcal{S}(N_i), p_0$ minimal, $p_i \in \mathcal{S}(N_i)$ extends p_0 and is non-algebraic then $\langle p_i : i \leq \alpha \rangle$ is increasing continuously.

Proof of 4.5. Easy. E.g.,

3) If $i < j \leq \alpha$ then $p_j \upharpoonright N_i$ is well defined, it belongs to $\mathcal{S}(N_i)$, also it is non-algebraic and extending p_0 hence by the uniqueness (=4.5(1)) we have $p_i = p_j \upharpoonright N_i$. If $\delta \leq \alpha, p_\delta \in \mathcal{S}(N_\delta)$ extends p_i for $i < \delta$; if $p'_\delta \in \mathcal{S}(N_\delta)$ extends each $p_i (i < \delta)$ then it extends p_0 and is non-algebraic hence by uniqueness $p'_\delta = p_\delta$.

4.6 Claim. *If $N \in K_\lambda, \mathcal{S} \subseteq \mathcal{S}(N)$ and $|\mathcal{S}| > \lambda^+$, then we can find N^*, N_i in K_λ , (for $i < \lambda^{++}$) such that:*

- (α) $N \leq_{\mathfrak{K}} N^* <_{\mathfrak{K}} N_i$
- (β) for no $i_0 < i_1 < \lambda^{++}$ and $c_\ell \in N_{i_\ell} \setminus N^*$ (for $\ell = 0, 1$) do we have $tp(c_0, N^*, N_{i_0}) = tp(c_1, N^*, N_{i_1})$
- (γ) there are $a_i \in N_i$ (for $i < \lambda^{++}$) such that $tp(a_i, N, N_i) \in \mathcal{S}$ (and they are pairwise distinct).

Remark. We use here less than Hypothesis 4.1:

- (*) $\mathfrak{K}$ is abstract elementary class with amalgamation in λ , categorical in λ , $K_{\lambda^+} \neq \emptyset$.

The same applies to 4.7.

Proof. Without loss of generality $|N| = \lambda$; now choose by induction on $\alpha < \lambda^{++}, \bar{N}^\alpha, N_\alpha, a_\alpha$ such that:

- (A) $N_\alpha \in K_{\lambda^+}$ has a set of elements $\lambda \times (1 + \alpha)$ and N_α is $\leq_{\mathfrak{K}}$ -increasing continuous in α
- (B) $\bar{N}^\alpha = \langle N_i^\alpha : i < \lambda^+ \rangle$ is a representation of N_α (i.e. is $\leq_{\mathfrak{K}}$ -increasing continuous, $\|N_i^\alpha\| \leq \lambda$ and $N_\alpha = \bigcup_{i < \lambda^+} N_i^\alpha$)
- (C) for $\alpha < \lambda^{++}$ successor, if $i < j < \lambda^+, p \in \mathcal{S}(N_i^\alpha)$ is realized in N_j^α and is λ -algebraic (see Definition 4.2(2)) then for no N', b do we have $N_j^\alpha \leq_{\mathfrak{K}} N' \in K_\lambda$ and $b \in N' \setminus N_j^\alpha$ realizes p (actually not needed)
- (D) $N = N_0^0 \leq_{\mathfrak{K}} N_0$ and $a_\alpha \in N_{\alpha+1} \setminus N_\alpha$ realizes some $p_\alpha \in \mathcal{S}$ not realized in N_α .
- (E) If $\aleph_0 < \text{cf}(\alpha) \leq \lambda$ then let for $j < \lambda$:
 $M_j^\alpha = \cap \{ \bigcup_{\beta \in C} N_j^\beta : C \text{ a club of } \alpha \}$ whenever possible, i.e. the result is in K_λ and $\leq_{\mathfrak{K}} N^\alpha$
- (F) for each $\alpha < \lambda^{++}$, for a club E_α^0 of ordinals $i < \lambda^+$ we have $(N_i^\alpha, N_i^{\alpha+1}, a_\alpha)$ is reduced;
hence (as $tp(a_\alpha, N_i^\alpha, N_i^{\alpha+1})$ extends $tp(a_\alpha, N, N_i^{\alpha+1})$ which is not realized in N_α):

- (G) for every $(i \in E_\alpha^0 \text{ and } b \in N_i^{\alpha+1} \setminus N_i^\alpha \text{ the type } \text{tp}(b, N_i^\alpha, N_i^{\alpha+1}) \text{ is not realized in } N_\alpha$
 (a key point).

There is no problem to carry out the construction ($\mathfrak{K}$ has amalgamation in λ).

Let $w_i^\alpha =: \{\beta : N_i^{\alpha+1} \cap N_{\beta+1} \not\subseteq N_\beta\}$, so necessarily $|w_i^\alpha| \leq \lambda$, w_i^α is increasing continuous in $i < \lambda^+$ and $\alpha = \bigcup_{i < \lambda^+} w_i^\alpha$ and for $\beta < \alpha$ let

$$\mathbf{i}(\beta, \alpha) = \text{Min}\{i : \beta \in w_i^\alpha\}.$$

Now for every $\alpha \in S^* =: \{\delta < \lambda^{++} : \text{cf}(\delta) = \lambda^+\}$, the set

$$E_\alpha =: \left\{ i < \lambda^+ : i \text{ limit, } N \leq_{\mathfrak{K}} N_i^{\alpha+1}, a_\alpha \in N_i^{\alpha+1}, \right. \\
\text{and for every } \beta < \alpha \text{ if} \\
\beta \in w_i^\alpha \text{ then} \\
N_i^\beta = N_i^\alpha \cap N_\beta \text{ and for } j < i \\
\text{the closure of } w_j^\alpha \text{ (in } \alpha) \text{ is included in } w_i^\alpha \\
\left. \text{and } \beta_1 < \beta_2 \ \& \ \beta_1 \in w_j^\alpha \ \& \ \beta_2 \in w_j^\alpha \Rightarrow \mathbf{i}(\beta_1, \beta_2) < i \right\}$$

is a club of λ^+ .

As we can assume $\lambda > \aleph_0$ (ignoring $\lambda = \aleph_0$ as was treated earlier in [Sh 88] though for a $\text{PC}_{\aleph_0}$ class, or see [Sh 600], §2), we can choose $j_\alpha \in E_\alpha$ such that $\text{cf}(j_\alpha) = \aleph_1$ and let $\delta_\alpha = \sup(w_{j_\alpha}^\alpha)$, now $w_{j_\alpha}^\alpha$ is closed under ω -limits (as $\langle w_j^\alpha : j \leq \alpha \rangle$ is increasing continuous, $j < \alpha \Rightarrow \text{closure}(w_j^\alpha) \subseteq w_{j+1}^\alpha$) and $\aleph_1 = \text{cf}(\text{otp } w_{j_\alpha}^\alpha)$ so there is $\langle \beta_\varepsilon : \varepsilon < \omega_1 \rangle$ increasing continuous with limit $\delta_\alpha, \beta_\varepsilon \in w_{j_\alpha}^\alpha$ so $\varepsilon < \zeta < \omega_1 \Rightarrow N_j^{\beta_\varepsilon} = N_j^{\beta_\zeta} \cap N_{\beta_\varepsilon}$ and easily $N_j^\alpha \cap N_{\beta_\varepsilon} = N_j^{\beta_\varepsilon}$ hence

$$\bigoplus M_{j_\alpha}^{\delta_\alpha} = \bigcap \left\{ \bigcup_{j \in C} N_j^\beta : C \text{ a club of } \delta_\alpha \right\}$$

so $N_{j_\alpha}^\alpha = M_{j_\alpha}^{\delta_\alpha}$ (see [Sh 351, §4]).

By Fodor lemma for some j^*, α^* and stationary $S \subseteq S^*, \alpha \in S^* \Rightarrow j_\alpha = j^* \ \& \ \delta_\alpha = \delta^*$. So for all $\alpha \in S, N_{j_\alpha}^\alpha$ are the same say N^* . So $N^* \in K_\lambda$, for $\alpha \in S$, $q_\alpha = \text{tp}(a_\alpha, N^*, N_{j_\alpha}^{\alpha+1})$ extend $p_\alpha \in \mathcal{S}$. Also if $r \in \mathcal{S}(N^*)$ is realized in $N_{j_\alpha}^{\alpha+1}$ say by b (for some $\alpha \in S$) then no member of $\bigcup \{N_{j^*}^{\beta+1} \setminus N_{j^*}^\beta : \beta \in S \cap \alpha\}$ realizes it (holds by clause (G), see clause (F)). So the sets $\Gamma_\alpha = \{\text{tp}(b, N_{j^*}^{\delta^*}, N_{j^*}^{\alpha+1}) : b \in N_{j^*}^{\alpha+1} \setminus N^*\}$ are pairwise disjoint and each has a member extending $p_\alpha \in \mathcal{S}$ (as exemplified by a_α and p_α is not extended by any $p \in \bigcup_{\beta < \alpha} \Gamma_\beta$ (as p_α is not realized in N_α)). $\square_{4.6}$

4.7 Claim. Assume p^* is a counterexample to 4.4.

1) If $N \in K_\lambda, \Gamma \subseteq \mathcal{S}_{\geq p^*}(N), |\Gamma| \leq \lambda^+$ then

$\{p \in \mathcal{S}_{\geq p^*}(N) : \text{for some } N', N \leq_{\mathfrak{K}} N' \in K_\lambda \text{ and some } b \in N' \\ \text{realizes } p \text{ but no } b \in N' \text{ realizes any } q \in \Gamma\}$

has cardinality $\geq \lambda^{++}$.

2) We can find $N \in K_{\lambda^+}$, and $N_i, N <_{\mathfrak{K}} N_i \in K_{\lambda^+}$ for $i < 2^{\lambda^+}$ such that the set $\Gamma_i = \{tp(a, N, N_i) : a \in N_i\}$ are pairwise distinct, in fact, no one embeddable into another (so we get $I(\lambda^+, \mathfrak{K}) = 2^{\lambda^+}$ and if $(2^\lambda)^+ < 2^{\lambda^+}$ then $IE(\lambda^+, \mathfrak{K}) = 2^{\lambda^+}$ thus contradicting the categoricity in λ^+ from the assumptions).

Proof. 1) Apply 4.6 with N and $\mathcal{S}_{\geq p^*}(N)$ here standing for N and $\mathcal{S}$ there, so we get $N^*, N_i (i < \lambda^{++})$ and M^* such that $N^* \leq_{\mathfrak{K}} N_i \in \mathfrak{K}_\lambda$, and $\Gamma_i = \{tp(a, N^*, N_i) : a \in N_i \setminus N^*\}$ are pairwise disjoint and there are $p_i \in \Gamma_i, p_i \upharpoonright N \in \mathcal{S}_{\geq p^*}(N)$ pairwise distinct; now p_i is not algebraic hence $p_i \in \mathcal{S}_{\geq p^*}(N^*)$. As K is categorical in λ , without loss of generality $N^* = N$, so all but $\leq \lambda^+$ of the models N_i can serve as the required N' .

2) Now by part (1) of 4.7 we can choose by induction on $i < \lambda^+$, $\langle (N_\eta, \Gamma_\eta) : \eta \in {}^i 2 \rangle$ such that:

- (a) $N_\eta \in K_\lambda$ and $\Gamma_\eta \subseteq \bigcup_{j < i} \mathcal{S}_{\geq p^*}(N_{\eta \upharpoonright j})$ and $|\Gamma_\eta| \leq \lambda$
- (b) if $\nu \triangleleft \eta$ then $N_\nu \leq_{\mathfrak{K}} N_\eta$ and $\Gamma_\eta \subseteq \Gamma_\nu$
- (c) some $p \in \Gamma_{\eta \wedge \langle 0 \rangle}$ is from $\mathcal{S}(N_\eta)$ and is realized in $N_{\eta \wedge \langle 1 \rangle}$
(similarly for $\Gamma_{\eta \wedge \langle 1 \rangle}, N_{\eta \wedge \langle 0 \rangle}$)
- (d) if i is a limit ordinal, then $N_\eta = \bigcup_{j < i} N_{\eta \upharpoonright j}$ and $\Gamma_\eta = \bigcup_{j < i} \Gamma_{\eta \upharpoonright j}$.

The successor case is done by 4.7(1) (you may object that the type in Γ_η are not from $\mathcal{S}_{\geq p^*}(N_\eta)$ but from $\bigcup_{j < i} \mathcal{S}_{\geq p^*}(N_{\eta \upharpoonright j})$? However they are minimal - see 4.5(1)).

For $\eta \in \lambda^+ 2$ let $N_\eta = \bigcup_{i < \lambda^+} N_{\eta \upharpoonright i}$. Now by 1.4(1), $\{N_\eta / \cong : \eta \in \lambda^+ 2\}$ has cardinality 2^{λ^+} , contradiction. $\square_{4.7}, \square_{4.4}, \square_{4.3}$

4.8 Claim. 1) If $M_0 \in K_\lambda, M_1 \in K_{\lambda^+}$, and $M_0 \leq_{\mathfrak{K}} M_1$ then every minimal $p \in \mathcal{S}(M_0)$ is realized in M .

2) Every $M_1 \in K_{\lambda^+}$ is saturated at least for minimal types (i.e. if $M_0 \leq_{\mathfrak{K}} M_1$, $M_0 \in K_\lambda$ and $M_1 \in K_{\lambda^+}$ then every minimal $p \in \mathcal{S}(M_0)$ is realized in M_1).

3) If $M \in K_\lambda$ then $\{p \in \mathcal{S}(M) : p \text{ minimal}\}$ has cardinality $\leq \lambda^+$.

Proof. 1) Let $\bar{N} = \langle N_\alpha : \alpha < \lambda^+ \rangle$ be a representation of $N_{\lambda^+} \in K_{\lambda^+}$. Let $N \in K_\lambda, p \in \mathcal{S}(N)$ be minimal we ask

$(*)_p$ is there a club of $\alpha < \lambda^+$ such that every $q \in \mathcal{S}_{\geq p}(N_\alpha)$ is realized in λ^+ ?

By 4.4 there is $N'_{\lambda^+} \in K_{\lambda^+}$ for which the answer is yes, hence, as $\mathfrak{K}$ is categorical in λ^+ , this holds for N_{λ^+} . So this holds for every minimal p . Now if $N' \leq_{\mathfrak{K}} N_{\lambda^+}$, $N' \in K_{\lambda}$ and $p \in \mathcal{S}(N')$ is minimal then for some α , $N' \leq_{\mathfrak{K}} N_{\alpha}$ and for every $\beta \in [\alpha, \lambda^+)$, p has a unique non-algebraic extension $p_{\beta} \in \mathcal{S}(N_{\beta})$ (which necessarily is minimal). Now $p_{\beta} \in \mathcal{S}_{\geq p}(N_{\beta})$ hence for a club of $\beta < \lambda^+$, p_{β} is realized in N_{λ^+} , so we have finished the proof of part (1).

2) By part (1).

3) Follows by part (1). $\square_{4.8}$

From 4.3, 2.9(1) we can conclude

4.9 Conclusion. Every $(M_0, M_1, a) \in K_{\lambda}^3$ has the extension property.

§5 INEVITABLE TYPES AND STABILITY IN λ

5.1 *Hypothesis.* Assume the model theoretic assumptions from 4.1 and

- (d) there is a minimal member of K_λ^3 (follows from the conclusion of 3.28).

5.2 Definition. We call $p \in \mathcal{S}(N)$ inevitable if:

$N \leq_{\mathfrak{K}} M$ & $N \neq M \Rightarrow$ some $c \in M$ realizes p .

We call $(M, N, a) \in K_\lambda^3$ inevitable if $\text{tp}(a, M, N)$ is inevitable.

So by 4.3 - 4.8 we shall deduce

5.3 Claim. 1) If there is a minimal triple in K_λ^3 , then there is an inevitable $p = \text{tp}(a, N, N_1)$ with $(N, N_1, a) \in K_\lambda^3$ minimal.

2) Moreover, if $p_0 \in \mathcal{S}(N_0)$ is minimal, $N_0 \in K_\lambda$ then we can find $N_1, N_0 \leq_{\mathfrak{K}} N_1 \in K_\lambda$ such that the unique non-algebraic extension p_1 of p_0 in $\mathcal{S}(N_1)$ is inevitable.

Proof of 5.3. 1) Follows by part (2).

2) Let $(M_0, M_1, a) \in K_\lambda^3$ be minimal and $p_0 = \text{tp}(a, M_0, M_1)$. We try to choose by induction on i a model N_i such that: $N_0 = M_0, N_i \in K_\lambda$ is $\leq_{\mathfrak{K}}$ -increasing continuously and N_i omits $p_0, N_i \neq N_{i+1}$. If we succeed, $\bigcup_{i < \lambda^+} N_i$ is a member of

K_{λ^+} which is non-saturated for minimal types, contradicting 4.8(2). As for $i = 0$, i limit we can define, necessarily for some i we have N_i but not N_{i+1} . Now p_0 has a unique extension in $\mathcal{S}(N_i)$ which we call p_i and p_0 has no algebraic extension in $\mathcal{S}(N_i)$.

[why? as N_i omits p_0]. So p_i is the unique extension of p_0 in $\mathcal{S}(N_i)$ [by 4.5(1)], and so

- (*) if $N_i \leq_{\mathfrak{K}} N' \in K_\lambda$ and $N' \neq N_i$, then p_i is realized in N' .

By L.S. we can omit " $N' \in K_\lambda$ ", so (N_i, p_i) are as required. □_{5.3}

5.4 *Fact.* Inevitable types have few ($\leq \lambda$) conjugates (i.e. for $p \in \mathcal{S}(M_0)$ inevitable $M_0 \in K_\lambda, M_1 \in K_\lambda$ we have $|\mathcal{S}_p(M_1)| \leq \lambda$), moreover $|\{p \in \mathcal{S}(N) : p \text{ inevitable}\}| \leq \lambda$ for $N \in K_\lambda$.

Proof. Easy.

The following construction plays a central role in what remains of this paper.

5.5 Claim. For any limit $\alpha < \lambda^+$, we can find $\langle N_i : i \leq \alpha \rangle$ and $\langle p_i : i \leq \alpha \rangle$ such that:

- (i) $N_i \in K_\lambda$,
- (ii) N_i is $\leq_{\mathfrak{K}}$ -increasing continuous
- (iii) $p_i \in \mathcal{S}(N_i)$ is minimal,

- (iv) p_i increases continuously (see 4.5(3))
- (v) p_0 is inevitable
- (vi) p_α is inevitable
- (vii) $N_i \neq N_{i+1}$ moreover some $c \in N_{i+1} \setminus N_i$ realizes p_0 (hence p_i).

Remark. Why not just build a non-saturated model in order to prove 5.5? Works, too.

Proof. Choose $N^0 <_{\mathfrak{K}} N^1$ in K_{λ^+} (so $N^0 \neq N^1$), such a pair exists as $K_{\lambda^+} \neq \emptyset$. Let $N^\ell = \bigcup_{i < \lambda^+} N_i^\ell$ with $N_i^\ell \in K_\lambda$ being $\leq_{\mathfrak{K}}$ -increasing continuously in i . Now $E_0 = \{\delta < \lambda^+ : N_\delta^1 \neq N_\delta^0 \text{ and } N_\delta^1 \cap N^0 = N_\delta^0\}$ is a club of λ^+ . Without loss of generality $E_0 = \lambda^+$.

For each $c \in N^1 \setminus N^0$, the set

$$X_c =: \{i < \lambda^+ : c \in N_i^1 \text{ and } (N_i^0, N_i^1, c) \text{ is minimal}\}$$

is empty or an end segment of λ^+ hence

$$E_1 = \left\{ \delta < \lambda^+ : \begin{array}{l} \text{(i) } \delta \text{ limit} \\ \text{(ii) if } i < \delta \text{ and } p \in \mathcal{S}(N_i^0) \text{ is minimal inevitable} \\ \quad \text{and realized in } N^0 \setminus N_\delta^0 \text{ then it is} \\ \quad \text{realized in } N_\delta^0 \setminus N_i^0 \text{ (actually automatic)} \\ \text{(iii) if } c \in N_\delta^1 \setminus N^0 \text{ (hence } \exists i < \delta, c \in N_i^1) \text{ and } X_c \\ \quad \text{is non-empty then } \delta \in X_c \text{ and } \min(X_c) < \delta \end{array} \right\}$$

is a club of λ^+ (see 5.4).

Now for $\delta \in E_1$, we have $N_\delta^0 <_{\mathfrak{K}} N_\delta^1$, so by 5.3(1) there is $c_\delta \in N_\delta^1 \setminus N_\delta^0$ such that:

$$\begin{array}{l} (N_\delta^0, N_\delta^1, c_\delta) \text{ is minimal} \\ \text{tp}(c_\delta, N_\delta^0, N_\delta^1) \text{ is inevitable} \end{array}$$

As δ is limit, for some $i < \delta, c \in N_i^1$, also $\delta \in X_c$ hence there is j such that: $i < j < \delta$ & $j \in X_c$ hence (N_j^0, N_j^1, c) is minimal; choose such j_δ, c_δ . Let $\kappa = \text{cf}(\kappa) = \text{cf}(\alpha) \leq \lambda$, so for some j^*, c^* we have

$$S = \{\delta \in E_1 : \text{cf}(\delta) = \kappa, j_\delta = j^*, c_\delta = c^*\}$$

is stationary in λ^+ .

Choose e closed $\subseteq E_1$ of order type $\alpha + 1$ with first element and last element in S ; for $\zeta \in [j^*, \lambda^+)$ let $p_\zeta = \text{tp}(c^*, N_\zeta^0, N_\zeta^1)$. (In fact, we could have: all non-accumulation member of e are in S ; no real help.)

Now $\langle N_\zeta^0, p_\zeta : \zeta \in e \rangle$ is as required (up to re-indexing)(clause (viii) holds by clause (ii) in the definition of E_1). $\square_{5.5}$

5.6 Claim. *Assume $\langle N_i, p_i : i \leq \alpha \rangle$ is as in 5.5, $\alpha < \lambda$ divisible by λ . Then any $p \in \mathcal{S}(N_0)$ is realized in N_α , moreover N_α is universal in K_λ over N_0 .*

Proof. (Similar to the proof of 0.25; which is [Sh 300, II, §3]).

Let $N_0 \leq_{\mathfrak{K}} M_0 \in K_\lambda$, $a \in M_0 \setminus N_0$ we shall show that $\text{tp}(a, N_0, M_0)$ is realized in N_α .

Let $\alpha = \bigcup_{i < \lambda} S_i$, $\langle S_i : i < \lambda \rangle$ pairwise disjoint, each S_i unbounded in α , λ divides $\text{otp}(S_i)$ and $\text{Min}(S_i) \geq i$. We choose by induction on $i \leq \alpha$ the following:

$$N_i^1, M_i^1, h_i, \langle a_\zeta : \zeta \in S_i \rangle \quad (\text{the last one only if } i < \alpha)$$

such that:

- (a) $N_i^1 \leq_{\mathfrak{K}} M_i^1$ are in K_λ
- (b) N_i^1 is $\leq_{\mathfrak{K}}$ -increasing continuous in i
- (c) M_i^1 is $\leq_{\mathfrak{K}}$ -increasing continuous in i
- (d) $(N_0^1, M_0^1) = (N_0, M_0)$
- (e) $\langle a_\zeta : \zeta \in S_i \rangle$ is a list of $\{c \in M_i^1 : c \text{ realizes } p_0\}$
- (f) h_i is an isomorphism from N_i onto N_i^1
- (g) $j < i \Rightarrow h_j \subseteq h_i$ and $h_0 = \text{id}_{N_0}$
- (h) $a_i \in N_{i+1}^1$ (note: $M_i^1 \cap N_{i+1}^1 \neq N_i^1$ in general).

For $i = 0$: See clauses (d), (g)

$$N_0^1 = N_0, M_0^1 = M_0, h_0 = \text{id}_{N_0}.$$

For $i = \text{limit}$: Let $N_i^1 = \bigcup_{j < i} N_j^1$ and $M_i^1 = \bigcup_{j < i} M_j^1$ and $h_i = \bigcup_{j < i} h_j$ and lastly choose $\langle a_\zeta : \zeta \in S_i \rangle$ by clause (e).

For $i = j + 1$: Note a_j is already defined, it belongs to M_j^1 and it realizes p_0 .

Case 1: $a_j \in N_j^1$ (so clause (h) is no problem).

Use amalgamation on N_j, N_i, M_j^1 and the mapping id_{N_j}, h_i , i.e.

$$\begin{array}{ccc} N_i & \longrightarrow & M_i^1 \\ \text{id}_{N_j} \uparrow & & \uparrow \text{id}_{N_0^1} \\ N_j & \xrightarrow{h_i} & N_j^1 \end{array}$$

Case 2: $a_j \notin N_j^1$.

Then $\text{tp}(a_j, N_j^1, M_j^1)$ is not algebraic, extending the minimal type $p_0 \in \mathcal{S}(N_0)$. Also by clause (viii) of 5.5 there is $c \in N_i \setminus N_j$ which realizes p_0 . As $p_0 \in \mathcal{S}(N)$ is minimal

$$h_j(\text{tp}(c, N_j, N_i)) = \text{tp}(a_j, N_j^1, M_j^1)$$

so acting as in Case 1 we can also guarantee $h_i(c) = a_j$, so $a_j \in \text{Rang}(h_i) = N_i^1$ as required.

* * *

In the end we have $N_\alpha^1 \leq_{\mathfrak{K}} M_\alpha^1$. If $N_\alpha^1 = M_\alpha^1$, then $h_\alpha^{-1} \upharpoonright M_0 = h_\alpha^{-1} \upharpoonright M_0^1 = h_\alpha^{-1} \upharpoonright N_0^1$ show that M_0 can be embedded into N_α over N_0 as required. So assume $N_\alpha^1 <_{\mathfrak{K}} M_\alpha^1$. Now $p_\alpha \in \mathcal{S}(N_\alpha)$ is inevitable hence $h_\alpha(p_\alpha) \in \mathcal{S}(N_\alpha^1)$ is inevitable. Hence some $d \in M_\alpha^1 \setminus N_\alpha^1$ realizes $h_\alpha(p_\alpha)$ hence d realizes $h_\alpha(p_\alpha) \upharpoonright N_0^1 = p_0$; also α is a limit ordinal so for some $i < \alpha$, $d \in M_i^1$ hence for some $\zeta \in S_i$ we have $a_\zeta = d$, hence

$$d = a_\zeta \in N_{\zeta+1}^1 \subseteq N_\alpha^1,$$

contradicting the choice of d .

So we are done. □_{5.6}

5.7 Conclusion. If $N \in K_\lambda$ then:

- (a) $|\mathcal{S}(N)| = \lambda$
- (b) there is $N_1, N <_{\mathfrak{K}} N_1 \in K_\lambda$ such that N_1 is universal over N in K_λ
- (c) for any regular $\kappa \leq \lambda$ we can demand that $(N_1, c)_{c \in N}$ is (λ, κ) -saturated (see 0.6(1))

5.8 Remark. In fact amalgamation in λ and stability in λ (i.e. (a) of 5.7) implies (b) and (c) of 5.7.

5.9 Conclusion. The $N \in K_{\lambda+}$ is saturated above λ (i.e. over models in $K_\lambda!$).

5.10 Claim. Assume $\kappa = \text{cf}(\kappa) \leq \lambda$.

There are $N_0, N_1, a, N_0^+, N_1^+$ such that

- (i) $(N_0, N_1, a) \in K_\lambda^3$ and
- (ii) $(N_0, N_1, a) \leq (N_0^+, N_1^+, a) \in K_\lambda^3$ and
- (iii) $(N_0^+, c)_{c \in N_0}$ is (λ, κ) -saturated,
- (iv) $\text{tp}(a, N_0, N_1)$ is minimal inevitable and
- (v) $\text{tp}(a, N_0^+, N_1^+)$ is minimal inevitable.

Proof. As in the proof of 5.5 as

$$E_2 = \left\{ \delta : \text{for every } i < \delta, (N_\delta^0, c)_{c \in N_i} \text{ is saturated} \right. \\ \left. \text{of cofinality } \text{cf}(\delta) \right\}$$

is a club of λ^+ .

□_{5.10}

5.11 Claim. 1) In K_λ we have disjoint amalgamations.

2) If $M \leq_{\mathfrak{K}} N$ are in K_λ and $p \in \mathcal{S}(M)$ non-algebraic then for some N', c we have: $N \leq_{\mathfrak{K}} N' \in K_\lambda$ and $c \in N' \setminus N$ realizes p .

Proof. 1) First note

⊗ if $M \leq_{\mathfrak{K}} N$ in K_λ we can find $\alpha < \lambda^+$, and sequence $\langle M_i : i \leq \alpha \rangle$ which is $\leq_{\mathfrak{K}}$ -increasing continuous, and $\langle a_i : i < \alpha \rangle$ such that (M_i, M_{i+1}, a_i) is minimal and reduced and $N \leq_{\mathfrak{K}} M_\alpha, M = M_0$.

[Why? There is a minimal reduced pair, hence we can find $\langle M_i : i < \lambda^+ \rangle \leq_{\mathfrak{K}}$ -increasing continuous, (M_i, M_{i+1}, a_i) minimal reduced and $M = M_0$. So by 5.9 we know $\bigcup_{i < \lambda^+} M_i \in K_{\lambda^+}$ is saturated, hence we can embed

N into $\bigcup_{i < \lambda^+} M_i$ over N so this embedding is into some $M_\alpha, \alpha < \lambda^+$.]

So given $M \leq_{\mathfrak{K}} M^1, M^2$, without loss of generality $M^\ell = M_{\alpha_\ell}^\ell, \langle (M_i^\ell, a_i^\ell) : i \leq \alpha_\ell \rangle$ as above, and start to amalgamate using the extension property and “reduced”.

□_{5.11}

5.12 Remark. We could prove 5.11 earlier using “reduced triples”. I.e. note that for some $\langle M_i^1 : i < \lambda^+ \rangle \in \text{Seq}_{\lambda^+}[\mathbf{C}_{\mathfrak{K}, \lambda^+}^1]$, for each i for some a the triple $(M_i^1, M_{i+1}^1, a) \in K_\lambda^3$ is reduced. Hence if $M \leq_{\mathfrak{K}} N$ from K_λ , for some $\bar{M} = \langle M_i : i \leq \alpha \rangle, \leq_{\mathfrak{K}}$ -increasing continuous, $\langle M_i, M_{i+1}, b_i \rangle \in K_\lambda^3$ is reduced, $M_0 = M, N \leq M_\alpha \in K_\lambda$ (otherwise find $\langle M_i^2 : i < \lambda^+ \rangle \in \text{Seq}_{\lambda^+}[\mathbf{C}_{\mathfrak{K}, \lambda^+}^1]$ with $(M_i, M_{i+1}) \cong (M, N)$, hence $M^1 = \bigcup_{i < \lambda^+} M_i^1, M^2 = \bigcup_{i < \lambda^+} M_i^2$ are non-isomorphic members of K_{λ^+} , contradiction). Now prove by induction on $\beta \leq \alpha$ that if $M \leq_{\mathfrak{K}} N_0 \in K_\lambda$ then N_0, M_β has disjoint amalgamation over $M_0 = M$ (i.e. we need to decompose only one side).

5.13 Question: If $M \in K_\lambda, p \in \mathcal{S}(M)$ is minimal, is it reduced? Or at least, if $M_0 \leq_{\mathfrak{K}} M_1$ are in $K_\lambda, p_1 \in \mathcal{S}(M_\ell)$, no algebraic $p_0 = p_1 \upharpoonright M_0, p_0$ is minimal and reduced is also p_1 reduced?

Probably true, but not needed for our aims (here and in [Sh 600], 7.6(2)) is an approximation. Can prove it if $\lambda < \lambda^{\aleph_0}$ or there are E.M. models.

§6 PROVING FOR $\mathfrak{K}$ CATEGORICAL IN λ^{+2}

6.1 Hypothesis. Assume the model theoretic assumptions from 4.1 + 5.1 and so the further model theoretic properties deduced in §4 + §5.

We use heavily 4.8.

6.2 Definition. 1) We say $(M_0, M_1, M_2) \in K_\lambda^{\text{uniq}}$ (has unique amalgamation in K_λ) when

- (a) $M_0, M_1, M_2 \in K_\lambda$
- (b) $M_0 \leq_{\mathfrak{K}} M_1$ and $M_0 \leq_{\mathfrak{K}} M_2$
- (c) if for $i = 1, 2$ we have $g_\ell^i : M_\ell \rightarrow N_i \in K_\lambda$ such that:
 - (i) g_ℓ^i a $\leq_{\mathfrak{K}}$ -embedding,
 - (ii) $g_0^i \subseteq g_1^i$ and $g_0^i \subseteq g_2^i$ for $i = 1, 2$
 - (iii) $\text{Rang}(g_1^i) \cap \text{Rang}(g_2^i) = \text{Rang}(g_0^i)$
(disjoint amalgamation) for $i = 1, 2$then we can find $N \in K_\lambda$ and $\leq_{\mathfrak{K}}$ -embeddings

$$f^i : N_i \rightarrow N \text{ for } i = 1, 2$$

such that

$$\bigwedge_{\ell < 3} f^1 \circ g_\ell^1 = f^2 \circ g_\ell^2.$$

2) Let $K_\lambda^{2,\text{uq}}$ is the class of pairs (M_0, M_2) such that $M_0 \leq_{\mathfrak{K}} M_2$ are both in K_λ and $[M_0 \leq_{\mathfrak{K}} M_1 \in K_\lambda \Rightarrow (M_0, M_1, M_2) \in K_\lambda^{\text{uniq}}]$ and let $K_\lambda^{3,\text{uq}}$ be the class of pairs $(M_0, M_2) \in K_\lambda^{2,\text{uq}}$ satisfying $M_0 \neq M_2$.

6.3 Claim. 1) If $(M_0, M_1, M_2) \in K_\lambda^{\text{uniq}}$ then

- (a) $(M_0, M_2, M_1) \in K_\lambda^{\text{uniq}}$
- (b) if $M_0 \leq_{\mathfrak{K}} M'_2 \leq_{\mathfrak{K}} M_2$ then $(M_0, M_1, M'_2) \in K_\lambda^{\text{uniq}}$.

2) Assume $M_0 \leq_{\mathfrak{K}} M_2$ are from K_λ and $M_1 \in K_\lambda$ is universal over M_0 . Then $(M_0, M_2) \in K_\lambda^{2,\text{uq}} \Leftrightarrow (M_0, M_1, M_2) \in K_\lambda^{\text{uniq}}$.

Proof. 1)a) Trivial.

b) Chase arrows (using disjoint amalgamation; i.e. 5.11).

2) Follows by 6.3(1)(a)+(b) and the definition. □_{6.3}

6.4 Lemma. *Suppose*

$\otimes$ *there is $(M_0, M_1, M_2) \in K_\lambda^{\text{uniq}}$ such that $M_0 \neq M_2$ and M_1 is universal over M_0 .*

Then: *there are $N^0 <_{\aleph} N^1$ in K_{λ^+} such that:*

- (a) $N^0 \neq N^1$
- (b) *for every $c \in N^1 \setminus N^0$ there is $M = M_c$ satisfying $N^0 \leq_{\aleph} M \leq_{\aleph} N^1$ and $N^0 \neq M$ and $c \in N^1 \setminus M$.*

Proof. Choose $\langle N_i^0 : i < \lambda^+ \rangle$, a sequence of members of K_λ which is $\leq_{\aleph}$ -increasing continuous, such that:

$$(N_i^0, N_{i+1}^0) \cong (M_0, M_2).$$

So $N_i^0 \neq N_{i+1}^0$ hence $N_0 = \bigcup_{i < \lambda^+} N_i^0 \in K_{\lambda^+}$ and without loss of generality

$$|N_0| = \lambda^+.$$

We now choose by induction $i < \lambda^+$, N_i^1 and $M_{i,c}$ for $c \in N_i^1 \setminus N_i^0$ such that:

- (a) $N_i^0 \leq_{\aleph} N_i^1 \in K_\lambda$ and $N_i^0 \neq N_i^1$
- (b) N_i^1 is $\leq_{\aleph}$ -increasing continuous in i
- (c) $j < i \Rightarrow N_j^1 \cap N_i^0 = N_j^0$; moreover $N_i^1 \cap |N_0| = N_i^0$
- (d) $N_i^0 \leq_{\aleph} M_{i,c} \leq_{\aleph} N_i^1$
- (e) $c \notin M_{i,c}$
- (f) $N_i^0 \neq M_{i,c}$
- (g) if $j < i$ and $c \in N_j^1 \setminus N_j^0$ then $M_{i,c} \cap N_j^1 = M_{j,c}$.

For $i = 0$: Choose N_i^1 such that

$N_i^0 \leq_{\aleph} N_i^1$, $(N_i^1, c)_{c \in N_i^0}$ saturated (any cofinality will do) then by disjoint amalgamation easy to define the $M_{0,c}$ (remembering clause (c)).

For i limit: Straightforward.

For $i = j + 1$: First we disjointly amalgamate getting $N'_i \in K_\lambda$ such that

$N_i^0 \leq N'_i$, $N_j^1 \leq_{\aleph} N'_i$ and $|N'_i| \cap |N_0| = |N_i^0|$ (as set of elements).

Let N_i^1 be such that:

$$\begin{aligned} N'_i &\leq_{\aleph} N_i^1 \in K_\lambda \\ (N_i^1, c)_{c \in N'_i} &\text{ is saturated (any cofinality will do)} \\ |N_i^1| \cap |N_0| &= |N_i^0|. \end{aligned}$$

Lastly we shall find the $M_{i,c}$'s, the point is that $(N_j^0, N_i^0, N_j^1) \in K_\lambda^{\text{uniq}}$ (by 6.3(2)).

By $\otimes$ and Claim 6.3 we could have done the amalgamation in two steps and use uniqueness. Then by uniqueness of saturated extensions embed the result inside N_i^1 and similarly deal with new c 's.

Now let $N_1 =: \bigcup_{1 < \lambda^+} N_i^1$ and for $c \in N_1 \setminus N_0$ let $M_c = \bigcup \{M_{i,c} : c \in N_i'\}$ they are as required. $\square_{6.4}$

Remark. The proof of 6.5 below is like [Sh 88, proof of 3.8 stage(c)]. The aim is to contradict that under $I(\lambda^+, \mathfrak{K}) = 0$ there are maximal triples.

6.5 Conclusion. Assume $\mathfrak{K}$ has amalgamation in λ^+ . If $\otimes$ of 6.4, then there is no maximal triple (M, N, a) in $K_{\lambda^+}^3$.

Proof. We can get by 6.4 a contradiction.

[Why? Assume $(N_0, N_2, a) \in K_{\lambda^+}^3$ maximal, (N^0, N^1) as in the conclusion (i.e. (a) + (b)) of 6.4; by categoricity in λ^+ without loss of generality $N_0 = N^0$ and let $N_1 = N^1$. Now $\mathfrak{K}$ has amalgamation for λ^+ so there are $N \in K_{\lambda^+}$ and f such that $f : N_2 \rightarrow N$ is a $\leq_{\mathfrak{K}}$ -embedding of N_2 into N over N_0 and $N_1 \leq_{\mathfrak{K}} N$. If $f(a) \notin N_1$, then $(N_0, N_2, a) <_f (N_1, N, f(a))$ contradict maximality. If $f(a) \in N_1$, then $M_{f(a)}$ is well defined (see 6.4) and $(N_0, N_2, a) <_f (M_{f(a)}, N, f(a))$ contradicts maximality.] $\square_{6.5}$

6.6 Remark. 1) Another proof is to replace the assumption “ $\mathfrak{K}$ has amalgamation in λ^+ ” by $I(\lambda^+, K) < 2^{\lambda^{++}}$. We start with N_0, N_1, N_2, a as above and building, for every $S \subseteq \lambda^{++}$, a sequence $\langle M_{\alpha}^S : \alpha < \lambda^{++} \rangle$ of members of K_{λ^+} , which is $\leq_{\mathfrak{K}}$ -increasing continuous, and $\alpha \in S \Rightarrow (M_{\alpha}^S, M_{\alpha+1}^S, a_{\alpha}^S) \cong (N_0, N_2, a)$, and $\alpha \in \lambda^{++} \setminus S \Rightarrow (M_{\alpha}^S, M_{\alpha+1}^S) \cong (N_0, N_1)$ which are as in (a) + (b) of 6.4. Let $M^S = \bigcup_{\alpha < \lambda^{++}} M_{\alpha}^S \in K_{\lambda^{++}}$ and from $M^S / \cong$ we can reconstruct $S / \mathcal{D}_{\lambda^{++}}$. So here we use $I(\lambda^{++}, K) < 2^{\lambda^{++}}$ but not the definitional weak diamond for λ^{++} .
2) Note that if $2^{\lambda^+} < 2^{\lambda^{++}}$ then the assumption of 6.6(1) implies the assumption of 6.5.

6.7 Claim. *Assume*

$$(*) \quad 2^{\lambda} < 2^{\lambda^+} < 2^{\lambda^{++}} \\ \text{(or at least the definitional weak diamond for } \lambda^+, \lambda^{++} \text{).}$$

If $\otimes$ of 6.4 fails, we get $I(\lambda^{++}, K) \geq \mu_{\text{wd}}(\lambda)$.

Proof. By 3.23 and 6.8 below.

The following serves to prove 6.7

6.8 Claim. Assume $M \in K_\lambda \Rightarrow |\mathcal{S}(M)| \leq \lambda$.

If $K_\lambda^{3, \text{uq}} = \emptyset$ (see Definition 6.2(2)), then there is F , an amalgamation choice function F for $\mathbf{C} = \mathbf{C}_{\mathfrak{K}, \lambda^+}^0$ with the weak λ^+ -coding property.

Proof. The point is that if $\bar{M} = \langle M_\alpha : \alpha < \lambda^+ \rangle \in \mathbf{Seq}_{\lambda^+}[\mathbf{C}]$ and $\alpha < \lambda^+, M_\alpha <_{\mathfrak{K}} N \in K_\lambda$, then for some $\beta \in (\alpha, \lambda^+)$ we have:

M_β is universal over M_α , so as $K_\lambda^{3, \text{uq}} = \emptyset$ necessarily
 $(M_\alpha, M_\beta, N) \notin K_\lambda^{\text{uniq}}$ and rest should be clear.

Of course, we use the extension property. □_{6.8}

6.9 Remark. We can work in the context of §3, we need the existence of a saturated (equivalent by super limit) $M \in K_{\lambda^+}$. We now say how to replace $\mu_{\text{wd}}(\lambda^{+2})$ by $2^{\lambda^{+2}}$.

6.10 Claim. 1) Assume each $M \in K_{\lambda^+}$ is saturated above λ .

If $(M, N, a) \in K_\lambda^3$ it and every $(M', N', a) \in K_\lambda^3$ above it has the extension property but for every $(M'', N'', a) \geq (N, N, a)$ (all in $K_{\lambda^+}^3$ for some $M^* \geq_{\mathfrak{K}} M''$ from K_{λ^+} , in amalgamation $(M^*, N^*, a) \geq (M'', N'', a)$ the type of $M^* \cup N$ inside N^* is not determined then some F (actually $\mathbf{F}^*$) has the λ^+ -coding.

2) If above we just require that the type of $M^* \cup N''$ inside N^* is not determined, then some F (actually $\mathbf{F}^*$) has weak λ^+ -coding.

3) We can restrict ourselves to disjoint embedding; i.e. use $(K_{\lambda^+}^3, \leq_{\text{dj}})$.

6.11 Discussion. 1) We get $IE(\lambda^{+2}, \mathfrak{K}) = 2^{\lambda^{+2}}$ when $(2^{\lambda^+})^+ < 2^{\lambda^{+2}}$. See more in [Sh 600].

6.12 Theorem. Assume $I(\lambda^{+2}, K) < \mu_{\text{wd}}(\lambda^{+2})$ and $(*)$ of 6.7 (or at least the conclusion of 6.7). Then $I(\lambda^{+2}, K) = 1 \Rightarrow I(\lambda^{+3}, K) > 0$.

Remark. As in [Sh 88, §3].

Proof. By 0.20(1) it is enough to show that for some $M \in K_{\lambda^{++}}$ there is M' , $M \leq_{\mathfrak{K}} M' \in K_{\lambda^{++}}, M \neq M'$.

[Why? As then we can choose by induction on $i < \lambda^{+3}$ models $M_i \in K_{\lambda^{+2}}$, $\leq_{\mathfrak{K}}$ -increasing continuous, $M_i \neq M_{i+1}$, for $i = 0$ use $K_{\lambda^{+2}} \neq \emptyset$, for i limit take union, for $i = j + 1$ use the previous sentence; so $M_{\lambda^{+3}} = \cup \{M_i : i < \lambda^{+3}\} \in K_{\lambda^{+3}}$ as required.]

By 6.7, the statement $\otimes$ of 6.4 holds so we can find (N^0, N^1) as there so by 6.5 there is in $K_{\lambda^+}^3$ no maximal member. This implies (easy, see 2.6(6)) that there are $M^* \leq_{\mathfrak{K}} N^*$ from $K_{\lambda^{+2}}$ such that $M^* \neq N^*$ which as said above (by categoricity in λ^+), suffices. □_{6.12}

§7 EXTENSIONS AND CONJUGACY

7.1 Hypothesis. Assume the model theoretic assumptions from 4.1 + 5.2 and the further model theoretic properties deduced since then (but not in 6.7, 6.12) or just

- (a) $\mathfrak{K}$ abstract elementary class
- (b) $\mathfrak{K}$ has amalgamation in λ
- (c) $\mathfrak{K}$ is categorical in λ (can be weakened)
- (d) $\mathfrak{K}$ is stable in λ (see 5.7, clause (a))
- (e) there is an inevitable $p \in \mathcal{S}(N)$ for $N \in K_\lambda$ (holds by 5.3)
- (f) the basic properties in type theory.

We now continue toward eliminating the use of $I(\lambda^{++}, K) = 1$ (in 6.12), and give more information. We first deal with the nice types in $\mathcal{S}(N)$, $N \in K_\lambda$ in particular the realize/materialize problem which is here: if $N_1 \leq_{\mathfrak{K}} N_2$ are in K_λ , $p_\ell \in \mathcal{S}(N_\ell)$ is minimal, $p_1 \leq p_2$ are they conjugate? (i.e. $p_2 \in \mathcal{S}_{p_1}(N_2)$).

7.2 Claim. *If $N \in K_\lambda$ and $p \in \mathcal{S}(N)$ is minimal and reduced or just p is reduced (see Definition 2.3(7)), then p is inevitable.*

Proof. Suppose N, p form a counterexample. We can then find N_1 and a such that $N \leq_{\mathfrak{K}} N_1 \in K_\lambda$, $a \in N_1 \setminus N$ and $p = \text{tp}(a, N, N_1)$ and (N, N_1, a) is reduced. As p is not inevitable, there is N_2 such that: $N \leq_{\mathfrak{K}} N_2 \in K_\lambda$, $N \neq N_2$ but no element of N_2 realizes p . By amalgamation in K_λ , without loss of generality there is $N_3 \in K_\lambda$ such that $\ell \in \{1, 2\} \Rightarrow N_\ell \leq_{\mathfrak{K}} N_3$. By 5.3 (i.e. 7.1(e)) there is $q \in \mathcal{S}(N)$ which is inevitable so there are $c_\ell \in N_\ell$ with $q = \text{tp}(c_\ell, N, N_\ell)$ for $\ell \in \{1, 2\}$. By the equality of types (and amalgamation in K_λ) there is $N^+ \in K$, a $\leq_{\mathfrak{K}}$ -extension of N_1 and a $\leq_{\mathfrak{K}}$ -embedding of N_2 into N^+ over N such that $f(c_2) = c_1$; so without loss of generality $N^+ = N_3$ and f is the identity, hence $c_1 = c_2$. Now $a \notin N_2$ as $p = \text{tp}(a, N, N_1)$ is not realized in N_2 . So $(N, N_1, a) \leq (N_2, N_3, a)$ and $N_2 \cap N_1 \setminus N \neq \emptyset$ contradicting “ (N, N_1, a) is reduced”. $\square_{7.2}$

7.3 Claim. 1) *If $\kappa = \text{cf}(\kappa) \leq \lambda$ and $\bar{N} = \langle N_i : i \leq \omega\kappa \rangle$ is an $\leq_{\mathfrak{K}}$ -increasingly continuous sequence, $N_i \in K_\lambda$, N_{i+1} universal over N_i , and $p \in \mathcal{S}(N_{\omega\kappa})$ is minimal reduced (or minimal inevitable) then for some $i < \omega\kappa$ we have $p \restriction N_i \in \mathcal{S}(N_i)$ is minimal (so p is the unique, non-algebraic extension of $p \restriction N_i$ in $\mathcal{S}(N_{\omega\kappa})$ (and of course, there is one)).*

2) *If $\lambda \geq \kappa = \text{cf}(\kappa)$, $\bar{N} = \langle N_i : i \leq \kappa \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous in K_λ and $p \in \mathcal{S}(N_\kappa)$ is minimal and reduced and the set $Y = \{i < \kappa : N_{i+1} \text{ is } (\lambda, \kappa)\text{-saturated over } N_i\}$ is unbounded in κ then for every large enough $i \in Y$ there is an isomorphism f from N_{i+1} onto N_κ which is the identity on N_i and⁵*

(*) *f maps $p \restriction N_{i+1} \in \mathcal{S}(N_{i+1})$ to $p \in \mathcal{S}(N_\kappa)$.*

⁵in fact, this implies (*)

Hence as p is minimal reduced, so is $p \upharpoonright N_{i+1}$.

Proof. 1) We can choose $(N_i^0, N_i^1, a) \in K_\lambda^3$ for $i < \lambda^+$ reduced, $\leq$ -increasing continuous such that $N_i^0 \neq N_{i+1}^0$. Let $N_\ell = \bigcup_{i < \lambda^+} N_i^\ell$. As in the proof of 5.5 for $c \in N_1 \setminus N_0$

$$I_c^* = \{j < \lambda^+ : c \in N_j^1 \text{ and } \text{tp}(c, N_j^0, N_j^1) \text{ is minimal}\}$$

is empty or is an end segment of λ^+ and

$$E = \left\{ \delta < \lambda^+ : \begin{array}{l} \text{if } c \in N_\delta^1 \text{ and } I_c^* \neq \emptyset \text{ then } I_c^* \cap \delta \\ \text{is an unbounded subset of } \delta; \text{ and if } \alpha < \delta \\ \text{then for some } \beta \in (\alpha, \delta), N_\beta^0 \text{ is universal over } N_\alpha^0 \\ \text{and if } Pr \text{ is one of the properties reduced and/or} \\ \text{inevitable and/or minimal and there is } i \geq \delta \text{ such that} \\ (N_i^0, N_i^1, c) \text{ has } Pr, \text{ then there are arbitrarily} \\ \text{large such } i < \delta \end{array} \right\}$$

is a club of λ^+ ; for the universality demand in the definition of E use categoricity in λ^+ . Let $\delta \in \text{acc}(\text{acc}(E))$, $\text{cf}(\delta) = \kappa$, let $\langle \alpha_\zeta : \zeta < \omega\kappa \rangle$ be an increasing continuous sequence of ordinals from E with limit δ , now set $\alpha_{\omega\kappa} = \delta$ and $N'_\zeta =: N_{\alpha_\zeta}^0$.

So there is an isomorphism f from $N_{\omega\kappa}$ onto $N_{\alpha_{\omega\kappa}}^0$ such that for every $\zeta < \omega\kappa$ we have $N_{\alpha_{2\zeta}}^0 \leq_{\mathfrak{R}} f(N_{\alpha_{2\zeta}}) \leq_{\mathfrak{R}} N_{\alpha_{2\zeta+1}}^0$ (so if ζ is a limit ordinal, then $N_{\alpha_\zeta}^0 = N_{\alpha_{2\zeta}}^0 = f(N_\zeta)$), so without loss of generality f is the identity. As $p \in \mathcal{S}(N_{\omega\kappa})$ is inevitable (by assumption or by 7.2) and $N_{\omega\kappa} = N_{\alpha_{\omega\kappa}}^0 <_{\mathfrak{R}} N_{\alpha_{\omega\kappa}}^1$, for some $c \in N_{\alpha_{\omega\kappa}}^1 \setminus N_{\alpha_{\omega\kappa}}^0$ we have $p = \text{tp}(c, N_{\alpha_{\omega\kappa}}^0, N_{\alpha_{\omega\kappa}}^1)$, so for some $\beta < \alpha_{\omega\kappa}$ we have $c \in N_\beta^1$. As p is minimal (by assumption) clearly $\delta \in I_c$, but $\delta \in E$ so $\text{Min}(I_c) < \delta$, but I_c is an end segment of λ^+ hence without loss of generality for some $\zeta < \omega\kappa$ we have $\beta = \alpha_\zeta \in I_c$. So for $\xi \in (\zeta, \omega\kappa)$, both $p \in \mathcal{S}(N_{\omega\kappa})$ and $p \upharpoonright N_\xi \in \mathcal{S}(N_\xi)$ are non-algebraic extensions of the minimal $p \upharpoonright N_{\alpha_\zeta}^0 \in \mathcal{S}(N_{\alpha_\zeta}^0)$ and $N_{\alpha_\zeta}^0 \leq_{\mathfrak{R}} N_\xi \leq_{\mathfrak{R}} N_{\omega\kappa}$, all in K_λ , so we have proved part (1).

2) Without loss of generality every $\zeta < \kappa$ is in Y . We can find $\langle N'_\zeta : \zeta \leq \omega\kappa \rangle$ as in part (1), moreover satisfying “ $N'_{\zeta+1}$ is (λ, κ) -saturated over N'_ζ ” and such that: for every $\zeta \leq \kappa$ we have $N_\zeta = N'_{\omega\zeta}$. So again choose $\zeta < \kappa$ as there, we set $\beta = \alpha_{\omega\zeta} \in I_c^*$. If $\xi \in Y$ & $\omega\xi > \zeta$ clearly by the uniqueness of (λ, κ) -saturated models there is an isomorphic f from $N_{\xi+1} = N'_{\omega(\xi+1)}$ onto $N_\kappa = N'_{\omega\kappa}$ over $N_\xi = N'_{\omega\kappa}$, and $f(p \upharpoonright N_{\xi+1}) = p$ is proved as above. $\square_{7.3}$

7.4 Claim. 1) If $M_0 \leq_{\mathfrak{K}} M_1$ are in K_λ and the types $p_\ell \in \mathcal{S}(M_\ell)$ are minimal reduced, for $\ell = 0, 1$ and $p_0 = p_1 \upharpoonright M_0$ then p_0, p_1 are conjugate; (i.e. there is an isomorphic f from M_0 onto M_1 such that $f(p_0) = p_1$).
 2) If in addition $M \leq_{\mathfrak{K}} M_0$ and M_0, M_1 are (λ, κ) -saturated over M , then p_0, p_1 are conjugate over N .

Remark. Note that p minimal (or reduced) implies that p is not algebraic.

Proof. 1) Let $\langle (N_i^0, N_i^1, a) : i < \lambda^+ \rangle$ and E be as in the proof of 7.3 and $\kappa = \text{cf}(\kappa) \leq \lambda$. For each $\delta \in S_\kappa =: \{\alpha < \lambda^+ : \alpha \in E \text{ and } \text{cf}(\alpha) = \kappa\}$, and minimal reduced $p \in \mathcal{S}(N_\delta^0)$, we know that for some $i_p < \delta$, $p \upharpoonright N_{i_p}^0$ is minimal reduced [why? by 7.3(1),(2)] and some $q_p \in \mathcal{S}(N_{i_p}^0)$ is conjugate to p . For $\kappa = \text{cf}(\kappa) \leq \lambda$, $q \in \mathcal{S}(N_i^0)$, $i < \lambda^+$, $r \in \mathcal{S}(N_i^0)$ minimal let $A_{q,r}^{\kappa,i} = \{\delta < \lambda^+ : \text{there is a type } p \text{ such that } r \subseteq p \in \mathcal{S}(N_\delta^0), p \text{ non-algebraic (this determines } p), p \text{ minimal reduced, } i_p = i, q_p = q \text{ (and clearly } p \upharpoonright N_i^0 = r)\}\}.$

Next let

$$E_1 = \left\{ \delta < \lambda^+ : \text{for every } \kappa = \text{cf}(\kappa) \leq \lambda, \right. \\ \left. r, q \in \mathcal{S}(N_i^0) \text{ and } i < \delta, \text{ if } A_{q,r}^{\kappa,i} \text{ is well defined and} \right. \\ \left. \text{unbounded in } \lambda^+ \text{ then it is unbounded in } \delta \right\}.$$

So if $\delta_1 \in E_1$, $\kappa = \text{cf}(\delta_1)$, $p_1 \in \mathcal{S}(N_{\delta_1}^0)$ is minimal reduced, then we can find $\delta_0 < \delta_1$, $\text{cf}(\delta_0) = \kappa$, and $p_0 \in \mathcal{S}(N_{\delta_0}^0)$ minimal reduced with $q_{p_1} = q_{p_0}$, $i_{p_1} = i_{p_0}$, $p_0 \upharpoonright N_{i_{p_0}}^0 = p_1 \upharpoonright N_{i_{p_1}}^0$ call it r , it is necessarily minimal.

As p_1, p_0 extend r , $N_{i_{p_0}}^0 = N_{i_{p_1}}^0 \leq_{\mathfrak{K}} N_{\delta_0}^0 \leq_{\mathfrak{K}} N_{\delta_1}^0$, necessarily $p_1 = p_0 \upharpoonright N_{\delta_0}^0$, and also they are both conjugate to $q_{p_0} = q_{p_1}$ hence they are conjugate. Next we prove

- (*) if $M_0 <_{\mathfrak{K}} M_1$ are in K_λ , M_1 is (λ, κ) -limit over M_0 , $p'_0 \in \mathcal{S}(M_0)$ is minimal reduced and $p'_0 \leq p'_1 \in \mathcal{S}(M_1)$, p'_1 non-algebraic, then p'_0, p'_1 are conjugate.

Above we have a good amount of free choice in choosing $p_1 \in \mathcal{S}(N_{\delta_1}^0)$ (it should be minimal and reduced) so we could have chosen p_1 to be conjugate to p'_0 (i.e. is in $\mathcal{S}_{p'_0}(N_{\delta_1}^0)$); now also the corresponding p_0 is conjugate to p_1 hence p_0 is conjugate to p'_0 , hence we can find an isomorphism f_0 from M_0 onto $N_{\delta_0}^0$, $f_0(p'_0) = p_0$, and extend it to an isomorphism f_1 from M_1 onto $N_{\delta_1}^0$, so necessarily $f(p'_1) = p_1$ (as p_1 is the unique non-algebraic extension). As p_0, p_1 are conjugate through f_1 also p'_0, p'_1 are conjugate. So (*) holds.

Now assume just $M_0 \leq_{\mathfrak{K}} M_1$ are in K_λ , $p_0 \in \mathcal{S}(M_0)$ minimal reduced, $p_1 \in \mathcal{S}(M_1)$ the unique non-algebraic extension of p_0 and it is reduced (and necessarily minimal). There is $M_2, M_1 \leq_{\mathfrak{K}} M_2 \in K_\lambda$, M_2 is (λ, κ) -limit over M_1 hence also over M_0

and let p_2 be the unique non-algebraic extension of p_1 in $\mathcal{S}(M_2)$ hence p_2 is also the unique non-algebraic extension of p_0 in $\mathcal{S}(M_2)$.

Using $(*)$ on (M_0, M_2, p_0, p_2) and on (M_1, M_2, p_1, p_2) and get that p_0, p_2 are conjugate and that p_1, p_2 are conjugate hence p_1, p_2 are conjugate, the required result.

2) Similar proof. $\square_{7.4}$

7.5 Claim. 1) Assume $M_1 \leq_{\mathfrak{K}} M_2$ are in K_λ and M_2 is (λ, κ) -saturated over M_1 . If $p_1 \in \mathcal{S}(M_1)$ is minimal and reduced, then p_2 , the unique non-algebraic extension of p_1 in $\mathcal{S}(M_2)$, is reduced (and, of course, minimal).

2) There is no need to assume “ p_1 reduced”.

Proof. 1) We can find $\langle N_i : i \leq \kappa \rangle$, an $\leq_{\mathfrak{K}}$ -increasingly continuous sequence in K_λ such that N_{i+1} is (λ, κ) -saturated over N_i and $N_\kappa = M_1$. So by 7.3(1),(2), for some $\zeta < \kappa$ we have: $p_2 \upharpoonright N_\zeta$ is minimal and for some isomorphism f from $N_{\zeta+1}$ onto N_κ we have $f(p_1 \upharpoonright N_\zeta) = p_1$ and $f \upharpoonright N_\zeta = \text{id}_{N_\zeta}$. Also $M_1, N_\zeta + 1$ are isomorphic over N_ζ (as both are (λ, κ) -saturated over it) hence there is an isomorphism g from $N_{\zeta+1}$ onto M_2 over N_ζ . Now $p_1 = f(p_1 \upharpoonright N_{\zeta+1})$ and $f_2 =: g(p_1 \upharpoonright N_{\zeta+1})$ are non-algebraic extensions of $p_1 \upharpoonright N_\zeta$ which is minimal, hence $p_1 = p_2 \upharpoonright M_1$ and p_2 is as mentioned in 7.5. Now $g \circ f^{-1}$ show that p_1, p_2 are conjugate so as p_1 is reduced also p_2 is reduced.

2) Easy as we can find $N, M_1 \leq_K N, q \in \mathcal{S}(N)$ minimal reduced; without loss of generality $N \leq_{\mathfrak{K}} M_2$ and M_2 is (λ, κ) -saturated over N , and apply part (1). $\square_{7.5}$

7.6 Claim. Assume

- (a) $N_{i,j} \in K_\lambda$ for $i \leq \delta_1, j \leq \delta_2$
- (b) $\langle N_{i,j} : j \leq \delta_2 \rangle$ is $\leq_{\mathfrak{K}}$ -increasingly continuous for each $i \leq \delta_1$
- (c) $\langle N_{i,j} : i < \delta_1 \rangle$ is $\leq_{\mathfrak{K}}$ -increasingly continuous for each $j \leq \delta_2$
- (d) $\langle N_{i,j} : i \leq \delta_1, j \in \delta_2 \rangle$ is smooth, i.e. $N_{i_1, j_1} \cap N_{i_2, j_2} = N_{\min\{i_1, i_2\}} \cap N_{\min\{j_1, j_2\}}$
- (e) $N_{i+1, j+1}$ is universal over $N_{i, j+1} \cup N_{i+1, j}$ (i.e. $N_{i+1, j+1}$ is universal over some $N'_{i+1, j+1}$ where $N_{i, j+1} \cup N_{i+1, j} \subseteq N'_{i+1, j+1}$),
- (f) δ_1 is divisible by $\text{cf}(\delta_2) \times \lambda \times \omega$
(and even easier if $\delta_1 = 1!$).

Then N_{δ_1, δ_2} is $(\lambda, \text{cf}(\delta_1))$ -saturated over N_{i, δ_1} for $i < \delta_1$.

Proof. Without loss of generality $\delta_2 = \text{cf}(\delta_2)$. (Why? let $\langle \alpha_\varepsilon : \varepsilon \leq \text{cf}(\delta_1) \rangle$ be increasingly continuous with limit δ_1 such that $[\varepsilon \text{ limit} \leftrightarrow \alpha_\varepsilon \text{ limit}]$, and use $N'_{i, \varepsilon} = N_{i, \alpha_\varepsilon}$).

For $i < \delta_1, j < \delta_2$ let $M_{i,j}, M'_{i,j}$ be such that $M_{i+1, j} \cup M_{i, j+1} \subseteq M_{i,j} \leq_{\mathfrak{K}} M'_{i,j} \leq M_{i+1, j+1}$ and $M'_{i,j}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $M_{i,j}$.

Now let $p \in \mathcal{S}(N)$ be minimal and reduced; and for $i \leq \delta_1, j \leq \delta_2$ let $p_{i,j} \in \mathcal{S}(N_{i,j})$ be the unique non-algebraic extension of p in $\mathcal{S}(N_{i,j})$ so it is minimal. Now for $i < \delta_1$, note that $\langle M'_{i+\varepsilon, \varepsilon} : \varepsilon < \delta_2 \rangle$ is $\leq_{\mathfrak{K}}$ -increasing (not continuous!) and $M'_{i+\varepsilon+1, \varepsilon+1}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $M'_{i+\varepsilon, \varepsilon}$ and $\bigcup_{\varepsilon < \delta} M'_{i+\varepsilon, \varepsilon} = \bigcup_{\varepsilon < \delta_1} N_{i+1, \varepsilon+1} = N_{i+\delta_2, \delta_2}$, and

$N_{0,0} \leq_{\mathfrak{k}} M'_{1,0}$, hence by 7.5 we know that $p_{i+\delta_2, \delta_2}$ is reduced (and minimal). In fact, similarly $\alpha < \delta_1$ & $\text{cf}(\alpha) = \text{cf}(\delta_2) \Rightarrow p_{\alpha, \delta_2}$ is reduced. As $N_{i+1, j+1} \neq N_{i+1, j} \cup N_{i, j+1}$ and clause (d) (smoothness) necessarily $N_{i+\delta_2, \delta_2} < N_{i+\delta_2+1, \delta_2}$, hence some $c \in N_{i+\delta_2+i, \delta_2} \setminus N_{i+\delta_2, \delta_2}$ realizes $p_{i+\delta_2, \delta_2}$. So if $\alpha \leq \delta_1$ is divisible by $\delta_2 \times \lambda$ and has cofinality $\text{cf}(\delta_2)$ and $\beta < \alpha$, then by 5.6 N_{α, δ_2} is universal over N_{β, δ_2} . As δ_1 is divisible by $\text{cf}(\delta_2) \times \lambda \times \omega$ we are done. $\square_{7.6}$

7.7 Lemma. 1) For every $N \in K_{\lambda^+}$ we can find a representation

$\bar{N} = \langle N_i : i < \lambda^+ \rangle$, with N_{i+1} being $(\lambda, \text{cf}(\lambda))$ -saturated over N_i .

2) If for $\ell = 1, 2$ we have $N^\ell = \langle N_i^\ell : i < \lambda^+ \rangle$ as in part (1) then there is an isomorphism f from N^1 onto N^2 mapping N_i^1 onto N_i^2 for each $i < \lambda^+$. Moreover for any $i < \lambda^+$ and isomorphism g from N_i^1 onto N_i^2 we can find an isomorphism f from N^1 onto N^2 extending g and mapping N_j^1 onto N_j^2 for each $j \in [i, \lambda^+)$.

3) If $N^0 \leq_{\mathfrak{K}} N^1$ are in K_{λ^+} then we can find representation $\bar{N}^\ell$ of N^ℓ as in (1) with $N_i^0 = N^0 \cap N_i^1$, (so $N_i^0 \leq_{\mathfrak{K}} N_i^1$).

4) For any strictly increasing function $\mathbf{f} : \lambda^+ \rightarrow \lambda^+$, we can find $N_{i, \varepsilon}$ for $i < \lambda^+, \varepsilon \leq \lambda \times (1 + \mathbf{f}(i))$ such that:

- (a) $N_{i, \varepsilon} \in K_\lambda$
- (b) $\langle N_{i, \varepsilon} : \varepsilon \leq \lambda \times (1 + \mathbf{f}(i)) \rangle$ is strictly $\leq_{\mathfrak{K}}$ -increasing continuous
- (c) for each ε , $\langle N_{i, \varepsilon} : i \in [i_\varepsilon, \lambda^+) \rangle$ is a representation as in (1) where $i_\varepsilon = \text{Min}\{i : \varepsilon \leq \lambda \times (1 + \mathbf{f}(i))\}$
- (d) if $\varepsilon < \lambda \times (1 + \mathbf{f}(i))$ and $i < j < \lambda^+$ then $N_{j, \varepsilon} \cap N_{i, \lambda \times (1 + \mathbf{f}(i))} = N_{i, \varepsilon}$
- (e) $N_{i+1, \varepsilon+1}$ is $(\lambda, \aleph_0)$ -saturated over $N_{i+1, \varepsilon} \cup N_{i, \varepsilon+1}$.

Proof. Straight.

4) First use $\mathbf{f}' : \lambda^+ \rightarrow \lambda^+$ which is $\mathbf{f}'(i) = \lambda^\omega \times \mathbf{f}(i)$. Then define the $N_{i, \varepsilon}$ ($\varepsilon < \lambda \times (1 + \mathbf{f}'(i); i < \lambda^+)$). “Forget” about “ $N_{i+1, \varepsilon}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $N_{i, \varepsilon}$ ”, remember we have disjoint amalgamation by 5.11. Now by 7.6, even for ε limit divisible by λ^3 we get $N_{i+\lambda, \varepsilon}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $N_{i, \varepsilon}$, so renaming all is O.K.). $\square_{7.7}$

We can deduce from 7.5, but to keep the door open to other uses we shall not use

7.8 Claim. If $\kappa_\ell = \text{cf}(\kappa_\ell) \leq \lambda$, and N_ℓ is (λ, κ_ℓ) -saturated over N for $\ell = 1, 2$ then N_1, N_2 are isomorphic over N .

Proof. We can define by induction on $i \leq \lambda \times \kappa_1$, and then by induction on $j \leq \lambda \times \kappa_2, M_{i, j}$ such that:

- (a) $M_{i, j} \in K_\lambda$
- (b) $M_{0, 0} = N$
- (c) $i_1 \leq i$ & $j_1 \leq j \Rightarrow M_{i_1, j_1} \leq_{\mathfrak{K}} M_{i, j}$
- (d) $M_{i_1, j_1} \cap M_{i_2, j_2} = M_{\min\{i_1, i_2\}, \min\{j_1, j_2\}}$
- (e) $M_{i, j}$ is $\leq_{\mathfrak{K}}$ -increasing continuous in i
- (f) $M_{i, j}$ is $<_{\mathfrak{K}}$ -increasing continuous in j

- (g) $M_{0,j} \neq M_{0,j+1}$
- (h) $M_{i+1,0} \neq M_{i,0}$
- (i) $M_{i+1,j+1}$ is universal over $M_{i+1,j} \cup M_{i,j+1}$.

There is no problem by 5.11(1) (using the existence of disjoint amalgamation).

Now $M_{\lambda \times \kappa_1, \lambda \times \kappa_2}$ is the union of the strictly $<_{\mathfrak{K}}$ -increasing sequence $\langle M_{0,0} \rangle^\wedge \langle M_{\lambda \times i, \lambda \times \kappa_2} : i < \kappa_1 \rangle$ hence by 7.6 is (λ, κ_1) -saturated over $M_{0,0} = N$ hence $M_{\lambda \times \kappa_1, \lambda \times \kappa_2} \cong_N N_1$. Similarly $M_{\lambda \times \kappa_1, \lambda \times \kappa_2}$ is the union of the strictly $\leq_{\mathfrak{K}}$ -increasing sequence $\langle M_{0,0} \rangle^\wedge \langle M_{\lambda \times \kappa_1, \lambda \times j} : j < \kappa_2 \rangle$ hence is (λ, κ_2) -saturated over $M_{0,0} = N$, hence $M_{\lambda \times \kappa_1, \lambda \times \kappa_2} \cong_N N_2$. Together N_1, N_2 are isomorphic over N .

□_{7.8}

§8 UNIQUENESS OF AMALGAMATION IN $\mathfrak{K}_\lambda$

We deal in this section only with K_λ .

We want to, at least, approximate unique amalgamation using as starting point $\otimes$ of 6.4 (see also 6.7), i.e. $K_\lambda^{3,\text{uq}} \neq \emptyset$.

8.1 Hypothesis. 1) Assume hypothesis 7.1, so

- (a) $\mathfrak{K}$ abstract elementary class
- (b) $\mathfrak{K}$ has amalgamation in λ
- (c) $\mathfrak{K}$ is categorical in λ (can be weakened)
- (d) $\mathfrak{K}$ is stable in λ (see 5.7, clause (a))
- (e) there is an inevitable $p \in \mathcal{S}(N)$ for $N \in K_\lambda$ (holds by 5.3)
- (f) the basic properties in type theory.

2) (M^*, N^*) is some pair in

$K_\lambda^{3,\text{uq}} = \{(M_0, M_2) : M_0 \leq_{\mathfrak{K}} M_2 \text{ are in } K_\lambda \text{ and for every } M_1, M_0 \leq_{\mathfrak{K}} M_1 \in K_\lambda \Rightarrow (M_0, M_1, M_2) \in K_\lambda^{\text{uq}}; \text{ equivalently for some } M_1, (M_0, M_1, M_2) \text{ are as in } \otimes \text{ of 6.4}\}$ (eventually the choice does not matter; if each time instead of $\cong (M^*, N^*)$ we write $\in K_\lambda^{2,\text{uq}}$, see 8.11; but if we start with this definition then the uniqueness theorems will be more cumbersome).

8.2 Definition. Assume $\bar{\delta} = \langle \delta_1, \delta_2, \delta_3 \rangle$, δ_2 a limit ordinal $< \lambda^+$ but δ_1, δ_3 are $< \lambda^+$ and may be 1. We say that $NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$ (we say N_1, N_2 are saturated and smoothly amalgamated in N_3 over N_0 for $\bar{\delta}$) when:

- (a) $N_\ell \in K_\lambda$ for $\ell \in \{0, 1, 2, 3\}$
- (b) $N_0 \leq_{\mathfrak{K}} N_\ell \leq_{\mathfrak{K}} N_3$ for $\ell = 1, 2$
- (c) $N_1 \cap N_2 = N_0$ (i.e. in disjoint amalgamation)
- (d) N_1 is $(\lambda, \text{cf}(\delta_1))$ -saturated over N_0
- (e) N_2 is $(\lambda, \text{cf}(\delta_2))$ -saturated over N_0 , if $\delta_1 = 1$ this means just $N_0 \leq_{\mathfrak{K}} N_2$
- (f) there are $N_{1,i}, N_{2,i}$ for $i \leq \lambda \times \delta_1$ (called the witness) such that:
 - (α) $N_{1,0} = N_0, N_{1, \lambda \times \delta_1} = N_1$
 - (β) $N_{2,0} = N_2$
 - (γ) $\langle N_{\ell,i} : i \leq \lambda \times \delta_1 \rangle$ is $<_{\mathfrak{K}}$ -increasing continuous for $\ell = 1, 2$
 - (δ) $(N_{1,i}, N_{1,i+1}) \cong (M^*, N^*)$
 - (ε) $N_{2,i} \cap N_1 = N_{1,i}$
 - (ζ) N_3 is $(\lambda, \text{cf}(\delta_3))$ -saturated over $N_{2, \lambda \times \delta_1}$; if $\delta_3 = 1$ this means just $N_{2, \lambda \times \delta_1} \leq_{\mathfrak{K}} N_3$

Discussion: Why this definition of NF? We need a nonforking notion with the usual properties. We first describe a version depending on $\langle \delta_0, \delta_1, \delta_2 \rangle$ and get $NF = NF_{\lambda, \bar{\delta}}$, $\bar{\delta}$ works like a scaffold – eventually $\bar{\delta}$ disappears.

8.3 Definition. 1) We say $N_1 \bigcup_{N_0}^{N_3} N_2$ (or N_1, N_2 are smoothly amalgamated over N_0 inside N_3 or $NF_\lambda(N_0, N_1, N_2, N_3)$) iff we can find $M_\ell \in K_\lambda$ (for $\ell < 4$) such that:

- (a) $NF_{\lambda, \langle \lambda, \lambda, \lambda \rangle}(M_0, M_1, M_2, M_3)$
- (b) $N_\ell \leq_{\mathfrak{K}} M_\ell$ for $\ell < 4$
- (c) $N_0 = M_0$
- (d) M_1, M_2 are $(\lambda, \text{cf}(\lambda))$ -saturated over N_0 (follows by (a) see clauses (d), (e) of 8.2).

8.4 Claim. 1) If $\bar{\delta} = \langle \delta_1, \delta_2, \delta_3 \rangle, \delta_\ell$ a limit ordinal $< \lambda^+$ and $N_\ell \in K_\lambda$ for $\ell < 3$, and N_1 is $(\lambda, \text{cf}(\delta_1))$ -saturated over N_0 and N_2 is $(\lambda, \text{cf}(\delta_2))$ -saturated over N_0 and $N_0 \leq_{\mathfrak{K}} N_1, N_0 \leq_{\mathfrak{K}} N_2$ and for simplicity $N_1 \cap N_2 = N_0$. Then we can find N_3 such that $NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$.

2) Moreover, we can choose any $\langle N_{1,i} : i \leq \lambda \times \delta_1 \rangle$ as in 8.2(f)(α), (γ), (δ) as part of the witness.

Proof. Straight (remembering 7.5 (and uniqueness of the $(\lambda, \text{cf}(\delta_1))$ -saturated model over N_0)). $\square_{8.4}$

8.5 Claim. If Definition 8.2, if δ_3 is a limit ordinal, then without loss of generality (even without changing $\langle N_{1,i} : i \leq \lambda \times \delta_1 \rangle$)

- (g) $N_{2,i+1}$ is $(\lambda, \text{cf}(\delta_2))$ -saturated over $N_{i+1}^1 \cup N_i^2$ (which means it is $(\lambda, \text{cf}(\delta_2))$ -saturated over some N , where $N_{i+1}^1 \cup N_i^2 \subseteq N \leq_{\mathfrak{K}} N_{2,i+1}$).

Proof. So assume $NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$ holds as witnessed by $\langle N_{\ell,i} : i \leq \lambda \times \delta_\ell \rangle$ for $\ell = 1, 2$. Now we choose by induction on $i \leq \lambda \times \delta_1$ a model $M_{2,i} \in K_\lambda$ such that:

- (i) $N_{2,i} \leq M_{2,i}$
- (ii) $M_{2,0} = N_2$
- (iii) $M_{2,i}$ is $\leq_{\mathfrak{K}}$ -increasing continuous
- (iv) $M_{2,i} \cap N_{2, \lambda \times \delta_1} = N_{2,i}$ moreover $M_{2,i} \cap N_3 = N_{2,i}$
- (v) $M_{2,i+1}$ is $(\lambda, \text{cf}(\delta_2))$ -saturated over $M_{2,i} \cup N_{2,i+1}$.

There is no problem to carry the definition. Let M_3 be such that $M_{2, \lambda \times \delta_1} \leq_{\mathfrak{K}} M_3 \in K_\lambda$ and M_3 is $(\lambda, \text{cf}(\delta_3))$ -saturated over $M_{2, \lambda \times \delta_1}$. So both M_3 and N_3 are $(\lambda, \text{cf}(\delta_3))$ -saturated over $N_{2, \lambda \times \delta_1}$, hence they are isomorphic over $N_{2, \lambda \times \delta_1}$ so let f be an isomorphism from M_3 onto N_3 which is the identity over $N_{2, \lambda \times \delta_1}$.

Clearly $\langle N_{1,i} : i \leq \lambda \times \delta_1 \rangle, \langle f(M_{2,i}) : i \leq \lambda \times \delta_1 \rangle$ are also witnesses for

$NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$ satisfying the extra demand (g). $\square_{8.5}$

8.6 Claim. (*Weak Uniqueness*).

Assume that for $x \in \{a, b\}$, we have $NF_{\lambda, \bar{\delta}^x}(N_0^x, N_1^x, N_2^x, N_3^x)$ holds as witnessed by $\langle N_{1,i}^x : i \leq \lambda \times \delta_1^x \rangle, \langle N_{2,i}^x : i \leq \lambda \times \delta_1^x \rangle$ and $\delta_1 =: \delta_1^a = \delta_1^b$, $cf(\delta_2^a) = cf(\delta_2^b)$ and $cf(\delta_3^a) = cf(\delta_3^b) \geq \aleph_0$.

Suppose further that f_ℓ is an isomorphism from N_ℓ^a onto N_ℓ^b for $\ell = 0, 1, 2$, moreover: $f_0 \subseteq f_1, f_0 \subseteq f_2$ and $f_1(N_{1,i}^a) = N_{1,i}^b$.

Then we can find an isomorphism f from N_3^a onto N_3^b extending $f_1 \cup f_2$.

Proof. Without loss of generality $N_{2,i+1}^x$ is $(\lambda, cf(\delta_2))$ -saturated over $N_{1,i+1}^x \cup N_{2,i}^x$ (by 8.5, note “without changing the $N_{1,i}$ ’s). Now we choose by induction on $i \leq \lambda \times \delta_1$ an isomorphism g_i from $N_{2,i}^a$ onto $N_{2,i}^b$ such that: g_i is increasing in i and g_i extends $(f_1 \upharpoonright N_{1,i}^a) \cup f_2$.

For $i = 0$ choose $g_0 = f_2$ and for i limit let g_i be $\bigcup_{j < i} g_j$ and for $i = j + 1$ use

$(N_{1,i}, N_{1,i+1}) \cong (M^*, N^*)$ (see 8.2) and the extra saturation clause (g). Now we can extend $g_{\lambda \times \delta_1}$ to an isomorphism from N_3^a onto N_3^b as N_3^x is $(\lambda, cf(\delta_3))$ -saturated from $N_{2,\lambda \times \delta_1}^x$ (for $x \in \{a, b\}$); note that knowing 8.6 possibly the choice of $\langle N_{1,i} : i \leq \lambda \times \delta_1 \rangle$ matters. $\square_{8.6}$

Now we prove an “inverted” uniqueness

8.7 Claim. *Suppose that*

- (a) for $x \in \{a, b\}$ we have $NF_{\lambda, \bar{\delta}^x}(N_0^x, N_1^x, N_2^x, N_3^x)$
- (b) $\bar{\delta}^x = \langle \delta_1^x, \delta_2^x, \delta_3^x \rangle, \delta_1^a = \delta_2^b, \delta_2^a = \delta_1^b, cf(\delta_3^a) = cf(\delta_3^b)$ all limit ordinals
- (c) f_0 is an isomorphism from N_0^a onto N_0^b
- (d) f_1 is an isomorphism from N_1^a onto N_2^b
- (e) f_2 is an isomorphism from N_2^a onto N_1^b
- (f) $f_0 \subseteq f_1$ and $f_0 \subseteq f_2$.

Then there is an isomorphism from N_3^a onto N_3^b extending $f_1 \cup f_2$.

Before proving

8.8 Subclaim. 1) For any limit ordinals $\delta_1^a, \delta_2^a, \delta_3^a \leq \lambda$ we can find $M_{i,j}$ (for $i \leq \lambda \times \delta_1^a$ and $j \leq \lambda \times \delta_2^a$) and M_3 such that:

- (A) $M_{i,j} \in K_\lambda$
- (B) $i_1 \leq i_2 \ \& \ j_1 \leq j_2 \Rightarrow M_{i_1,j_1} \leq_{\aleph} M_{i_2,j_2}$
- (C) if $i \leq \lambda \times \delta_1$ is a limit ordinal and $j \leq \lambda \times \delta_2$ then $M_{i,j} = \bigcup_{\zeta < i} M_{\zeta,j}$
- (D) if $i \leq \lambda \times \delta_1$ and $j \leq \lambda \times \delta_2$ is a limit ordinal then $M_{i,j} = \bigcup_{\xi < j} M_{i,\xi}$
- (E) $(M_{0,j}, M_{0,j+1}) \cong (M^*, N^*)$
- (F) $(M_{i,0}, M_{i+1,0}) \cong (M^*, N^*)$

(G) for $i_1, i_2 \leq \lambda \times \delta_1$ and $j_1, j_2 \leq \lambda \times \delta_2$ we have

$$M_{i_1, j_1} \cap M_{i_2, j_2} = M_{\min\{i_1, i_2\}, \min\{j_1, j_2\}},$$

(H) $M_{\lambda \times \delta_1, \lambda \times \delta_2} \leq_{\mathfrak{K}} M_3 \in K_\lambda$ moreover

M_3 is $(\lambda, \text{cf}(\delta_3))$ -saturated over $M_{\lambda \times \delta_1, \lambda \times \delta_2}$.

2) Moreover, it is O.K. if $\langle M_{0,j} : j \leq \lambda \times \delta_2^a \rangle, \langle M_{i,0} : i \leq \lambda \times \delta_1^a \rangle$ are pregiven as long as both $\leq_{\mathfrak{K}}$ -increasing continuous in $\mathfrak{K}_\lambda$ satisfying (E) + (F) and $M_{0, \lambda \times \delta_2^a} \cap M_{\lambda \times \delta_1^a, 0} = M_{0,0}$.

Proof. 1) This is done by induction on i and for fixed i by induction on j . For $i = 0, j = 0$ let $M_{0,0} \in K_\lambda$ be arbitrary, for $i = 0, j$ limit use clause (D), for $i = 0, j = \xi + 1$ use clause (E). For i limit use clause (C) (and check). For $i = \zeta + 1$, if $j = 0$ use clause (F), if j limit use clause (D) and if $j = \xi + 1$ use the existence of disjoint amalgamation (i.e. 5.11).

Lastly, choose $M_3 \in K_\lambda$ which is $(\lambda, \text{cf}(\delta_3^a))$ -saturated over $M_{\lambda \times \delta_1^a, \lambda \times \delta_2^a}$.

2) Similarly. $\square_{8.8}$

Proof of 8.7. Let $M_{i,j}, M_3$ be as in 8.8. For $x \in \{a, b\}$ as $NF_{\lambda, \bar{\delta}^x}(N_0^x, N_1^x, N_2^x, N_3^x)$, we know that there are witnesses $\langle N_{1,i}^x : i \leq \lambda \times \delta_1^x \rangle, \langle N_{2,i}^x : i \leq \lambda \times \delta_1^x \rangle$ for this, so $\langle N_{1,i}^x : i \leq \lambda \times \delta_1^x \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous and $(N_{1,i}^x, N_{1,i+1}^x) \cong (M^*, N^*)$. So $\langle N_{1,i}^a : i \leq \lambda \times \delta_1^a \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous sequences with each successive pair isomorphic to (M^*, N^*) hence by 8.8(2) without loss of generality there is an isomorphism g_1 from $N_{1, \lambda \times \delta_1^a}^a$ onto $M_{\lambda \times \delta_1^a}$, mapping $N_{1,i}^a$ onto $M_{i,0}$; remember $N_{1, \lambda \times \delta_1^a}^a = N_1^a$. Let $g_0 = g_1 \upharpoonright N_0^a = g_1 \upharpoonright N_{1,0}^a$ so $g_0 \circ f_0^{-1}$ is an isomorphism from N_0^b onto $M_{0,0}$.

As $\delta_1^b = \delta_2^a$, using 8.8(2) fully without loss of generality there is an isomorphism g_2 from $N_{1, \lambda \times \delta_2^a}^b$ onto $M_{0, \lambda \times \delta_2^a}$ mapping $N_{1,j}^b$ onto $M_{0,j}$ (for $j \leq \lambda \times \delta_2^a$) and g_2 extends $g_0 \circ f_0^{-1}$.

Now we want to use the weak uniqueness 8.6 and for this note:

- (α) $NF_{\lambda, \bar{\delta}^a}(N_0^a, N_1^a, N_2^a, N_3^a)$ as witnessed by $\langle N_{1,i}^a : i \leq \lambda \times \delta_1^a \rangle, \langle N_{2,i}^a : i \leq \lambda \times \delta_1^a \rangle$
[why? an assumption]
- (β) $NF_{\lambda, \bar{\delta}^a}(M_{0,0}, M_{\lambda \times \delta_1^a, 0}, M_{0, \lambda \times \delta_2^a}, M_3)$ as witnessed by the sequences $\langle M_{i,0} : i \leq \lambda \times \delta_1^a \rangle, \langle M_{i, \lambda \times \delta_2^a} : i \leq \lambda \times \delta_2^a \rangle$
[why? check]
- (γ) g_0 is an isomorphism from N_0^a onto $M_{0,0}$
[why? see its choice]
- (δ) g_1 is an isomorphism from N_1^a onto $M_{\lambda \times \delta_1^a, 0}$ mapping $N_{1,i}^a$ onto $M_{i,0}$ for $i \leq \lambda \times \delta_1^a$ and extending g_0
[why? see the choice of g_1 and of g_0]
- (ε) $g_2 \circ f_2$ is an isomorphism from N_2^a onto $M_{0, \lambda \times \delta_2^a}$ extending g_0
[why? f_2 is an isomorphism from N_2^a onto N_1^b and g_2 is an isomorphism from N_1^b onto $M_{0, \lambda \times \delta_1^a}$ extending $g_0 \circ f_0^{-1}$ and $f_0 \subseteq f_2$].

So there is by 8.6 an isomorphism g_3^a from N_3^a onto M_3 extending g_1 and $g_2 \circ f_2$.

We next want to apply 8.6 to the N_1^b 's; so note:

- (α)' $NF_{\lambda, \bar{\delta}^b}(N_0^b, N_1^b, N_2^b, N_3^b)$ as witnessed by $\langle N_{1,i}^b : i \leq \lambda \times \delta_2^a \rangle$,
 $\langle N_{2,i}^b : i \leq \lambda \times \delta_2^a \rangle$
- (β)' $NF_{\lambda, \bar{\delta}^b}(M_{0,0}, M_{0,\lambda \times \delta_2^a}, M_{\lambda \times \delta_1^a, 0}, M_3)$ as witnessed by the sequences
 $\langle M_{0,j} : j \leq \lambda \times \delta_2^a \rangle, \langle M_{\lambda \times \delta_1^a, j} : j \leq \lambda \times \delta_1^a \rangle$
- (γ)' $g_0 \circ (f_0)^{-1}$ is an isomorphism from N_0^b onto $M_{0,0}$
 [why? Check.]
- (δ)' g_2 is an isomorphism from N_1^b onto $M_{0,\lambda \times \delta_2^a}$ mapping $N_{1,j}^b$ onto $M_{0,j}$ for
 $j \leq \lambda \times \delta_2^a$ and extending $g_0 \circ (f_1)^{-1}$
 [why? see the choice of g_2 : it maps $N_{1,j}^b$ onto $M_{0,j}$]
- (ε)' $g_1 \circ (f_1)^{-1}$ is an isomorphism from N_2^b onto $M_{\lambda \times \delta_1^a, 0}$ extending g_0
 [why? remember f_1 is an isomorphism from N_1^a onto N_2^b extending f_0 and
 the choice of g_1 : it maps N_1^a onto $M_{\lambda \times \delta_1^a, 0}$].

So there is an isomorphism g_3^b from N_3^b onto M_3 extending $g_2, f_1 \circ (f_1)^{-1}$.

Lastly $(g_3^b)^{-1} \circ g_3^a$ is an isomorphism from N_3^a onto N_3^b (chase arrows). Also

$$\begin{aligned}
 ((g_3^b)^{-1} \circ g_3^a) \upharpoonright N_1^a &= (g_3^b)^{-1}(g_3^a \upharpoonright N_1^a) \\
 &= (g_3^b)^{-1}g_1 = ((g_3^b)^{-1} \upharpoonright M_{\lambda \times \delta_1^a, 0}) \circ g_1 \\
 &= (g_3^b \upharpoonright N_2^b)^{-1} \circ g_1 = ((g_1 \circ (f_1)^{-1})^{-1}) \circ g_1 \\
 &= (f_1 \circ (g_1)^{-1}) \circ g_1 = f_1.
 \end{aligned}$$

Similarly $((g_3^b)^{-1} \circ g_3^a) \upharpoonright N_2^a = f_2$.

So we have finished. □_{8.7}

8.9 Claim. [Uniqueness]. Assume for $x \in \{a, b\}$ we have

$NF_{\lambda, \bar{\delta}^x}(N_0^x, N_1^x, N_2^x, N_3^x)$ and $cf(\delta_1^a) = cf(\delta_1^b), cf(\delta_2^a) = cf(\delta_2^b), cf(\delta_3^a) = cf(\delta_3^b)$, all δ_ℓ^x limit ordinals.

If f_ℓ is an isomorphism from N_ℓ^a onto N_ℓ^b for $\ell < 3$ and $f_0 \subseteq f_1, f_0 \subseteq f_2$ then there is an isomorphism f from N_3^a onto N_3^b extending f_1, f_2 .

Proof. Let $\bar{\delta}^c = \langle \delta_1^c, \delta_2^c, \delta_3^c \rangle = \langle \delta_2^a, \delta_1^a, \delta_3^a \rangle$; by 8.4 there are N_ℓ^c (for $\ell \leq 3$) such that $NF_{\lambda, \bar{\delta}^c}(N_0^c, N_1^c, N_2^c, N_3^c)$. There is for $x \in \{a, b\}$ an isomorphism g_0^x from N_0^x onto N_0^c (as K_λ is categorical in λ) and without loss of generality $g_0^b = g_0^a \circ f_0$. Similarly for $x \in \{a, b\}$ there is an isomorphism g_1^x from N_1^x onto N_2^c extending g_0^x (as N_1^x is $(\lambda, cf(\delta_1^x))$ -saturated over N_0^x and also N_2^c is $(\lambda, cf(\delta_2^c))$ -saturated over N_0^c and $cf(\delta_2^c) = cf(\delta_1^a) = cf(\delta_1^b)$) and without loss of generality $g_1^b = g_1^a \circ f_1$. Similarly for $x \in \{a, b\}$ there is an isomorphism g_2^x from N_2^x onto N_1^c extending g_0^x (as N_2^x is $(\lambda, cf(\delta_2^x))$ -saturated over N_0^x and also N_1^c is $(\lambda, cf(\delta_1^c))$ -saturated over N_0^c and $cf(\delta_1^c) = cf(\delta_2^a) = cf(\delta_2^b)$) and without loss of generality $g_2^b = g_2^a \circ f_2$.

So by 8.7 for $x \in \{a, b\}$ there is an isomorphism g_3^x from N_3^x onto N_3^c extending g_1^x

and g_2^x . Now $(g_3^b)^{-1} \circ g_3^a$ is an isomorphism from N_3^a onto N_3^b extending f_1, f_2 as required. $\square_{8.9}$

8.10 Conclusion. [Symmetry].

If $NF_{\lambda, \langle \delta_1, \delta_2, \delta_3 \rangle}(N_0, N_1, N_2, N_3)$ then $NF_{\lambda, \langle \delta_2, \delta_1, \delta_3 \rangle}(N_0, N_2, N_1, N_3)$.

Proof. By 8.7 (and 8.9).

8.11 Claim. 1) In Definition 8.2 we can replace $(N_{1,i}, N_{1,i+1}) \cong (M^*, N^*)$ by $(N_{1,i}, N_{1,i+1}) \in K^{3, \text{uq}}$.

Proof. Like the proof of 8.7 (get $(M_{i,0}, M_{i+1,0}) \cong (M^*, N^*)$, $(M_{0,j}, M_{0,j+1}) \in K_\lambda^{3, \text{uq}}$), but as we shall not use it, we do not elaborate. $\square_{8.11}$

Now we turn to smooth amalgamation (not necessarily saturated, see Definition 8.3).

8.12 Claim. 1) If $NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$ each δ_ℓ limit then $NF_\lambda(N_0, N_1, N_2, N_3)$ (see Definition 8.3).

2) In Definition 8.3 we can add:

- (d)⁺ M_ℓ is $(\lambda, \text{cf}(\lambda))$ -saturated over N_0 and moreover over N_ℓ ,
- (e) M_3 is $(\lambda, \text{cf}(\lambda))$ -saturated over $M_1 \cup M_2$ (actually this is given by (f)(ζ) of Definition 8.2).

Proof. 1) By 8.8 we can find $M_{i,j}$ for $i \leq \lambda \times (\delta_1 + \lambda), j \leq \lambda \times (\delta_2 + \lambda)$ for $\bar{\delta}' = \langle \delta_1 + \lambda, \delta_2 + \lambda, \delta_3 \rangle$ and choose $M'_3 \in K_\lambda$ which is $(\lambda, \text{cf}(\delta_3))$ -saturated over $M_{\lambda \times \delta_1, \lambda \times \delta_2}$. So $NF_{\lambda, \bar{\delta}}(M_{0,0}, M_{\lambda \times \delta_1, 0}, M_{0, \lambda \times \delta_2}, M_3)$, hence by 8.9 without loss of generality $M_{0,0} = N_0, M_{\lambda \times \delta_1, 0} = N_1, M_{0, \lambda \times \delta_2} = N_2$, and $N_3 = M'_3$. Lastly, let M_3 be $(\lambda, \text{cf}(\lambda))$ -saturated over M'_3 . Now clearly also $NF_{\lambda, \langle \delta_1 + \lambda, \delta_2 + \lambda, \delta_3 + \lambda \rangle}(M_{0,0}, M_{\lambda \times (\delta_1 + \lambda), 0}, M_{0, \lambda \times (\delta_2 + \lambda)}, M_3)$ and $N_0 = M_{0,0}, N_1 = M_{\lambda \times \delta_2, 0} \leq_{\mathfrak{K}} M_{\lambda \times (\delta_2 + \lambda), 0}, N_2 = M_{0, \lambda \times \delta_2} \leq_{\mathfrak{K}} M_{0, \lambda \times (\delta_2 + \lambda)}$ and $M_{\lambda \times (\delta_1 + \lambda), 0}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $M_{\lambda \times \delta_1, 0}$ and $M_{0, \lambda \times (\delta_2 + \lambda)}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $M_{0, \lambda \times \delta_2}$ and $N_3 = M'_3 \leq_{\mathfrak{K}} M_3$. So we get all the requirements for $NF_\lambda(N_0, N_1, N_2, N_3)$ (as witnessed by $\langle M_{0,0}, M_{\lambda \times (\delta_1 + \lambda), 0}, M_{0, \lambda \times (\delta_2 + \lambda)}, M_3 \rangle$).

2) Similar proof. $\square_{8.12}$

8.13 Claim. [Uniqueness of smooth amalgamation]: If $NF_\lambda(N_0^x, N_1^x, N_2^x, N_3^x)$ for $x \in \{a, b\}$, f_ℓ an isomorphism from N_ℓ^a onto N_ℓ^b for $\ell < 3$ and $f_0 \subseteq f_1, f_0 \subseteq f_2$ then $f_1 \cup f_2$ can be extended to a $\leq_{\mathfrak{K}}$ -embedding of N_3^a into some $\leq_{\mathfrak{K}}$ -extension of N_3^b (so if N_3^x is (λ, κ) -saturated over $N_1^x \cup N_2^x$ for $x = a, b$, we can extend $f_1 \cup f_2$ to an isomorphism from N_3^a onto N_3^b).

Proof. For $x \in \{a, b\}$ let the sequence $\langle M_\ell^x : \ell < 4 \rangle$ be a witness to $NF_\lambda(N_0^x, N_1^x, N_2^x, N_3^x)$ as in 8.3, 8.12(2) so in particular

$NF_{\lambda, \langle \lambda, \lambda, \lambda \rangle}(M_0^x, M_1^x, M_2^x, M_3^x)$. By chasing arrows and uniqueness, i.e. 8.7 without loss of generality $M_\ell^a = M_\ell^b$ for $\ell < 4$ and $f_0 = \text{id}_{N_0^a}$. As M_1^a is $(\lambda, \text{cf}(\lambda))$ -saturated over N_1^a and also over N_1^b and f_1 is an isomorphism from N_1^a onto N_1^b , clearly there is an automorphism g_1 of M_1^a such that $f_1 \subseteq g_1$, hence also $\text{id}_{N_0^a} = f_0 \subseteq f_1 \subseteq g_1$. Similarly there is an automorphism g_2 of M_1^a extending f_1 hence f_0 . So $g_\ell \in \text{AUT}(M_\ell^a)$ for $\ell = 1, 2$ and $g_1 \upharpoonright M_0^a = f_0 = g_2 \upharpoonright M_0^a$. By the uniqueness of $NF_{\lambda, \langle \lambda, \lambda, \lambda \rangle}$ (i.e. Claim 8.9) there is an automorphism g_4 of M_4^a extending $g_1 \cup g_2$. This proves the desired conclusion. $\square_{8.13}$

8.14 Claim. *Assume*

- (a) $\bar{\delta} = \langle \delta_1, \delta_2, \delta_3 \rangle, \delta_\ell < \lambda^+$ is a limit ordinal for $\ell = 1, 2, 3$;
 $N_0 \leq_{\aleph} N_\ell \leq_{\aleph} N_3$ for $\ell = 1, 2$ and
- (b) N_ℓ is $(\lambda, \text{cf}(\delta_\ell))$ -saturated over N_0
- (c) N_3 is $\text{cf}(\delta_3)$ -saturated over $N_1 \cup N_2$.

Then $NF_\lambda(N_0, N_1, N_2, N_3)$ iff $NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$.

Proof. The “if” direction holds by 8.13. For the “only if” direction, by the proof of 8.12(1) (and Definition 8.2, 8.3) we can find $M_\ell (\ell \leq 3)$ such that $NF_{\lambda, \bar{\delta}}(M_0, M_1, M_2, M_3)$ and clauses (b), (c), (d) of Definition 8.3 holds so by 8.12 also $NF_\lambda(M_0, M_1, M_2, M_3)$. Easily there are for $\ell < 3$, an isomorphism f_ℓ from M_ℓ onto N_ℓ such that $f_0 = f_\ell \upharpoonright M_\ell$. By the uniqueness for smooth amalgamation (i.e. 8.13) we can find an isomorphism f_3 from M_3 onto N_3 extending $f_1 \cup f_2$. So as $NF_{\lambda, \bar{\delta}}(M_0, M_1, M_2, M_3)$ holds also $NF_{\lambda, \bar{\delta}}(f_0(M_0), f_3(M_1), f_3(M_2), f_3(M_3))$;
i.e. $NF_{\lambda, \bar{\delta}}(N_0, N_1, N_2, N_3)$ as required. $\square_{8.14}$

8.15 Claim. [*Monotonicity*]: If $NF_\lambda(N_0, N_1, N_2, N_3)$ and $N_0 \leq_{\aleph} N'_1 \leq_{\aleph} N_1$ and $N_0 \leq_{\aleph} N'_2 \leq_{\aleph} N_2$ and $N'_1 \cup N'_2 \subseteq N'_3 \leq_{\aleph} N_3$ then $NF_\lambda(N_0, N'_1, N'_2, N'_3)$.

Proof. Read Definition 8.3.

8.16 Claim. [*Symmetry*]: $NF_\lambda(N_0, N_1, N_2, N_3)$ holds if and only if $NF_\lambda(N_0, N_2, N_1, N_3)$ holds.

Proof. By Claim 8.10 (and Definition 8.3).

8.17 Claim. Assume $\alpha < \lambda^+$ is an ordinal and for $x \in \{a, b, c\}$ the sequence $\langle N_i^x : i \leq \alpha \rangle$ is a $\leq_{\aleph}$ -increasing sequence of members of K_λ , for $x = a, b$ the sequence is $\leq_{\aleph}$ -increasing continuous for $i \leq \alpha$, $N_i^b \cap N_\alpha^a = N_i^a$, $N_i^c \cap N_\alpha^a = N_i^a$, $N_i^a \leq_{\aleph} N_i^b \leq_{\aleph} N_i^c$ and N_i^c is (λ, κ_i) -saturated over N_i^b and $NF_{\lambda, \bar{\delta}^i}(N_i^a, N_{i+1}^a, N_i^c, N_{i+1}^b)$ (so $i < \alpha \Rightarrow N_i^c \leq_{\aleph} N_{i+1}^b$) where

$\bar{\delta}^i = \langle \delta_1^i, \delta_2^i, \delta_3^i \rangle$ sequence of limit ordinals, $i < \alpha \Rightarrow \delta_2^{i+1} = \delta_i^3$, and for $i < 0$ limit, $cf(\delta_i^3) = \sum_{j < i} \delta_j^3$, $\delta_1 = \sum_{\beta < \alpha} \delta_\beta^1$ and $\delta_3 = \kappa_\alpha$, $\bar{\delta} = \langle \delta_1, \delta_2^0, \delta_3 \rangle$. Then $NF_{\lambda, \bar{\delta}}(N_0^a, N_\alpha^a, N_0^b, N_\alpha^c)$.

Proof. Use uniqueness of 8.9, lastly use 8.9 to show N_α^b is $(\lambda, cf(\alpha))$ -saturated over $N_\alpha^a \cup N_0^b$. $\square_{8.17}$

8.18 Claim. Assume that $\alpha < \lambda^+$ and for $x \in \{a, b\}$ we have $\langle N_i^x : i \leq \alpha \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuously sequence of members of K_λ .

- 1) If $NF_\lambda(N_i^a, N_{i+1}^a, N_i^b, N_{i+1}^b)$ for each $i < \alpha$ then $NF_\lambda(N_0^a, N_\alpha^a, N_0^b, N_\alpha^b)$.
- 2) If $\alpha_1 < \lambda^+, \alpha_2 < \lambda^+$ and $M_{i,j}$ ($i \leq \alpha_1, j \leq \alpha_2$) are as in 8.8, and for each $i < \alpha_1, j < \alpha_2$ we have:

$$M_{i,j+1} \quad \bigcup_{M_{i,j}}^{M_{i+1,j+1}} \quad M_{i+1,j}$$

$$\text{then } M_{i,0} \quad \bigcup_{M_{0,0}}^{M_{\alpha_1, \alpha_2}} \quad M_{0,j} \text{ for } i \leq \alpha_1, j \leq \alpha_2.$$

Proof. 1) We first prove special cases and use them to prove more general cases.

Case A: N_{i+1}^a is (λ, δ_i^1) -saturated over N_i^a and N_{i+1}^b is (λ, δ_i^2) -saturated over $N_{i+1}^a \cup N_i^b$ for $i < \alpha$.

We can choose by induction on $i \leq \alpha$, $N_i^c \in K_\lambda$ such that

- (a) $N_0^b \leq_{\mathfrak{K}} N_0^c \leq_{\mathfrak{K}} N_{i+1}^b$, N_0^c is (λ, δ_0^2) -saturated over N_0^b , and $NF_{\lambda, \langle \delta_1^0, \delta_2^0, \delta_3^0 \rangle}(N_0^a, N_1^a, N_0^c, N_1^b)$
- (b) $N_\alpha^c \in K_\lambda$ is $(\lambda, \delta_3^\alpha)$ -saturated over N_α^b .

(Possible by uniqueness; i.e. 8.13 and monotonicity, i.e. 8.15). Now we can use 8.17.

Case B: For each $i < \alpha$ we have: N_{i+1}^a is (λ, κ_i) -saturated over N_i^a .

Let $\bar{\delta}^i = (\kappa_i, \lambda, \lambda)$.

We can find a $\leq_{\mathfrak{K}}$ -increasing sequence $\langle M_i^x : i \leq \alpha \rangle$ for $x \in \{a, b, c\}$, continuous for $x = a, b$ such that $i < \alpha \Rightarrow M_i^b \leq_{\mathfrak{K}} M_i^c \leq_{\mathfrak{K}} M_{i+1}^b$ and $M_\alpha^b \leq_{\mathfrak{K}} M_\alpha^c$ and $NF_{\lambda, \bar{\delta}^i}(M_i^a, M_{i+1}^a, M_i^c, M_{i+1}^b)$ by choosing M_i^a, M_i^b, M_i^c by induction on i . By Case A we know that $NF_\lambda(M_0^a, M_\alpha^a, M_0^b, M_\alpha^c)$ holds.

We can now choose an isomorphism f_0^a from N_0^a onto M_0^a (exists as K is categorical in λ) and then a $\leq_{\mathfrak{K}}$ -embedding of N_0^b into M_0^b extending f_0^a . Next we choose by induction on $i \leq \alpha$, f_i^a an isomorphism from N_i^a onto M_i^a such that: $j < i \Rightarrow f_j^a \subseteq f_i^a$, possible by “uniqueness of the (λ, κ_i) -saturated model over M_i^a ”. (0.29)

Now we choose by induction on $i \leq \alpha$, a $\leq_{\mathfrak{K}}$ -embedding f_i^b of N_i^b into M_i^b extending f_i^a and f_j^b for $j < i$. For $i = 0$ we have done it, for i limit use $\bigcup_{j < i} f_j^b$,

lastly for i a successor ordinal let $i = j + 1$, now we have

- (*)₂ $NF_\lambda(M_i^a, M_{i+1}^a, f_i^b(N_i^b), M_{i+1}^b)$
 [why? because $NF_{\lambda, \delta^i}(M_i^a, M_{i+1}^a, M_i^c, M_{i+1}^b)$ by the choice of the M_ζ^x 's hence by 8.14 we have $NF_\lambda(M_i^a, M_{i+1}^a, M_i^c, M_{i+1}^b)$ and as $M_i^a \leq_{\mathfrak{K}} f_i^b(N_i^b) \leq M_i^b, M_i^c$ by 8.15 we get (*)₂].

By (*)₂ and the uniqueness of smooth amalgamation 8.13 there is f_i^b as required. So without loss of generality f_α^b is the identity, so we have $N_0^a = M_0^a, N_\alpha^a = M_\alpha^a, N_0^b \leq_{\mathfrak{K}} M_0^b, N_\alpha^b \leq_{\mathfrak{K}} M_\alpha^b$; also as said above $NF_\lambda(M_0^a, M_\alpha^a, M_0^b, M_\alpha^b)$ holds so by monotonicity i.e. 8.15 we get $NF_\lambda(N_0^a, N_\alpha^a, N_0^b, N_\alpha^b)$ as required.

Case C: General case.

We can find M_i^ℓ for $\ell < 3, i \leq \alpha$ such that:

- (a) $M_i^\ell \in K_\lambda$
- (b) for each $\ell < 3, M_i^\ell$ is $\leq_{\mathfrak{K}}$ -increasing in i
- (c) $M_i^0 = N_i^a$
- (d) $M_{i+1}^{\ell+1}$ is (λ, λ) -saturated over $M_{i+1}^\ell \cup M_i^{\ell+1}$ for $\ell < 2, i < \alpha$
- (e) $NF_\lambda(M_i^\ell, M_{i+1}^\ell, M_i^{\ell+1}, M_{i+1}^{\ell+1})$ for $\ell < 2, i < \alpha$
- (f) $M_0^{\ell+1}$ is (λ, λ) -saturated over M_0^ℓ
- (g) for $\ell < 2$ and $i < \alpha$ limit we have

$$M_i^{\ell+1} \text{ is } (\lambda, \lambda)\text{-saturated over } \bigcup_{j < i} M_j^{\ell+1} \cup M_i^\ell$$

- (h) for $i < \alpha$ limit we have

$$NF_\lambda \left(\bigcup_{j < i} M_j', M_i^1, \bigcup_{j < i} M_j^2, M_i^2 \right).$$

[how? as in the proof of 8.8].

Now note:

- (*)₄ $M_i^{\ell+1}$ is $(\lambda, \text{cf}(\lambda \times (1+i)))$ -saturated over M_i^ℓ
 [why? If $i = 0$ by clause (f), if i a successor ordinal by clause (d) and if i is a limit ordinal then by clause (g)]
- (*)₅ for $i < \alpha, NF_\lambda(M_i^0, M_{i+1}^0, M_i^2, M_{i+1}^2)$
 [why? we use Case B for $\alpha = 2$ with $M_i^0, M_{i+1}^0, M_i^1, M_{i+1}^1, M_i^2, M_{i+1}^2$ here standing for $N_0^a, N_0^b, N_1^a, N_1^b, N_2^a, N_2^b$ there].

Now we continue as in Case B (using $f_i^a = \text{id}_{N_i^a}$ and defining by induction on i a $\leq_{\mathfrak{K}}$ -embedding f_i^b of N_i^b into M_i^c).

2) By part (1) for each i the sequences $\langle M_{\beta, i} : \beta \leq \alpha_1 \rangle, \langle M_{\beta, i+1} : \beta \leq \alpha_1 \rangle$ we get

$M_{\alpha_1} \bigcup_{M_{0,i}}^{M_{\alpha_1,i+1}} M_{0,i+1}$ hence by symmetry (i.e. 8.13) we have $M_{0,i+1} \bigcup_{M_{0,i}}^{M_{\alpha_1,i+1}} M_{\alpha_1,i}$.
 Applying part (1) to the sequences $\langle M_{0,j} : j \leq \alpha_2 \rangle, \langle M_{\alpha_1,j} : j \leq \alpha_2 \rangle$ we get
 $M_{0,\alpha_2} \bigcup_{M_{0,0}}^{M_{\alpha_1,\alpha_2}} M_{\alpha_1,0}$ hence by symmetry (i.e. 8.13) we have $M_{\alpha_1,0} \bigcup_{M_{0,0}}^{M_{\alpha_1,\alpha_2}} M_{0,\alpha_2}$; by
 monotonicity, i.e 8.15 (or restriction of the matrix) we get the desired conclusion.
 $\square_{8.18}$

8.19 Conclusion. Assume $\langle N_i^\ell : i \leq \alpha \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous for $\ell = 0, 1$ where $N_i^\ell \in K_\lambda$ and N_{i+1}^1 is (λ, κ_ℓ) -saturated over $N_{i+1}^0 \cup N_i^1$ and $\text{NF}_\lambda(N_i^0, N_i^1, N_{i+1}^0, N_{i+1}^1)$.

Then N_α^1 is $(\lambda, \text{cf}(\sum_{i < \alpha} \kappa_i))$ -saturated over $N_\alpha^0 \cup N_0^1$ (if α is a limit ordinal, “ N_{i+1}^1 is universal over $N_{i+1}^0 \cup N_i^1$ ” suffice).

Proof. The case α not limit is trivial so assume α is a limit ordinal. We choose by induction on $i \leq \alpha$, a sequence $\langle M'_{i,\varepsilon} : \varepsilon \leq \varepsilon(i) \rangle$ such that:

- (a) $\langle M_{i,\varepsilon} : \varepsilon \leq \varepsilon(i) \rangle$ is (strictly) $<_{\mathfrak{K}}$ -increasing continuous
- (b) $N_i^0 \leq_{\mathfrak{K}} M_{i,\varepsilon} \leq_{\mathfrak{K}} N_i^1$
- (c) $N_i^0 = M_{i,0}$
- (d) $\varepsilon(i)$ is (strictly) increasing continuous in i ,
- (e) $j < i$ & $\varepsilon \leq \varepsilon(j) \Rightarrow M_{i,\varepsilon} \cap N_j^1 = M_{j,\varepsilon}$
- (f) $\varepsilon(0) = 1, M_{i,1} = N_0^1$
- (g) for $i > 0, \lambda$ divide $\varepsilon(i)$
- (h) $N_\ell^i \leq_{\mathfrak{K}} M_{i+1,\varepsilon(i)+1}$.

If we succeed, then $\varepsilon(\alpha)$ is divisible by λ and $\langle M_{i,\varepsilon} : \varepsilon \leq \varepsilon(\alpha) \rangle$ is (strictly) $<_{\mathfrak{K}}$ -increasing continuous, $M_{\alpha,0} = N_\alpha^0$, and $M_{\alpha,\varepsilon(\alpha)} \leq_{\mathfrak{K}} N_\alpha^1$ but it includes N_i^1 for $i < \alpha$ hence (as α is a limit ordinal) it includes $\bigcup_{i < \alpha} N_i^1 = N_\alpha^1$; and by 7.5 we conclude that

$N_\alpha^1 = M_{\alpha,\varepsilon(\alpha)}$ is $(\lambda, \text{cf}(\alpha))$ -saturated over $M_{\alpha,1}$ hence over $N_\alpha^0 \cup N_0^1$ (both $< M_{\alpha,i}$).

For $i = 0$, and i limit there is not much to do. For i successor we use 8.20 below.

8.20 Conclusion. 1) If $\text{NF}_\lambda(N_0, N_1, N_2, N_3)$ and $\langle M_{0,\varepsilon} : \varepsilon \leq \varepsilon(*) \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous, $N_0 \leq_{\mathfrak{K}} M_{0,\varepsilon} \leq_{\mathfrak{K}} M_2$ then we can find $\langle M_{1,\varepsilon} : \varepsilon \leq \varepsilon(*) \rangle$ and N'_3 such that:

- (a) $N_3 \leq_{\mathfrak{K}} N'_3 \in K_\lambda$
- (b) $\langle M_{1,\varepsilon} : \varepsilon \leq \varepsilon(*) \rangle$ is $\leq_{\mathfrak{K}}$ -increasing continuous
- (c) $M_{1,\varepsilon} \cap N_2 = M_{0,\varepsilon}$
- (d) $N_1 \leq_{\mathfrak{K}} M_{1,\varepsilon} \leq_{\mathfrak{K}} N'_3$
- (e) if $M_{0,0} = N_0$ then $M_{1,0} = N$.

2) if N_3 is universal over $N_1 \cup N_2$, then without loss of generality $N_3 = N_2$.

Proof. 1) Straight by uniqueness.

2) Follows by (1).

$\square_{8.19}, \square_{8.20}$

§9 NICE EXTENSIONS IN K_{λ^+}

9.1 Hypothesis. Assume the hypothesis 8.1.

So by §8 we have reasonable control on smooth amalgamation in K_λ . We use this to define “nice” extensions in K_{λ^+} . This is treated again in §10.

9.2 Definition. 1) Let $M_0 \leq_{\lambda^+}^* M_1$ mean:

- (a) $M_\ell \in K_{\lambda^+}$, for $\ell = 0, 1$
- (b) we can find $\bar{M}^\ell = \langle M_i^\ell : i < \lambda^+ \rangle$, a representation of M^ℓ , so $M_i^\ell \in K_\lambda$
(and M_i^ℓ is $\leq_{\aleph}$ -increasing continuously and $M_\ell = \bigcup_{i < \lambda^+} M_i^\ell$) such that:

$$NF_\lambda(M_i^0, M_{i+1}^0, M_i^1, M_{i+1}^1) \text{ for } i < \lambda^+.$$

2) Let $M_0 <_{\lambda^+, \kappa}^+ M_1$ mean $M_0 \leq_{\lambda^+}^* M_1$ by some witnesses M_i^ℓ (for $i < \lambda^+, \ell < 2$) such that $NF_{\lambda, \langle \kappa, 1, \kappa \rangle}(M_i^0, M_{i+1}^0, M_i^1, M_{i+1}^1)$. If $\kappa = \lambda$, we omit it.

9.3 Claim. 1) If $M_0 \leq_{\lambda^+}^* M_1$ and $\bar{M}^\ell = \langle M_i^\ell : i < \lambda^+ \rangle$ is a representation of M_ℓ (as in 8.18) then for some club E of λ^+ , for every $\alpha < \beta$ from E we have $NF_\lambda(M_\alpha^0, M_\beta^0, M_\alpha^1, M_\beta^1)$.

2) Similarly for $<_{\lambda^+, \kappa}^+$; if $M_0 <_{\lambda^+, \kappa_\ell}^* M_1$, $\bar{M}^\ell = \langle M_i^\ell : i < \lambda^+ \rangle$ a representation of M_ℓ for $\ell = 1, 2$ then for some club E of λ^+ for every $\alpha < \beta$ from E we have $NF_{\lambda, \langle \text{cf}(\alpha), 1, \text{cf}(\alpha) \rangle}(M_\alpha^0, M_\beta^0, M_\alpha^1, M_\beta^1)$.

3) The κ in Definition 9.2(2) does not matter. In fact, if $\langle M_i^\ell : i < \lambda^+ \rangle$ are as in 9.2(1) then for some club E of λ^+ we have: $\alpha \in E \Rightarrow M_\alpha^1 \cap M_0 = M_\alpha^0$ and $\alpha < \beta \ \& \ \alpha \in E \ \& \ \beta \in E \Rightarrow [M_\beta^1 \text{ is } \text{cf}(\beta)\text{-saturated over } M_\beta^0 \cup M_\alpha^1]$.

Proof. 1) Straight by 8.18.

2) Easy.

3) By 8.19. (Could have used 7.8). □_{9.3}

9.4 Claim. 1) For every $\kappa = \text{cf}(\kappa) \leq \lambda$ and $M_0 \in K_{\lambda^+}$ for some $M_1 \in K_{\lambda^+}$ we have $M_0 <_{\lambda^+, \kappa}^+ M_1$.

2) $<_{\lambda^+}^*$ are $<_{\lambda^+, \kappa}^+$ are transitive.

3) If $M_0 \leq_{\aleph} M_1 \leq_{\aleph} M_2$ and $M_0 \leq_{\lambda^+}^* M_2$ then $M_0 \leq_{\lambda^+}^* M_1$.

4) [transitivity] If $M_0 \leq_{\lambda^+}^* M_1 <_{\lambda^+, \kappa}^+ M_2$ then $M_0 <_{\lambda^+, \kappa}^* M_2$.

Proof. 1) Let $\langle M_i^0 : i < \lambda^+ \rangle$ be a representation of M_0 such that M_{i+1}^0 is (λ, κ) -saturated over M_i^0 . We choose by induction on i , $M_i^1 \in K_\lambda$ such that $\langle M_i^1 : i < \lambda^+ \rangle$ is $<_{\aleph}$ -increasing continuously, $M_i^0 \leq_{\aleph} M_i^1$, $M_i^1 \cap M_0 = M_i^0$ and $NF_{\lambda, \langle \kappa, 1, \kappa \rangle}(M_i^0, M_{i+1}^0, M_i^1, M_{i+1}^1)$. We can do it by 7.7(4).

2) Concerning $<_{\lambda^+, \kappa}^+$ use 9.3 and 8.18 (i.e. transitivity for smooth amalgamations). Now the proof for $<_{\lambda^+}^*$ is similar.

3) By monotonicity for smooth amalgamations; i.e. 8.15.

4) Check. □_{9.4}

9.5 Claim. 1) If $M_0 \leq_{\lambda^+}^* M_\ell$ for $\ell = 1, 2, \kappa = \text{cf}(\kappa) \leq \lambda$ and $a \in M_2 \setminus M_0$ then for some M_3 and f we have: $M_1 <_{\lambda^+, \kappa}^+ M_3$ and f is an $\leq_{\mathfrak{R}}$ -embedding of M_2 into M_3 over M_0 with $f(a) \notin M_1$, moreover, $f(M_2) \leq_{\lambda^+}^* M_3$.
 2) [uniqueness] Assume $M_0 <_{\lambda^+, \kappa}^+ M_\ell$ for $\ell = 1, 2$ then there is an isomorphism f from M_1 onto M_2 over M_0 .

Proof. We first prove part (2).

2) By 9.3(1) + (2) there are representations $\bar{M}^\ell = \langle M_i^\ell : i < \lambda^+ \rangle$ of M_ℓ for $\ell < 3$ such that: $M_i^\ell \cap M_0 = M_0^\ell$ and $NF_{\lambda, \langle \kappa, \text{cf}(\kappa \times i), \kappa \rangle}(M_i^0, M_{i+1}^0, M_i^\ell, M_{i+1}^\ell)$.

Now we choose by induction on $i < \lambda$ an isomorphism f_i from M_i^1 onto M_i^2 , increasing with i and being the identity over M_i^0 . For $i = 0$ use “ M_0^ℓ is (λ, κ) -saturated over M_0^0 for $\ell = 1, 2$ ” which holds by 8.2. For i limit take unions, for i successor ordinal use uniqueness Claim 8.9.

Proof of part (1). Let $\kappa = \aleph_0$, by 9.4(1) there are for $\ell = 1, 2$ models $N_\ell^* \in K_{\lambda^+}$ such that $M_\ell <_{\lambda^+, \kappa}^+ N_\ell^*$. Now let $\bar{M}^\ell = \langle M_i^\ell : i < \lambda^+ \rangle$ be a representation of M_ℓ for $\ell = 0, 1, 2$ and let $\bar{N}^\ell = \langle N_i^\ell : i < \lambda^+ \rangle$ be a representation of N_ℓ^* for $\ell = 1, 2$. By 9.4(4) and 9.3(3) without loss of generality $NF_\lambda(M_i^0, M_{i+1}^0, M_i^\ell, M_{i+1}^\ell)$ for $\ell = 1, 2$ and $NF_{\lambda, \langle \kappa, 1, \kappa \rangle}(M_i^\ell, M_{i+1}^\ell, N_i^\ell, N_{i+1}^\ell)$ respectively. Now clearly N_0^ℓ is (λ, κ) -saturated over M_0^ℓ hence over M_0^0 (for $\ell = 1, 2$) so there is an isomorphism f_0 from N_0^2 onto N_0^1 extending $\text{id}_{M_0^0}$ and $f(a) \notin M_0^1$.

We continue as in the proof of part (2). In the end $f = \bigcup_{i < \lambda^+} f_i$ is an isomorphism of N_2 onto N_1 over M_0 and as $f_0^1(a)$ is well defined and in $N_0^1 \setminus M_0^1$ clearly $f(a) = f_0(a) \notin M_1$, as required. $\square_{9.5}$

9.6 Claim. If δ is a limit ordinal $< \lambda^+$ and $\langle M_i : i < \delta \rangle$ is a $\leq_{\lambda^+}^*$ -increasing continuous then $M_i \leq_{\lambda^+}^* \bigcup_{j < \delta} M_j$ for each $i < \delta$.

Proof. We prove it by induction on δ . Now if C is a club of δ with $i \in C$, then we can replace $\langle M_j : j < \delta \rangle$ by $\langle M_j : j \in C \rangle$ so without loss of generality $\delta = \text{cf}(\delta)$, so $\delta \leq \lambda^+$; clearly it is enough to prove $M_0 \leq_{\lambda^+}^* \bigcup_{j < \delta} M_j$. Let $\langle M_\zeta^i : \zeta < \lambda^+ \rangle$ be a representation of M_i .

Case A: $\delta < \lambda^+$.

Without loss of generality (see 9.3(1)) for every $i < j < \delta$ and $\zeta < \lambda^+$ we have:

$M_\zeta^j \cap M_i = M_\zeta^i$ and $NF_\lambda(M_\zeta^i, M_{\zeta+1}^i, M_\zeta^j, M_{\zeta+1}^j)$. Let $M_\zeta^\delta = \bigcup_{i < \delta} M_\zeta^i$, so

$\langle M_\zeta^\delta : \zeta < \lambda^+ \rangle$ is $\leq_{\mathfrak{R}}$ -increasingly continuous sequence of members of K_λ with limit M_δ , and for $i < \delta, M_\zeta^\delta \cap M_i = M_\zeta^i$. By symmetry (see 8.16) we have $NF_\lambda(M_\zeta^i, M_\zeta^{i+1}, M_{\zeta+1}^i, M_{\zeta+1}^{i+1})$ so as $\langle M_\zeta^i : i \leq \delta \rangle, \langle M_{\zeta+1}^i : i \leq \delta \rangle$ are $\leq_{\mathfrak{R}}$ -increasingly continuous by 8.18 we know $NF_\lambda(M_\zeta^0, M_\zeta^\delta, M_{\zeta+1}^0, M_{\zeta+1}^\delta)$ hence by symmetry (8.16)

we have $NF_\lambda(M_\zeta^0, M_{\zeta+1}^0, M_\zeta^\delta, M_{\zeta+1}^\delta)$.

So $\langle M_\zeta^0 : \zeta < \lambda^+ \rangle, \langle M_\zeta^\delta : \zeta < \lambda^+ \rangle$ are witnesses to $M_0 \leq_{\lambda^+}^* M_\delta$.

Case B: $\delta = \lambda^+$.

By 9.3(1) (using normality of the club filter, restricting to a club of λ^+ and renaming), without loss of generality for $i < j \leq 1 + \zeta < 1 + \xi < \lambda^+$ we have $M_\zeta^j \cap M_i = M_\zeta^i$, and $NF_\lambda(M_\zeta^i, M_\xi^i, M_\zeta^j, M_\xi^j)$. Let us define $M_i^{\lambda^+} = \bigcup_{j < 1+i} M_i^j$. So

$\langle M_i^{\lambda^+} : i < \lambda^+ \rangle$ is a representation of $M_{\lambda^+}^\lambda = M_\delta$ and continue as before. $\square_{9.6}$

9.7 Claim. Assume $M_0 <_{\lambda^+, \kappa}^+ M_2$ and $a \in M_2 \setminus M_0$, and for some $N \leq_{\mathfrak{K}} M_0$ we have: $N \in K_\lambda$ and $tp(a, N, M_2)$ is minimal. Then we can find M_1 , $M^0 = \langle M_{0,i} : i < \lambda^+ \rangle, M^1 = \langle M_{1,i} : i < \lambda^+ \rangle$ such that:

- (a) $\bar{M}^0$ is a $\leq_{\mathfrak{K}}$ -representation of M_0
- (b) $\bar{M}^1$ is a representation of $M_1 (\in K_{\lambda^+})$, $a \in M_{1,i}$, for all i
- (c) $M_0 \leq_{\mathfrak{K}} M_1 \leq_{\mathfrak{K}} M_2$
- (d) for $i < \lambda^+$ we have $NF_{\lambda, \langle \lambda, 1, 1 \rangle}(M_{0,i}, M_{0,i+1}, M_{1,i}, M_{1,i+1})$
(hence $M_{\ell,i} = M_{\ell+1,i} \cap M_\ell$)
- (e) $(M_{0,i}, M_{1,i}, a) \in K_\lambda^3$ is minimal and reduced.

Proof. Let $\langle M_{0,i} : i < \lambda^+ \rangle, \langle M_{2,i} : i < \lambda^+ \rangle$ be representations of M_0, M_2 respectively, as required in 9.2(2) and without loss of generality $N \leq_{\mathfrak{K}} M_{0,0}$ and $a \in M_{2,0}$. We now choose by induction on $\zeta < \lambda^+$, an ordinal $i(\zeta)$ and models $M_{1,i(\zeta)}, M_{3,i(\zeta)}$ such that:

- (A) $i(\zeta) < \lambda^+$ is increasing continuous in ζ and $a \in M_{2,i(0)} \setminus M_{0,i(0)}, N \leq_{\mathfrak{K}} M_{0,i(0)}$
- (B) $M_{0,i(\zeta)} \leq_{\mathfrak{K}} M_{1,i(\zeta)} \leq_{\mathfrak{K}} M_{3,i(\zeta)}$ and $M_{2,i(\zeta)} \leq_{\mathfrak{K}} M_{3,i(\zeta)}$
- (C) $a \in M_{1,i(0)}$ and $(M_{0,i(\zeta)}, M_{1,i(\zeta)}, a)$ is minimal and reduced
- (D) for $\xi < \zeta$ and $(\ell, m) \in \{(0, 1), (0, 2), (1, 3), (2, 3)\}$ we have $NF_\lambda(M_{\ell,i(\xi)}, M_{\ell,i(\zeta)}, M_{m,i(\xi)}, M_{m,i(\zeta)})$
- (E) $M_{1,i(\zeta)}, M_{3,i(\zeta)}$ is $\leq_{\mathfrak{K}}$ -increasing continuous in ζ .

For $\zeta = 0$ note that for $i(0) < \lambda^+, a \in M_{2,i(0)}$ and $M_{2,i(0)}$ is universal over $M_{0,i(0)}$.

For ζ limit let $i(\zeta) = \bigcup_{\xi < \zeta} i(\xi)$ and $M_{1,i(\zeta)} = \bigcup_{\xi < \zeta} M_{1,i(\xi)}$.

For $\zeta = \xi + 1$, there is $i(\zeta) \in (i(\xi), \lambda^+)$ and a model N_ζ such that $M_{1,i(\xi)} \leq_{\mathfrak{K}} N_\zeta \in K_\lambda$ and $\leq_{\mathfrak{K}}$ -embedding f of $M_{0,i(\zeta)}$ into $N_\zeta, f \upharpoonright M_{0,i(\zeta)}$ the identity and $(f(M_{0,i(\xi)}), N_\zeta, a)$ is minimal and reduced. By uniqueness (i.e. Claim 8.2) we can find such N satisfying $(\exists M)(N \leq_{\mathfrak{K}} M \in K_\lambda \ \& \ M_{1,i(\zeta)} \leq_{\mathfrak{K}} M)$. So we can carry the induction.

Lastly, by uniqueness of $<_{\lambda^+, \kappa}^+$ we can make $M_3 = \bigcup_{\zeta < \lambda^+} M_{3,i(\zeta)}$ to be $\leq_{\mathfrak{K}} M_2$ as required. $\square_{9.7}$

9.8 Definition. If (M_0, M_1, a) are as in 9.7 (a)-(e) we say (M_0, M_1, a) is λ^+ -locally reduced nice and minimal (λ^+ -l.r.n.m.). We omit “nice” if we omit clause (d).

9.9 Claim. *If (M_0, M_1, a) is λ^+ -l.r.n.m. then $(M_0, M_1, a) \in K_{\lambda^+}^3$ is reduced.*

Proof. Check.

Using also 7.5:

9.10 Claim. $M_0 <_{\lambda^+, \kappa}^+ M_1$ if and only if we can find $\langle M_j^*, a_j : j < \lambda^+ \times \kappa \rangle$ such that:

- (a) M_j^* is $\leq_{\mathfrak{R}}$ -increasing continuous (in K_{λ^+})
- (b) (M_j^*, M_{j+1}^*, a_j) is λ^+ -l.r.n.m.
- (c) $M_0^* = M_0, M_j^* \cup M_j^* = M_1$
- (d) for some $N \leq M_0, N \in K_{\lambda}$ and minimal reduced $p \in S(N)$, for every j .

Proof. We can find $\langle M_j^* : j \leq \lambda^+ \times \kappa \rangle$ satisfying clauses (a), (b) and (d). By 5.6 easily $M_0^* <_{\lambda}^+ M_{\lambda^+}^*$. Now by the uniqueness (= 9.5(2)) + categoricity of K in λ^+ , we are done. $\square_{9.10}$

9.11 Claim. *In $(K_{\lambda^+}, <_{\lambda^+}^*)$ we have disjoint amalgamation.*

Proof. First redo 9.5 assuming $(M_0, M_{\ell}, a_{\ell})$ for $\ell = 1, 2$ is λ^+ -l.r.n.m., and getting $a_1 \notin f(M_2), f(a_2) \notin M_1$ (and we can start with this). By 9.9 we get $M_1 \cap f(M_2) = M_3$, so we have disjoint amalgamation. By 9.10 and chasing arrows we get it in general. $\square_{9.11}$

Remark. This is like the proof of disjoint amalgamations in 5.11.

§10 NON-STRUCTURE FOR $\leq_{\lambda^+}^*$

10.1 Hypothesis. Assume the hypothesis 8.1 and the further model theoretic properties deduced since then (including 6.7 but not 6.12.)

It would have been nice to prove all disjoint amalgamations in K_λ are NF_λ , but this is, at this point, not clear. But as we look upward (i.e. want to prove statement on $K_{>\lambda^+}$) and $<_{\lambda^+}^*$ is very nice, it will be essentially just as well if for $M, N \in K_{\lambda^+}$ we have $M \leq_{\aleph} N \Rightarrow M \leq_{\lambda^+}^* N$. Our intention is to assume $M^* \leq_{\aleph} N^*$ is a counterexample of this statement and we would like to say that in a sense this implies the existence of many types over M^* so that we can construct many models in λ^{+2} . Note: building models in $K_{\lambda^+}, K_{\lambda^{++}}$ by approximations in K_λ is nice if we use the smooth amalgamation but we do not have it for non-smooth ones. So we shall use $M^* \in K_{\lambda^+}$ being saturated so it has many automorphisms.

10.2 Claim. 1) Assume $M_1 \leq_{\aleph} M_2$ are in K_{λ^+} . Then we can find $M_0 \in K_{\lambda^+}$ such that $M_0 <_{\lambda, \kappa}^+ M_1$ and $M_0 \leq_{\lambda^+}^* M_2$.

2) Also we can find $\langle M_{0,i} : i < \lambda^+ \rangle$, an $\leq_{\aleph}$ -increasingly continuous sequence of members of K_{λ^+} such that $M_{0,i} <_{\lambda^+}^* M_{0,i+1}$ and $\bigcup_{i < \lambda^+} M_{0,i} = M_1$ and

$i < \lambda^+ \Rightarrow M_{0,i} \leq_{\lambda^+}^* M_2$.

Proof. Let $(M^*, N^*) \in K_\lambda^{3, \text{uq}}$ be from 8.1(2). Let $\langle M_{\ell,i} : i < \lambda^+ \rangle$ be a representation of M_ℓ for $\ell = 1, 2$ and without loss of generality $M_{\ell,i+1}$ is (λ, λ) -saturated over $M_{\ell,i}$ for $\ell = 1, 2$ and $M_{2,i} \cap M_1 = M_{1,i}$.

1) Now choose by induction on $i, M_{0,i}$ such that:

- (a) $M_{0,i} \leq_{\aleph} M_{1,i}$
- (b) $M_{0,i}$ is $\leq_{\aleph}$ -increasing continuous
- (c) $M_{0,i+1} \cap M_{1,i} = M_{0,i}$
- (d) $M_{1,i+1}$ is $(\lambda, \text{cf}(\lambda))$ -saturated over $M_{1,i} \cup M_{0,i+1}$
- (e) $(M_{0,i}, M_{0,i+1}) \cong (M^*, N^*)$.

There is no problem to carry the definition. Now let $M_0 = \bigcup_{i < \lambda^+} M_{0,i}$ so $M_0 <_{\lambda^+}^+ M_1$

and $M_0 \leq_{\lambda^+}^* M_2$ are checked by their definitions.

2) We choose by induction on $i < \lambda^+, \langle M_{\varepsilon,i}^* : \varepsilon \leq 1+i \rangle$ such that:

- (a) $M_{1+i,i}^* = M_{1, \lambda \times (1+\varepsilon) \times i}$
- (b) For each ε the sequence $\langle M_{\varepsilon,j}^* : \varepsilon \leq j \leq i \rangle$ is $\leq_{\aleph}$ -increasing continuous
- (c) For each i , the sequence $\langle M_{\varepsilon,i}^* : \varepsilon \leq 1+i \rangle$ is $\leq_{\aleph}$ -increasing continuous
- (d) $M_{\varepsilon,i}^* \cap M_{\zeta,j}^* = M_{\min\{\varepsilon, \zeta\}, \min\{i, j\}}^*$
- (e) $M_{\varepsilon+1,i+1}^*$ is $(\lambda, \text{cf}(\lambda \times (1+\varepsilon)))$ -saturated over $M_{\varepsilon,i+1}^* \cup M_{\varepsilon+1,i}^*$
- (f) $NF_{\lambda, \langle \lambda, 1, \lambda \rangle}(M_{\varepsilon,i}^*, M_{\varepsilon+1,i}^*, M_{\varepsilon,i+1}^*, M_{\varepsilon+1,i+1}^*)$
- (g) for $\varepsilon < 1+i$ we have $NF_\lambda(M_{\varepsilon,i}^*, M_{\varepsilon,i+1}^*, M_{2, \lambda \times \lambda \times i}, M_{2, \lambda \times \lambda \times (i+1)})$.

For $i = 0$, i limit no problem, for $i = j + 1$ first choose $N_{i,\varepsilon,\zeta} \leq_{\mathfrak{K}} M_{1,\lambda \times \lambda \times i + \lambda \times \zeta}$ for $\zeta \leq \lambda \times (1 + i)$, $\leq_{\mathfrak{K}}$ -increasing continuous, $N_{i,\varepsilon,\zeta} \in K_\lambda$, $N_{i,\varepsilon,0} = M_{1,j} (= M_{1+i,i}^*)$, $(N_{i,\varepsilon,\zeta}, N_{i,\varepsilon,\zeta+1}) \cong (M^*, N^*)$ and $N_{i,\varepsilon,\zeta+1} \cap M_{2,\lambda \times \lambda \times i + \lambda \times \zeta} = N_{i,\varepsilon,\zeta}$.

Now by 5.6, without loss of generality $N_{i,\varepsilon,\lambda}$ is (λ, λ) -saturated over $M_{1,j}$, and we choose it as $M_{1+j,i}^*$, and we choose $M_{1,i+1}^*$ as $M_{1+i,i}$ note that clauses (a) and (f) holds. Now we can find N_{ε,j_1} for $j_1 < 1 + j$ as in 8.8 and use uniqueness of the $(\lambda, \lambda \times (1 + i))$ -saturated model over $M_{1,j}$. $\square_{10.3}$

10.3 Conclusion. Assume $M \leq_{\mathfrak{K}} N$ are from K_{λ^+} . If $\langle M_i : i < \lambda^+ \rangle$ is $\leq_{\lambda^+}^*$ -increasing continuous and for each i for some N_i we have $M_i <_{\lambda^+}^+ N_i \leq_{\lambda^+}^* M_{i+1}$ then for some (M', N') we have:

$$(M', N') \cong (M, N)$$

$$M' = \bigcup_{i < \lambda} M_i$$

$$i < \lambda^+ \Rightarrow M_i \leq_{\lambda^+}^* N'.$$

Proof. By 10.2(2) and as we know $M^a \leq_{\lambda^+}^* M^b \leq_{\lambda^+}^+ M^c \Rightarrow M^a <_{\lambda^+}^+ M^c$ and the uniqueness of M'' over M' when $M' <_{\lambda^+}^+ M''$. $\square_{10.3}$

10.4 Lemma. If $\leq_{\mathfrak{K}} \restriction K_{\lambda^+}$ is not $\leq_{\lambda^+}^*$ then $I(\lambda^{+2}, K) = 2^{\lambda^{+2}}$.

Proof. Let $S \subseteq \{\delta < \lambda^{+2} : \text{cf}(\delta) = \lambda^+\}$ be stationary. We shall below construct a model $M_S \in K_{\lambda^{+2}}$ such that from the isomorphism type of M_1^S we can reconstruct $S/\mathcal{D}_{\lambda^{+2}}$; this clearly suffices. Choose $M \leq_{\mathfrak{K}} N$ from K_{λ^+} such that $\neg(M \leq_{\lambda^+}^* N)$ so by 9.4(3) without loss of generality $|N \setminus M| = \lambda^+$.

We choose by induction on $\alpha < \lambda^{+2}$ a model M_α^S such that:

- (a) $M_\alpha^S \in K_{\lambda^+}$ has universe $\lambda \times (1 + \alpha)$
- (b) for $\beta < \alpha$ we have $M_\beta^S \leq_{\mathfrak{K}} M_\alpha^S$
- (c) if $\alpha = \beta + 1$, $\beta \notin S$ then $M_\beta^S <_{\lambda^+}^+ M_\alpha^S$
- (d) if $\alpha = \beta + 1$, $\beta \in S$ then $(M_\beta^S, M_\alpha^S) \cong (M, N)$
- (e) if $\beta < \alpha$, $\beta \notin S$ then $M_\beta^S \leq_{\lambda^+}^* M_\alpha^S$.

We use freely the transitivity (9.4(4)) and continuity (9.6) of $\leq_\lambda^*$ and $[M^a \leq_{\mathfrak{K}} M^b \leq_{\mathfrak{K}} M^c \text{ in } K_\lambda, \neg(M^a \leq_{\lambda^+}^* M^b \Rightarrow \neg(M^a \leq_{\lambda^+}^* M^c)]$ 9.4(3).

For $\alpha = 0$ no problem.

For α limit no problem.

For $\alpha = \beta + 1$, $\beta \notin S$ no problem.

For $\alpha = \beta + 1$, $\beta \in S$ so $\text{cf}(\beta) = \lambda^+$, let $\langle \gamma_i : i < \lambda^+ \rangle$ be increasing continuous with limit β and $\text{cf}(\gamma_i) \leq \lambda^+$, hence $\gamma_i \notin S$. By 9.3(2) without loss of generality $M_{\gamma_i} <_{\lambda^+}^* M_{\gamma_{i+1}}$. Now use 10.3 (and the uniqueness (9.5(2))). $\square_{10.4}$

10.5 Conclusion. Assume $I(\lambda^{+2}, K) < 2^{\lambda^{+2}}$ (in addition to hypothesis 10.1).

1) $\leq_{\lambda^+}^* = \leq_{\mathfrak{K}} \restriction K_{\lambda^+}$.

2) $(K_{\lambda^+}, \leq_{\mathfrak{K}})$, has disjoint amalgamation so no $M \in K_{\lambda^{+2}}$ is $\leq_{\mathfrak{K}}$ -maximal.

3) $K_{\lambda^{+3}} \neq \emptyset$.

Proof. 1) By 10.4.

2) By 9.11 (and part (1)).

3) By 10.5(2) and 2.6(8) with λ there replaced by λ^+ here. $\square_{10.5}$

* * *

So we have finally proved the main theorem. Though not directly contributing to our main theme, we remark some more consequences of $\leq_{\mathfrak{K}} \restriction K_{\lambda^+} \neq \leq_{\lambda^+}^*$.

10.6 Claim. $(*)_0 \Leftrightarrow (*)_1$ where

$(*)_0$ for some $M \leq_{\mathfrak{K}} N$ from K_{λ^+} , we do not have $M \leq_{\lambda^+}^* N$

$(*)_1$ for some $M \leq_{\mathfrak{K}} N$ from K_{λ^+} we have:

if $a \in N \setminus M$ then $tp(a, M, N)$ is not realized in any M
such that $M^* \leq_{\lambda^+}^* M \in K_{\lambda^+}$

10.7 Definition. Assume $M_0 <_{\mathfrak{K}} M_1$ are in K_{λ^+} , and $\bar{M}^\ell = \langle M_{\ell, i} : i < \lambda^+ \rangle$ is a $\leq_{\mathfrak{K}}$ -representation of M_ℓ for $\ell = 0, 1$. Let

$$(a) \quad \mathcal{S}_0(\bar{M}^0, \bar{M}^1) = \left\{ \delta < \lambda^+ : M_{1, \delta} \cap M_0 = M_{0, \delta} \right. \\ \left. \text{and not } NF_\lambda(M_{0, \delta}, M_{0, \delta+1}, M_{1, \delta}, M_1) \right\}$$

$$(b) \quad \mathcal{S}_1(M_0, M_1) = \mathcal{S}_0(\bar{M}^0, \bar{M}^1) / D_{\lambda^+} \text{ (well defined).}$$

$$(c) \quad J \text{ is the normal ideal on } \lambda^{++} \text{ generated by sets of the form } \mathcal{S}_0(M^0, M^1) \text{ where } M^0, M^1 \text{ are as above.}$$

10.8 Comment: An earlier try for 10.4 was:

1) For every $S \in J$ for some $\bar{M}^0, M^1$ as in 10.7 we have $S = \mathcal{S}_0(\bar{M}^0, \bar{M}^1)$.

2) If $S_1 = S_1(M^0, M^1)$ is stationary then for some $\bar{M} = \langle M_i : i < \lambda^+ \rangle$ a representation of $M = \bigcup_{i < \lambda^+} M_i \in K_{\lambda^+}$ for every $S \subseteq S_1$ for some $\bar{M}'$, we have

$$\bar{M}^1 = \langle M_i^1 : i < \lambda^+ \rangle, M_1 <_{\mathfrak{K}} M^1 = \bigcup_{i < \lambda^+} M_i^1, M_i^1 \cap M = M_i \text{ and}$$

$$S_0(\bar{M}, \bar{M}^1) = S \text{ mod } \mathcal{D}_{\lambda^+}.$$

3) If $\leq_{\lambda^+}^* \neq \leq_{\aleph} \restriction K_{\lambda^+}$ and $I(\lambda^{++}, K) < 2^{\lambda^{++}}$ then for some stationary $S \subseteq \lambda^+$ we have:

- (a) $\mathcal{D}_{\lambda^+} \restriction S$ is λ^{++} -saturated
- (b) $\bar{M}^0, \bar{M}^1$ as in 10.4 implies $S_1(\bar{M}^0, \bar{M}^1) \subseteq S \bmod D_{\lambda^+}$.

4) If $\leq_{\lambda^+}^* \neq \leq_{\aleph} \restriction K_{\lambda^+}$ and $2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}}$ then $I(\lambda^{++}, K) = 2^{\lambda^{++}}$.

Proof. 1) First we prove only “ $S \subseteq S_0(\bar{M}^0, \bar{M}^1)$ ”. Easy as $\aleph_{\lambda^+}$ has amalgamation and

$$\bigotimes \text{ if } M_0 \leq_{\aleph} M_1 \leq_{\aleph} M_2 \text{ are in } K_{\lambda^+}, \bar{M}^\ell \\ \text{representing the } S_0(\bar{M}^0, \bar{M}^1) \subseteq S_0(\bar{M}^0, \bar{M}^2).$$

Now for equality use part (2).

2) Similar to the proof of 10.2.

3) Suppose $S^* = S_1(\bar{M}^0, \bar{M}^2)$ is stationary, let $\bar{S} = \langle S'_\alpha : \alpha < \lambda^{++} \rangle$ be such that $S'_\alpha \in J$. We can build a model $M^{\bar{S}} \in K_{\lambda^{++}}$ and a representation $\langle M_\alpha^{\bar{S}} : \alpha < \lambda^{++} \rangle$ such that

$$S_1(M_\alpha, M_{\alpha+1}) = S'_\alpha / D_{\lambda^{++}}.$$

4) By §3 (using the proof of part (2)).

REFERENCES.

- [DvSh 65] Keith J. Devlin and Saharon Shelah. A weak version of $\diamond$ which follows from $2^{\aleph_0} < 2^{\aleph_1}$. *Israel Journal of Mathematics*, **29**:239–247, 1978.
- [GrSh 266] Rami Grossberg and Saharon Shelah. $***$. . Does not exist.
- [Ha61] Andras Hajnal. Proof of a conjecture of S.Ruziewicz. *Fundamenta Mathematicae*, **50**:123–128, 1961.
- [HaSh 323] Bradd Hart and Saharon Shelah. Categoricity over P for first order T or categoricity for $\phi \in L_{\omega_1\omega}$ can stop at $\aleph_k$ while holding for $\aleph_0, \dots, \aleph_{k-1}$. *Israel Journal of Mathematics*, **70**:219–235, 1990.
- [KlSh 362] Oren Kolman and Saharon Shelah. Categoricity of Theories in $L_{\kappa,\omega}$, when κ is a measurable cardinal. Part 1. *Fundamenta Mathematicae*, **151**:209–240, 1996.
- [MaSh 285] Michael Makkai and Saharon Shelah. Categoricity of theories in $L_{\kappa\omega}$, with κ a compact cardinal. *Annals of Pure and Applied Logic*, **47**:41–97, 1990.
- [Mw85] Johann A. Makowsky. Compactnes, embeddings and definability. In J. Barwise and S. Feferman, editors, *Model-Theoretic Logics*, pages 645–716. Springer-Verlag, 1985.
- [MkSh 313] Alan H. Mekler and Saharon Shelah. Diamond and λ -systems. *Fundamenta Mathematicae*, **131**:45–51, 1988.
- [Sh 600] Saharon Shelah. Categoricity in abstract elementary classes: going up inductive step. *in preparation*.
- [Sh 394] Saharon Shelah. Categoricity of abstract classes with amalgamation. *Annals of Pure and Applied Logic*, **accepted**.
- [Sh 472] Saharon Shelah. Categoricity of Theories in $L_{\kappa^*\omega}$, when κ^* is a measurable cardinal. Part II. *Fundamenta Mathematicae*, **submitted**.
- [Sh 638] Saharon Shelah. More on Weak Diamond. *in preparation*.
- [Sh:e] Saharon Shelah. *Non-structure theory*, accepted. Oxford University Press.
- [Sh 460] Saharon Shelah. The Generalized Continuum Hypothesis revisited. *Israel Journal of Mathematics*, **accepted**.
- [Sh:b] Saharon Shelah. *Proper forcing*, volume 940 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, xxix+496 pp, 1982.
- [Sh 87b] Saharon Shelah. Classification theory for nonelementary classes, I. The number of uncountable models of $\psi \in L_{\omega_1,\omega}$. Part B. *Israel Journal of Mathematics*, **46**:241–273, 1983.
- [Sh 87a] Saharon Shelah. Classification theory for nonelementary classes, I. The number of uncountable models of $\psi \in L_{\omega_1,\omega}$. Part A. *Israel Journal of Mathematics*, **46**:212–240, 1983.

- [Sh 88] Saharon Shelah. Classification of nonelementary classes. II. Abstract elementary classes. In *Classification theory (Chicago, IL, 1985)*, volume 1292 of *Lecture Notes in Mathematics*, pages 419–497. Springer, Berlin, 1987. Proceedings of the USA–Israel Conference on Classification Theory, Chicago, December 1985; ed. Baldwin, J.T.
- [Sh 300] Saharon Shelah. Universal classes. In *Classification theory (Chicago, IL, 1985)*, volume 1292 of *Lecture Notes in Mathematics*, pages 264–418. Springer, Berlin, 1987. Proceedings of the USA–Israel Conference on Classification Theory, Chicago, December 1985; ed. Baldwin, J.T.
- [Sh:c] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, xxxiv+705 pp, 1990.
- [Sh 351] Saharon Shelah. Reflecting stationary sets and successors of singular cardinals. *Archive for Mathematical Logic*, **31**:25–53, 1991.
- [Sh:f] Saharon Shelah. *Proper and improper forcing*. Perspectives in Mathematical Logic. Springer, 1998.